

CULTIVOS
HERBACEOS

17th INTERNATIONAL SUNFLOWER CONFERENCE

Vol. 2



Proceedings of the
**17th International
Sunflower Conference**

Vol. 2



Córdoba, Spain
June 8-12, 2008

Sponsored by The International Sunflower Association, Paris, France



Instituto de Investigación y Formación Agraria y Pesquera
CONSEJERÍA DE AGRICULTURA Y PESCA



International Sunflower
Association

Proceedings of the 17th International Sunflower Conference
Córdoba, Spain. June 8-12, 2008

Editor: Leonardo Velasco

Editorial Committee:

José M. Fernández-Martínez

Luís García-Torres

Pedro González

José M. Melero-Vara

Francisco Orgaz

Begoña Pérez-Vich

Francisco Villalobos

Organizing Committee:

Chairman: Juan Domínguez: Institute for Research and Training in Agriculture and Fisheries (IFAPA).

Deputy Chairman: Juan Fernández Pérez LIMAGRAIN IBERICA

ISA Secretariat: Laurecine Lot

ISA Secretary General: André Pouzet

Luis Carlos Alonso Arnedo. KOIPESOL S.A.

José M. Fernández Martínez. Institute for Sustainable Agriculture (CSIC)

José Antonio García de Tejada. ARLESA Semillas

José Rafael García Ruiz. Institute for Research and Training in Agriculture and Fisheries (IFAPA)

Luis López Bellido. University of Cordoba

José M. Melero Vara. Institute for Sustainable Agriculture (CSIC)

Leonardo Velasco. Institute for Sustainable Agriculture (CSIC)

Photography: *Feral sunflowers growing in a sunflower field in Santa Cruz, Córdoba, Spain.* Courtesy of Marie-Hélène Muller.

Foreword

The proceedings of the 17th *International Sunflower Conference* contain 142 contributions from scientists of 24 countries. They include plenary lectures in several disciplines and regular communications presented in posters during the conference and discussed in the corresponding workshops. The manuscripts are classified by disciplines. They offer a good picture of the current state of the art of sunflower research and cultivation around the world.

The manuscripts in the *Proceedings* have been reviewed by an editorial committee with the main objective of helping the authors to improve their manuscripts through a critical reading. The authors received the edited manuscripts together with the comments of the reviewers and then went on to draft their final version. All the manuscripts received have been published in the *Proceedings*. The contents of the manuscripts are the responsibility of the authors. They should be considered as being privileged communications that require the express consent of the authors to be reprinted in part or as a whole. We wish to thank both the members of the Editorial Committee for their dedication to the task of editing such a large number of manuscripts, as well as all the authors for their collaboration throughout the whole edition process.

The Organizing Committee would also like to thank Diana Badder and José A. Palacios for their excellent editorial assistance in the preparation of these *Proceedings*. We are indebted to the Spanish Association of Sunflower Breeders (Asociación Española de Mejoradores de Girasol), which collaborated actively in the organization of the conference, and, very especially, to Juan Parejo, who was in charge of the financial side.

Finally, we would like to thank all the participants in the conference, who have contributed to its success by a careful preparation and revision of manuscripts and posters, presentation of their research in the workshops, and stimulating discussions throughout the conference on the scientific and technical aspects of sunflower research and cultivation in the world.

The Organizing Committee
17th International Sunflower Conference
Córdoba, Spain. June 8-12, 2008

Volume 2

Table of Contents

CROP PRODUCTION – PHYSIOLOGY

Early responses to high crop population density in sunflower: Controls and effects of the crop self-organization process	
Mónica López Pereira, Nora Trápani, Jorge Casal, Antonio Hall.....	417
Early sowing as a means of drought escape in sunflower: effects on vegetative and reproductive stages	
Clémentine Allinne, Nedjoua Ghoribi, Pierre Maury, Rhym Maougal, Ahmad Sarrafi, Nadia Ykhlef, Philippe Grieu	423
SUNFLO: A joint phenotyping and modelling approach to analyse and predict the differences in yield potential of sunflower genotypes	
Jérémie Lecoeur, Richard Poiré-Lassus, Angélique Christophe, Lydie Guilioni.....	429
Root system and water extraction variability for sunflower hybrids	
Lydie Guilioni, Ando M. Radanielson, Angélique Christophe, Jérémie Lecoeur.....	435
Effects of high water table conditions on sunflower growth and quality	
Satoko Yasumoto, Yukari Terakado, Morio Matsuzaki, Kensuke Okada.....	441

CROP PRODUCTION – FERTILIZER

Optimizing of potassium and magnesium fertilizers in sunflower production	
E. Sepehr, F. Nourgolipour, M.J. Malakouti.....	447
Sunflower response to mineral nitrogen, organic and bio-fertilizers under two different levels of salinity	
M.M. Keshta, T.Y. Rizk, E.T. Abdou.....	451
The effect of different amounts of animal manure on qualitative and quantitative traits of sunflower hybrid varieties	
A. Faramarzi, B. Mirshekari, H. Mohammadi, A. Ahmadi.....	455
Nitrogen fertilization of high oleic sunflower in wet climate	
Gian Paolo Vannozzi, Maurizio Turi, Sattar Tahmasebi Enferadi, Zohreh Rabiei.....	459

CROP PRODUCTION – WEEDS

Efficiency of modeling sunflower and <i>Amaranthus retroflexus</i> L. competition	
Bahram Mirshekari, Ali Faramarzi, Mahmood Poor Yousef.....	463
Sunflower protection from negative effects of 2,4-D	
Vladimir Strelkov, Ludmila Fyadyuchenko, Lidia Isakova.....	469
Análisis del crecimiento de genotipos de girasol resistentes y susceptibles a herbicidas imidazolinonas	
Fernando S. Adegas, Marcelo F. Oliveira, Alexandre M. Brighenti.....	473

- Assessment of sunflower yield maps and discrimination of late-season weed patches by using field spectroradiometry and remote sensing: the case of *Ridolfia segetum* Moris
F. López-Granados, J.M. Peña-Barragán, M. Jurado-Exposito, L. García-Torres..... 477
- Control of *Cirsium* and *Xanthium* in sunflower hybrids resistant to express 50 SX
Alexandrina Popescu..... 483
- Development of CLHA-Plus: a novel herbicide tolerance trait in sunflower conferring superior imidazolinone tolerance and ease of breeding
Carlos Sala, Mariano Bulos, Mariel Echarte, Sherry Whit, Gregory Budziszewski, William Howie, Bijay Singh, Brigitte Weston..... 489
- BREEDING AND GENETICS**
- Selection of sunflower hybrids for Banja Luka area in Bosnia and Herzegovina
Jovan Kondić, Krsto Mijanović..... 495
- Principal component analysis as a reflector of combining abilities
Mehdi Ghaffari, Ebrahim Farrokhi..... 499
- New sunflower hybrids tolerant of Tribenuron-Methyl
Sinisa Jocić, Vladimir Miklić, Goran Malidza, Nada Hladni, Sandra Gvozdenovic..... 505
- Genetic improvement of oil quality in sunflower mutants under water stressed conditions
A. Ebrahimi, P. Maury, M. Berger, F. Shariati, P. Giteu, A. Sarrafi..... 509
- Estimation of genetic diversity of sunflower single cross hybrids using principal component analysis
Hossein Zehmalzadeh Tabrizi, Hassan Monirfar, Varahram Rashidi, Mehdi Ghaffari..... 513
- Relationship between genetic distance and heterosis based on quantitative traits and SSR markers in sunflower
Sandra Gvozdenovic, Dejana Satic-Pankovic, Sinisa Jocić, Dragan Skoric..... 519
- The Perenets mutation in sunflower knocks out the wild microsomal oleate desaturase gene and leads to high oleic acid content in the seed oil
Severine Lacombe, Irène Souyris, André Berville..... 525
- Vegetation period and hybrid sunflower productivity in breeding for earliness
Sergey Gontcharov, Maria Zaharova..... 531
- Homo- and heterozygous longitudinal gradient of oleic acid content in sunflower seeds
Yakov Demurin, Oxana Borisenko, Nikolay Bochkarev..... 535
- White rot resistance, seed weight and seed oil content in sunflower test crosses
M^a Antonela Giussani, Fernando Castaño, Raúl Rodríguez, Facundo Quiroz..... 539
- Gene effects and combining abilities of sunflower morphophysiological traits
Nada Hladni, Sinisa Jocić, Vladimir Miklić, Marija Kraljajević-Balalić, Dragan Skoric..... 545

Ichraq: Première variété de tournesol d'automne au Maroc	
Abdelghani Nabloussi, Bassou Akhtouch, Mohamed Boujghagh, Mohamed El Asri, Mohamed El Fechtali.....	551
Estimation of breeding potential for tocopherols and phytosterols in sunflower	
Alicia Ayerdi Gotor, Monique Berger, Françoise Labalette, Sylvie Centis, Jean Daydé, Anne Calmon.....	555
Studies on general and specific combining abilities in sunflower	
Ebrahim Farrokhi, Abolghasem Khodabandeh, Mehdi Ghaffari.....	561
Heredabilidad de componentes de rendimiento en dos poblaciones de girasol de la EEA Pergamino	
Julio González, Nora Mancuso, Pedro Ludueña.....	567
General combining ability analysis in sunflower maintainer lines using line x tester crosses	
Ebrahim Farrokhi, Bahram Alizadeh, Mehdi Ghaffari	571
Selection of sunflower genotypes for Central Brazil	
Claudio Guilherme Portela de Carvalho, Ana Cláudia Barneche de Oliveira, Anna Karolina Grunvald, Francielle Pereira da Silva.....	575
Selection of sunflower genotypes for sowing dates in August/September in Southern region of Brazil	
Ana Cláudia Barneche de Oliveira, Cláudio Guilherme Portela de Carvalho, Anna Karolina Grunvald, Francielle Pereira da Silva.....	579
Identification of a new CMS cytoplasm and localization of its fertility restoration gene in sunflower	
Jiuhuan Feng, C.C. Jan.....	583
Germoplasma mejorado de girasol de la EEA Pergamino	
Julio González, Nora Mancuso, Pedro Ludueña.....	589
Heterosis for yield and oil content of sunflower lines developed from bi-parental populations	
G. Chigeza, P. Shanahan, M.J. Savage, K. Mashingaidze.....	595
A modifying gene affecting gamma-tocopherol content in sunflower	
María J. García-Moreno, José M. Fernández-Martínez, Begoña Pérez-Vich, Leonardo Velasco.....	601
MAPPING AND MARKER ASSISTED SELECTION	
QTL for capitulum resistance to <i>Sclerotinia sclerotiorum</i> in sunflower	
Felicity Vear, Isabelle Jouan-Dufournel, Pierre-François Bert, Frédéric Serre, Florence Cambon, Caroline Pont, Pascal Walser, Sylvie Roche, Denis Tourvieille de Labrouhe, Patrick Vincourt.....	605
HeliaGene, a bioinformatics portal for <i>Helianthus</i> sp. genomics	
Sébastien Carrere, Jérôme Gouzy, Nicolas Langlade, Pascal Gamas, Patrick Vincourt.....	611
Mapping a novel fertility restoration gene in sunflower	
Gustavo Abratti, María Eugenia Bazzalo, Alberto León.....	617
Verification of positive BAC clones near the <i>Rf1</i> gene restoring pollen fertility in the presence of the PET1 cytoplasm in sunflower (<i>Helianthus annuus</i> L.) and direct isolation of BAC ends	
Sonia Hamrit, Barbara Kusterer, Wolfgang Friedt, Renate Horn.....	623

- 629 The efficiency of different molecular indices in sunflower breeding
 Maria Duca, Ana Capatana.....
- 635 Differential gene expression in SuCMoV-tolerant and susceptible
 sunflower lines
 Daniel Mallo, Monica Poverene, Fabian Giolitti, Sergio Lenardon.....
- 641 Molecular mapping of a new induced gene for nuclear male sterility in
 sunflower (*Helianthus annuus* L.)
 Ana Capatana, Jihuan Feng, Brady A. Vick, Maria Duca, C.C. Jan.....
- 645 Fine mapping of the downy mildew resistance locus *PlARG* in sunflower
 Silke Wieckhorst, Volker Hahn, Christina M. Duble, Steven J. Knapp, Chris Carolin
 Schön, Eva Bauer.....
- 651 Identification of molecular markers linked to a new nuclear male-sterility
 gene *ms7* in sunflower (*Helianthus annuus* L.)
 Caifeng Li, Jihuan Feng, Fengming Ma, Brady A. Vick, C.C. Jan.....
- 655 Construction of a linkage map with TRAP markers and identification of
 QTL for four morphological traits in sunflower (*Helianthus annuus* L.)
 Bing Yue, Brady A. Vick, Jerry F. Miller, Xiwen Cai, Jinguo Hu.....
- 661 Candidate gene analysis and identification of TRAP and SSR markers
 linked to the *Or5* gene, which confers sunflower resistance to race E of
 broomrape (*Orobanchae cumana* Wallr.)
 Angustias Marquez-Lema, Philippe Delavault, Patricia Letouzey, Jinguo Hu, Begoña
 Pérez-Vich.....
- 667 Tribenuron-methyl resistance in accessions of annual wild sunflower
 species from the Novi Sad germplasm collection
 Stren Terzic, Jovanka Atlagic.....
- 673 Hybridization between cultivated sunflower *Helianthus annuus* L. and
 wild perennial species *Helianthus pumilus* Nuttall
 Miroslava M. Hristova-Cherbadzi, Michail Christov.....
- 679 Studies on some morphological characters of wild *Helianthus annuus* L.
 accessions with different origin
 Daniela Valkova, Miroslava Hristova-Cherbadzi, Michail Christov, Emil Penchev.....
- 685 Weedy sunflowers in France: Prevalence and first inferences on their
 origin
 Marie-Hélène Muller, Vincent Lecomte, Bernard Garric, Pierre Jouffret, Martine Letlon,
 Florent Pourgeaux, Richard Séguin.....
- 691 Characterization of hybrids from crosses between cultivated *Helianthus
 annuus* L. and subspecies *rydbergii* (Britton) Long of perennial diploid
Helianthus nuttallii
 Miroslava M. Hristova-Cherbadzi, Michail Christov.....
- 697 Preventing botanical contamination risk of sunflower hybrid seed in the
 Valle Bonaerense del Río Colorado, Argentina
 Miguel Cantamutto, Alejandro Presotto, Juan Pablo Renzi, Mónica Poverene.....

Seed morphology and oil composition of wild *Helianthus ann.*

Argentina

Miguel Cantamutto, Daniel Alvarez, Alejandro Presotto, Ivana Fernandez-Moron, Seiler, Mónica Poverene.....

***Helianthus* species in breeding research on sunflower**

Michail Christov.....

Wild sunflower species from the southeastern United States as potential sources for improving oil content and quality in cultivated sunflower

Gerald J. Seiler, Tom J. Gulya, Gary Kong.....

Cytogenetic study of an F₁ sunflower interspecific hybrid (*Helianthus annuus* x *Helianthus praecox*)

Jovanka Atlagić, Sreten Terzić..... 721

Sunflower nested core collections for association studies and phenomics

Marie Coque, Sébastien Mesnildrey, Michel Romestant, Bruno Grezes-Beset, Félicity Vear, Nicolas B. Langlade, Patrick Vincourt..... 725

Using interspecific hybrids with *Helianthus tuberosus* L. to transfer genes for quantitative traits into cultivated sunflower, *H. annuus* L.

Brent S. Hulke, Donald L. Wyse..... 729

2008 update: The USDA sunflower collection at the north central regional plant introduction station, Ames, IA, USA

Laura Fredrick Marek, Charles C. Block, Candice C.A. Gardner..... 735

***Helianthus annuus* natural populations to increase the whole genetic diversity of domesticated sunflower: the concept of neodomestication**

Hervé Serieys, Hamidreza Nooryazdan, François Kaan, Roberto Bacilieri, Jacques David, Marie-France Ostrowski, Marie-Hélène Muller, André Bervillé..... 741

VARIATION IN OIL AND MEAL QUALITY

Effect of the environment on the chemical composition and some other parameters of sunflower seed quality

Velimir Radić, Siniša Jocić, Jelena Mrda 747

Variability and genetic analysis of sterols content in sunflower seeds

Marion Alignan, Jane Roche, Felicity Vear, Patrick Vincourt, Andrée Bouniols, Muriel Cerny, Zephirin Mouloungui, Othmane Merah..... 751

Caractérisation par infra-rouge des teneurs en acides gras de la graine entière décortiquée de tournesol

Alicia Ayerdi Gotor, Philippe Moreau, Antoine Gaillard, Anne Calmon..... 757

Near infrared spectrometry (NIRS) prediction of minor components in sunflower seeds

Alicia Ayerdi Gotor, Monique Berger, Françoise Labalette, Sylvie Centis, Jean Daydé, Anne Calmon..... 763

CURRENT STATUS AND NEW USES OF THE CROP

Expansion of sunflower crop production in Brazil: a survey of future trends

Nilza Patrícia Ramos, Cláudio César de A. Buschinelli, Ariovaldo Luchiani Junior, Adriana M. Moreno Pires..... 769

775	Breeding of sunflower as a biogas substrate Volker Hahn, Martin Ganssmann.....
779	Veinte años de ensayos de girasol en Andalucía: evolución del rendimiento de semilla y riqueza grasa J. R. García Ruiz, J. Domínguez Giménez, J. García López.....
785	The situation and future directions of sunflower production in the Black Sea Region Yalcin Kaya, Mukadder Ustun Kaya, Veyzel Kaya, Ibrahim Sahin.....
791	The future potential of oleic type sunflower in Turkey Yalcin Kaya, Veyzel Kaya, Ibrahim Sahin, Mukadder Ustun Kaya, Nesrin Citak.....
797	Oil type sunflower production in Turkey Yalcin Kaya, Veyzel Kaya, Goksel Evci, Ibrahim Sahin, Mukadder Ustun Kaya.....
803	Oilcake as a fuel alternative to wood pellets Yuichi Kobayashi, Hitoshi Kato, Genta Kanai.....

New sunflower hybrids tolerant of Tribenuron-Methyl

Sinisa Jocić, Vladimir Miklic, Goran Malidza, Nada Hladni, Sandra Gvozdenovic

Institute of Field and Vegetable Crops, Maksima Gorkog 30, 21 000 Novi Sad, Serbia,

E-mail: sinJocić@ifvcns.ns.ac.yu

ABSTRACT

Discovery of a tribenuron-methyl resistant wild *Helianthus annuus* L. population (ANN-KAN) created an opportunity for expansion of a sunflower herbicide resistance breeding program. The aim of this work was the creation of sunflower hybrids resistant to tribenuron-methyl. Creation of tribenuron-methyl resistant hybrids would enable the use of a wider palette of herbicides for sunflower, more efficient chemical control of *Cirsium arvense* and more economically profitable post-emergence control of some annual broad-lea weeds in sunflower. Original populations SURES-1 and SURES-2 are homozygous for resistance to tribenuron-methyl. F₁ generations produced from the crossings are completely resistant to tribenuron-methyl, pointing to the dominant way of inheritance of this trait. Studies on the exact number of genes controlling the resistance are in progress. Tribenuron-methyl resistance was transferred from original populations into a number of mother and restorer inbred lines of cultivated sunflower. These inbred lines could enable creation of a number of hybrids resistant to tribenuron-methyl. Hybrids NS-H-2017-SU, NS-H-2018-SU and NS-H-2019-SU are resistant to doubled application dose of tribenuron-methyl. Agronomical characteristics of these hybrids are on the level with the leading conventional sunflower hybrids.

Key words: hybrid – sunflower – tolerance – tribenuron-methyl.

INTRODUCTION

The main aim of plant breeding is to develop new varieties and hybrids to meet the needs of people and domestic animals. Due to the rapid growth of the human population, loss of arable land, global climate change, and water supply problems, the production of sufficient amounts of food will be a challenge in the future. The increase of yields of cultivated plants requires not only the development of new, more productive genotypes but the advancement of growing technology as well. Plant breeding for tolerance to herbicides covers both of these aspects.

The development of plants with herbicide tolerance has been made possible by the latest insights into the mechanism and target place of herbicide action at the molecular level and by the development of new biotechnology methods. In the 1990s, a number of crop genotypes resistant to herbicides have been developed as a result (Table 1). Although it is theoretically possible to develop a plant tolerant of any kind of herbicide, only combinations of major economically important crops and herbicides possessing favorable characteristics (glyphosate, glufosinate ammonium, sulfonylurea, imidazolinones, etc.) have found an actual commercial application (Malidza et al., 1999).

Table 1. Year of first registration of herbicide tolerant crops (Malidza et al., 1999).

Year	Company	Crop
1992	Cyanamid	IMI, IR, IT Maize
1992	Du Pont	STS Soybeans
1995	Calgene	BXN Cotton
1995	AgrEvo	Liberty Link Canola
1996	Monsanto	Roundup Ready Soybeans
1996	Monsanto	Roundup Ready Canola
1997	Monsanto	Roundup Ready Cotton
1997	AgrEvo	Liberty Link Maize
1997	AgrEvo	Liberty Link Soybeans
1999	Monsanto	Roundup Ready Maize

The initial stages of plant breeding for herbicide resistance did not include any work on sunflower. Crop species for which herbicide-tolerant genotypes had been developed began to be grown more widely thanks primarily to the improved economy of their production. A result of this was a decrease of area in sunflower in South and North America, where the new technologies had been accepted without any legal limitations. Additionally, weed killing herbicides are developed less rapidly in sunflower than in the rest of field crops. Weeds cause significant yield losses in sunflower due to a lack of effective herbicides for

the suppression of broadleaf weeds and use after crop emergence. The currently existing chemical measures are ineffective against large-seeded broadleaf weeds, while the present soil herbicides are often not effective enough in the suppression of small-seeds weed species, especially in years with rainfall deficits occurring after herbicide application (Malidza et al., 2004). All this prompted sunflower researchers to begin working on the crop's tolerance to herbicides. The first major breakthrough came when Al-Khatib et al. (1998) found a population of wild *Helianthus annuus* L. (ANN-PUR) originating from Rossville, Kansas (USA) that was resistant to imidazolinone-based herbicides. Once the genetics of the resistance were studied and understood (Miller and Al-Khatib, 2000; Jocić et al., 2001), this population was used to develop the first sunflower hybrids tolerant of imidazolinone herbicides. These were developed in the USA in 2003 and Serbia and Turkey in 2004 (Jocić et al., 2004).

The discovery in Kansas, USA of a wild *Helianthus annuus* L. (ANN-KAN) population (Al-Khatib et al., 1999) resistant to a sulfonylurea herbicide (tribenuron-methyl) opened up the possibility of expanding the scope of sunflower breeding for tolerance to herbicides. The present study was aimed at the development of sunflower hybrids possessing tolerance of tribenuron-methyl. The introduction of such hybrids provides multiple benefits, including a broadened range of available herbicides in sunflower, more effective control of Canada thistle (*Cirsium arvense*), and greater cost-efficiency in the suppression of some annual broadleaf weeds after sunflower emergence (Zollinger, 2003; Malidza et al., 2006).

MATERIALS AND METHODS

The herbicide Granstar 75 WG was used in the study in two doses, the normal, recommended one (30 g/ha) and twice that (60 g/ha). In the latter years of the program, another herbicide was also used in the study to test the tolerance of the newly developed hybrids. This was Express 50-SX (500 g/kg tribenuron-methyl), a new and improved tribenuron-methyl-based herbicide manufactured by Du Pont. Express 50-SX was applied at 45 g/ha (standard dose) and 90 g/ha (double dose).

The sources of genes for tolerance to tribenuron-methyl were the populations SURES-1 and SURES-2. SURES-1 is a population of B lines obtained from the cross HA 424/3HA 406 // HA 89/ ANN-KAN, while SURES-2 is a population of restorer lines originating from the cross RHA377/3 RHA 392 // RHA 376/ ANN-KAN (Miller and Al-Khatib, 2004). Of cultivated sunflower genotypes, we used the self-pollinated B lines HA-26, VL-A-8 and HA-48 for crosses with SURES-1 and the restorer lines RHA-583, RHA-SES and RHA-N-49 for crossing with SURES-2.

The tolerance of SURES-1 and SURES-2 towards tribenuron-methyl was tested in the greenhouse during September through December 2000. In parallel with this, initial crosses were made between the two populations and the self-pollinated lines chosen for the study. During the 2001 growing season, the tolerance of the resultant F₁ generations was tested under field conditions using the double dose of tribenuron-methyl. After determining the mode of inheritance, pedigree selection was employed, with each inbred generation being treated with the double dose of Granstar 75-WG (60 g/ha). The most tolerant plants from the most tolerant progenies were selected for further breeding work. Treatment with herbicides was performed at the stage of 2-6 leaves using the knapsack sprayer Solo, 350 l/ha of water and a pressure of 2 bars. Twenty days after the treatment, phytotoxicity was assessed visually on a scale of 0 to 100% (0% - no symptoms, 100% - complete plant necrosis). Thanks to the use of a greenhouse, three inbred generations were obtained per year, which enabled us to develop the first experimental hybrids as early as 2004 and to test the general (GCA) and specific (SCA) combining abilities of the newly developed restorer lines. The testing was done using line x tester method (Singh and Choudhary, 1976). The comparative trial was carried out on a well-prepared chernozem soil at the Rimski Sancevi Experiment Field of the Institute of Field and Vegetable Crops using a randomized block design with three replications. The planting dates were optimal, intensive cultural practice was implemented during the growing season, and harvesting was done manually. The best hybrid combinations were selected and tested for tolerance to tribenuron-methyl and performance characteristics in a network of small-plot trials in 2005.

RESULTS AND DISCUSSION

Tribenuron-methyl is a herbicide that inhibits the acetolactate synthase enzyme (ALS), which is responsible for the synthesis of the amino acids valin, leucine and isoleucine. It is also one of the oldest sulfonylurea herbicides in existence (Ferguson et al., 1985) and has been among the most important herbicides in small grains for the past two decades. In Serbia, it is used in wheat crops and is the active ingredient of the Granstar 75-WG formulation (75% tribenuron-methyl) (Mitic, 2004). According to

Kolkman et al. (2004), the SURES-1 and SURES-2 populations have been found to contain the Pro197 mutation. This mutation is one of the most common mutations found in crop species tolerant of herbicides inhibiting ALS. It provides several-fold tolerance towards such herbicides compared with the susceptible genotypes. During the 2001 growing season, progenies of the source populations were found to possess full tolerance to tribenuron-methyl, meaning these populations are fully homozygous for this trait. Full susceptibility of the conventional inbred lines was confirmed as well. The F₁ generations exhibited full tolerance along with slight chlorosis, but there was absolutely no lagging behind in growth of any sort relative to the control treatment, which indicates the dominant mode of inheritance of tolerance to Granstar 75-WG. Determining the genetic basis of herbicide tolerance is a very sensitive kind of research. The first requirement is to use the double dose of the active ingredient. Environmental factors have a great influence on the expression of herbicide tolerance, as does the genetic basis of the lines receiving the tolerance genes. Because the donor populations possess many traits characteristic of the source population of wild *Helianthus annuus*, the determination of the genetics of the tolerance requires prior development of inbred lines tolerant of tribenuron-methyl. Pedigree selection was used to develop 52 inbred lines from crosses between SURES-2 and the restorer lines RHA-583, RHA-SES and RHA-N-49 as well as 46 female inbreds obtained by crossing SURES-1 and the lines Ha-26, VL-A-8 and Ha-48. All these self-pollinated lines are tolerant of the double dose of tribenuron-methyl, since the herbicide was applied at the 2-6-leaves stage in each generation during their development. Besides the herbicide tolerance, the newly developed selfed lines also have other favorable agronomic characteristics (most importantly tolerance to *Phomopsis helianthi*), as these were selected for these characteristics as well during the selection process.

The development of these lines also enabled the development of the first hybrids tolerant of tribenuron-methyl. The GCA and SCA of the new lines were tested and then the experimental hybrids were developed in 2004. All the hybrids were tested for performance characteristics and resistance to the common diseases and treated each year with the double dose of tribenuron-methyl. Based on the results, three of the hybrids were chosen for commercial production.

Due to the large volume of this research program, the present paper shows only the results for the newly developed SU hybrids NS-H-2017-SU, NS-H-2018-SU and NS-H-2019-SU. Table 2 shows the results produced by the three hybrids in two years of testing. The main requirement these hybrids must meet is to have a sufficient level of tolerance to tribenuron-methyl. What this means in concrete terms is that they have to be able to withstand the double dose of the standard, recommended dose of the active ingredient per unit area without showing any signs of phytotoxicity or any significant losses of yield or yield components. The results achieved by our hybrids have shown that they have a sufficient level of tolerance, as there were no statistically significant yield losses or reductions in the other studied traits in the treatment with the double dose of tribenuron-methyl relative to the treatment in which no herbicide was used (Table 2). Additionally, there were no visible signs of phytotoxicity either. The only thing observed was that there was some slight chlorosis seven days after the treatment, but these symptoms disappeared completely after two weeks. The second important condition the new hybrids have to fulfil is to have good performance characteristics in addition to tolerance to tribenuron-methyl. Thus, they have to have a high yield potential, a high oil content, and resistance to the common diseases so as to be able to compete with the standard sunflower hybrids used in commercial production. The check hybrids in our trials were NS-H-111, the leading sunflower hybrid in Serbia, and NS-H-43, which is a hybrid that domestic sunflower growers are well familiar with, as it has been present in Serbian sunflower production for a considerable number of years already. The results of the trials have shown that the new SU hybrids are completely on a par with the standard ones in terms of performance. The performance of NS-H-2017-SU and NS-H-2019-SU completely matched that of the class-leading NS-H-111 in terms of seed yield, oil content and oil yield, while NS-H-2018-SU performed as well as NS-H-43 despite being two weeks earlier in terms of maturation (Table 2).

Our results indicate that the new SU hybrids NS-H-2017-SU, NS-H-2018-SU and NS-H-2019-SU will find their niche in the domestic sunflower market very soon. This has been confirmed by their results in the official variety trials of the Serbian Variety Commission and their subsequent registration in the Serbian Variety List.

The source populations SURES-1 and SURES-2 are homozygously tolerant of tribenuron-methyl. The F₁ generations produced in the program are completely tolerant of tribenuron-methyl, indicating the presence of the dominant mode of inheritance. Studies to determine the exact number of genes controlling this resistance are in progress. Resistance to tribenuron-methyl has been transferred from the source populations to a number of female and self-pollinated sunflower lines. This makes it possible to develop a larger number of hybrids tolerant of tribenuron-methyl. The hybrids NS-H-2017-SU, NS-H-2018-SU and

NS-H-2019-SU are tolerant of twice the recommended dose of tribenuron-methyl per hectare and are also as good as the leading sunflower hybrids in the domestic market in terms of agronomic performance.

Table 2. Mean values of several traits in tribenuron-tolerant sunflower hybrids

Hybrid	Treatment	Plant height (cm)	Maturity (days)	Seed yield (kg/ha)	Oil content (%)	Oil yield (kg/ha)
NS-H-2017	Untreated	176.43	122.5	4 265.33	46.36	1 976.43
	Tribenuron-methyl (45 g/ha)	178.29	123	4 230.57	46.25	1 956.20
NS-H-2018	Untreated	161.40	110.4	3 702.26	47.18	1 747.45
	Tribenuron-methyl (45 g/ha)	162.30	109.9	3 806.34	48.53	1 847.28
NS-H-2019	Untreated	189.55	127.8	3 926.57	48.30	1 896.53
	Tribenuron-methyl (45 g/ha)	188.75	127.5	4 157.52	49.25	2 046.11
NS-H-43	Untreated		129	3 938.04	46.49	1 830.35
NS-H-111	Untreated		123	4 258.12	48.53	2 066.36
			LSD	476.93	4.27	260.65

ACKNOWLEDGEMENTS

This work was supported by Ministry of Science of Republic of Serbia.

REFERENCES

- Al-Khatib, K., J.R. Baumgartner, D.E. Peterson, and R.S. Currie. 1998. Imazethapyr resistance in common sunflower (*Helianthus annuus*). *Weed Sci.* 46:403-407.
- Al-Khatib, K., J.R. Baumgartner, and R.S. Currie. 1999. Survey of common sunflower (*Helianthus annuus*) resistance to ALS inhibiting herbicides in northeast Kansas. p 210-215. In: Proceedings of 21th Sunflower Research Workshop. National Sunflower Association, Bismark, N.D.
- Ferguson, D.T., S.E. Schehl, L.H. Hageman, and G.E. Lepone. 1985. DPX-L5300 – A new cereal herbicide. p. 43-48 in: The 1985 British Crop Protection Conference – Weeds.
- Malidza, G., D. Ivanovic, G. Bekavac, and S. Jasic. 1999. Znacaj genetički modifikovanih biljaka u suzbijanju stetnih organizama. *Pesticidi* 14:125-152.
- Malidza, G., S. Jocić, D. Škorić, and B. Orbovic. 2004. Clearfield sistem proizvodnje suncokreta. *Zbornik radova Instituta za ratarstvo i povrtarstvo* 40:279-290.
- Malidza, G., S. Jocić, D. Škorić, and B. Orbovic. 2006. Suzbijanje korova u suncokretu tolerantnom prema tribenuron-metilju. *Zbornik radova Instituta za ratarstvo i povrtarstvo* 42:323-331.
- Jocić, S., D. Škorić, and G. Malidza. 2001. Oplemenjivanje suncokreta na otpornost prema herbicidima. *Zbornik radova Naučnog instituta za ratarstvo i povrtarstvo* 35:223-233.
- Jocić, S., D. Škorić, and G. Malidza. 2004. Suncokret tolerantan na herbicide iz grupe imidazolinona. *J.Sci.Agric. Research/Arh. poljopr. Nauke* 65(229):81-89.
- Kolkman, M.J., B.M. Slabaugh, M.J. Bruniard, S. Berry, B.S. Bushman, C. Olungu, N. Maes, G. Abratti, A. Zambelli, F.J. Miller, A. Leon, and J.S. Knapp. 2004. Acetohydroxyacid synthase mutations conferring resistance to imidazolinone or sulfonyleurea herbicides in sunflower. *Theor. Appl.Genet.* 109:1147-1159.
- Miller, F.J., and K. Al-Khatib. 2000. Development of herbicide resistant germplasm in sunflower. p. 37-42 (Vol. II). In: Proc. 15th Int. Sunflower Conf., Toulouse, France.
- Miller, F.J., and K. Al-Khatib. 2004. Registration of two oilseed sunflower genetic stock, SURES-1 and SURES-2, resistant to tribenuron herbicide. *Crop Sci.* 39:301-302.
- Mitic, N. 2004. Pesticidi u poljoprivredi i sumarstvu u Srbiji i Crnoj Gori. *Drustvo za zastitu bilja Srbije*. Beograd.
- Singh, R.K. and B. D. Chouduary. 1976. *Biometrical Techniques in Genetics and Breeding*. Int. Bioscience Publishers. Hisar. India.
- Zollinger, R. 2003. Innovaciones en Control de Malezas en Girasol. p. 20-28. In: 2^o Congreso Argentino de Girasol, 12-13 de agosto de 2003, Buenos Aires.