

CULTIVOS  
HERBACEOS


# 17<sup>th</sup> INTERNATIONAL SUNFLOWER CONFERENCE

Vol. 2



Proceedings of the  
**17th International  
Sunflower Conference**

**Vol. 2**



**Córdoba, Spain**  
June 8-12, 2008

Sponsored by The International Sunflower Association, Paris, France



Instituto de Investigación y Formación Agraria y Pesquera  
CONSEJERÍA DE AGRICULTURA Y PESCA



International Sunflower  
Association



*Proceedings of the 17<sup>th</sup> International Sunflower Conference*  
Córdoba, Spain. June 8-12, 2008

**Editor:** Leonardo Velasco

**Editorial Committee:**

José M. Fernández-Martínez  
Luís García-Torres  
Pedro González  
José M. Melero-Vara  
Francisco Orgaz  
Begoña Pérez-Vich  
Francisco Villalobos

**Organizing Committee:**

Chairman: Juan Domínguez: Institute for Research and Training in Agriculture and Fisheries (IFAPA).  
Deputy Chairman: Juan Fernández Pérez LIMAGRAIN IBERICA  
ISA Secretariat: Laurecine Lot  
ISA Secretary General: André Pouzet  
Luis Carlos Alonso Arnedo. KOIPESOL S.A.  
José M. Fernández Martínez. Institute for Sustainable Agriculture (CSIC)  
José Antonio García de Tejada. ARLESA Semillas  
José Rafael García Ruiz. Institute for Research and Training in Agriculture and Fisheries (IFAPA)  
Luis López Bellido. University of Cordoba  
José M. Melero Vara. Institute for Sustainable Agriculture (CSIC)  
Leonardo Velasco. Institute for Sustainable Agriculture (CSIC)

Photography: *Feral sunflowers growing in a sunflower field in Santa Cruz, Córdoba, Spain.* Courtesy of Marie-Hélène Muller.

## Foreword

The proceedings of the 17<sup>th</sup> *International Sunflower Conference* contain 142 contributions from scientists of 24 countries. They include plenary lectures in several disciplines and regular communications presented in posters during the conference and discussed in the corresponding workshops. The manuscripts are classified by disciplines. They offer a good picture of the current state of the art of sunflower research and cultivation around the world.

The manuscripts in the *Proceedings* have been reviewed by an editorial committee with the main objective of helping the authors to improve their manuscripts through a critical reading. The authors received the edited manuscripts together with the comments of the reviewers and then went on to draft their final version. All the manuscripts received have been published in the *Proceedings*. The contents of the manuscripts are the responsibility of the authors. They should be considered as being privileged communications that require the express consent of the authors to be reprinted in part or as a whole. We wish to thank both the members of the Editorial Committee for their dedication to the task of editing such a large number of manuscripts, as well as all the authors for their collaboration throughout the whole edition process.

The Organizing Committee would also like to thank Diana Badder and José A. Palacios for their excellent editorial assistance in the preparation of these *Proceedings*. We are indebted to the Spanish Association of Sunflower Breeders (Asociación Española de Mejoradores de Girasol), which collaborated actively in the organization of the conference, and, very especially, to Juan Parejo, who was in charge of the financial side.

Finally, we would like to thank all the participants in the conference, who have contributed to its success by a careful preparation and revision of manuscripts and posters, presentation of their research in the workshops, and stimulating discussions throughout the conference on the scientific and technical aspects of sunflower research and cultivation in the world.

The Organizing Committee  
17<sup>th</sup> International Sunflower Conference  
Córdoba, Spain. June 8-12, 2008



## Volume 2

### Table of Contents

#### CROP PRODUCTION – PHYSIOLOGY

<b>Early responses to high crop population density in sunflower: Controls and effects of the crop self-organization process</b>	
Mónica López Pereira, Nora Trápani, Jorge Casal, Antonio Hall.....	417
<b>Early sowing as a means of drought escape in sunflower: effects on vegetative and reproductive stages</b>	
Clémentine Allinne, Nedjoua Ghoribi, Pierre Maury, Rhym Maougal, Ahmad Sarrafi, Nadia Ykhlef, Philippe Grieu .....	423
<b>SUNFLO: A joint phenotyping and modelling approach to analyse and predict the differences in yield potential of sunflower genotypes</b>	
Jérémie Lecoeur, Richard Poiré-Lassus, Angélique Christophe, Lydie Guilioni.....	429
<b>Root system and water extraction variability for sunflower hybrids</b>	
Lydie Guilioni, Ando M. Radanielson, Angélique Christophe, Jérémie Lecoeur.....	435
<b>Effects of high water table conditions on sunflower growth and quality</b>	
Satoko Yasumoto, Yukari Terakado, Morio Matsuzaki, Kensuke Okada.....	441

#### CROP PRODUCTION – FERTILIZER

<b>Optimizing of potassium and magnesium fertilizers in sunflower production</b>	
E. Sepehr, F. Nourgolipour, M.J. Malakouti.....	447
<b>Sunflower response to mineral nitrogen, organic and bio-fertilizers under two different levels of salinity</b>	
M.M. Keshta, T.Y. Rizk, E.T. Abdou.....	451
<b>The effect of different amounts of animal manure on qualitative and quantitative traits of sunflower hybrid varieties</b>	
A. Faramarzi, B. Mirshekari, H. Mohammadi, A. Ahmadi.....	455
<b>Nitrogen fertilization of high oleic sunflower in wet climate</b>	
Gian Paolo Vannozzi, Maurizio Turi, Sattar Tahmasebi Enferadi, Zohreh Rabiei.....	459

#### CROP PRODUCTION – WEEDS

<b>Efficiency of modeling sunflower and <i>Amaranthus retroflexus</i> L. competition</b>	
Bahram Mirshekari, Ali Faramarzi, Mahmood Poor Yousef.....	463
<b>Sunflower protection from negative effects of 2,4-D</b>	
Vladimir Strelkov, Ludmila Fyadyuchenko, Lidia Isakova.....	469
<b>Análisis del crecimiento de genotipos de girasol resistentes y susceptibles a herbicidas imidazolinonas</b>	
Fernando S. Adegas, Marcelo F. Oliveira, Alexandre M. Brighenti.....	473

- Assessment of sunflower yield maps and discrimination of late-season weed patches by using field spectroradiometry and remote sensing: the case of *Ridolfia segetum* Moris  
F. López-Granados, J.M. Peña-Barragán, M. Jurado-Exposito, L. García-Torres..... 477
- Control of *Cirsium* and *Xanthium* in sunflower hybrids resistant to express 50 SX  
Alexandrina Popescu..... 483
- Development of CLHA-Plus: a novel herbicide tolerance trait in sunflower conferring superior imidazolinone tolerance and ease of breeding  
Carlos Sala, Mariano Bulos, Mariel Echarte, Sherry Whit, Gregory Budziszewski, William Howie, Bijay Singh, Brigitte Weston..... 489
- BREEDING AND GENETICS**
- Selection of sunflower hybrids for Banja Luka area in Bosnia and Herzegovina  
Jovan Kondić, Krsto Mijanović..... 495
- Principal component analysis as a reflector of combining abilities  
Mehdi Ghaffari, Ebrahim Farrokhi..... 499
- New sunflower hybrids tolerant of Tribenuron-Methyl  
Sinisa Jocić, Vladimir Miklič, Goran Malidza, Nada Hladni, Sandra Gvozdenovic..... 505
- Genetic improvement of oil quality in sunflower mutants under water stressed conditions  
A. Ebrahimi, P. Maury, M. Berger, F. Shariati, P. Giteu, A. Sarrafi..... 509
- Estimation of genetic diversity of sunflower single cross hybrids using principal component analysis  
Hossein Zehmalzadeh Tabrizi, Hassan Monirfar, Varahram Rashedi, Mehdi Ghaffari..... 513
- Relationship between genetic distance and heterosis based on quantitative traits and SSR markers in sunflower  
Sandra Gvozdenovic, Dejana Satić-Pankovic, Sinisa Jocić, Dragan Skoric..... 519
- The Perenets mutation in sunflower knocks out the wild microsomal oleate desaturase gene and leads to high oleic acid content in the seed oil  
Severine Lacombe, Irène Souyris, André Berville..... 525
- Vegetation period and hybrid sunflower productivity in breeding for earliness  
Sergey Gontcharov, Maria Zaharova..... 531
- Homo- and heterozygous longitudinal gradient of oleic acid content in sunflower seeds  
Yakov Demurin, Oxana Borisenko, Nikolay Bochkarev..... 535
- White rot resistance, seed weight and seed oil content in sunflower test crosses  
M<sup>a</sup> Antonela Giussani, Fernando Castaño, Raúl Rodríguez, Facundo Quiroz..... 539
- Gene effects and combining abilities of sunflower morphophysiological traits  
Nada Hladni, Sinisa Jocić, Vladimir Miklič, Marija Kraljajević-Balalić, Dragan Skoric..... 545



<b>Ichraq: Première variété de tournesol d'automne au Maroc</b>	
Abdelghani Nabloussi, Bassou Akhtouch, Mohamed Boujghagh, Mohamed El Asri, Mohamed El Fechtali.....	551
<b>Estimation of breeding potential for tocopherols and phytosterols in sunflower</b>	
Alicia Ayerdi Gotor, Monique Berger, Françoise Labalette, Sylvie Centis, Jean Daydé, Anne Calmon.....	555
<b>Studies on general and specific combining abilities in sunflower</b>	
Ebrahim Farrokhi, Abolghasem Khodabandeh, Mehdi Ghaffari.....	561
<b>Heredabilidad de componentes de rendimiento en dos poblaciones de girasol de la EEA Pergamino</b>	
Julio González, Nora Mancuso, Pedro Ludueña.....	567
<b>General combining ability analysis in sunflower maintainer lines using line x tester crosses</b>	
Ebrahim Farrokhi, Bahram Alizadeh, Mehdi Ghaffari .....	571
<b>Selection of sunflower genotypes for Central Brazil</b>	
Claudio Guilherme Portela de Carvalho, Ana Cláudia Barneche de Oliveira, Anna Karolina Grunvald, Francielle Pereira da Silva.....	575
<b>Selection of sunflower genotypes for sowing dates in August/September in Southern region of Brazil</b>	
Ana Cláudia Barneche de Oliveira, Cláudio Guilherme Portela de Carvalho, Anna Karolina Grunvald, Francielle Pereira da Silva.....	579
<b>Identification of a new CMS cytoplasm and localization of its fertility restoration gene in sunflower</b>	
Jiuhuan Feng, C.C. Jan.....	583
<b>Germoplasma mejorado de girasol de la EEA Pergamino</b>	
Julio González, Nora Mancuso, Pedro Ludueña.....	589
<b>Heterosis for yield and oil content of sunflower lines developed from bi-parental populations</b>	
G. Chigeza, P. Shanahan, M.J. Savage, K. Mashingaidze.....	595
<b>A modifying gene affecting gamma-tocopherol content in sunflower</b>	
María J. García-Moreno, José M. Fernández-Martínez, Begoña Pérez-Vich, Leonardo Velasco.....	601
<b>MAPPING AND MARKER ASSISTED SELECTION</b>	
<b>QTL for capitulum resistance to <i>Sclerotinia sclerotiorum</i> in sunflower</b>	
Felicity Vear, Isabelle Jouan-Dufournel, Pierre-François Bert, Frédéric Serre, Florence Cambon, Caroline Pont, Pascal Walser, Sylvie Roche, Denis Tourvieille de Labrouhe, Patrick Vincourt.....	605
<b>HeliaGene, a bioinformatics portal for <i>Helianthus</i> sp. genomics</b>	
Sébastien Carrere, Jérôme Gouzy, Nicolas Langlade, Pascal Gamas, Patrick Vincourt.....	611
<b>Mapping a novel fertility restoration gene in sunflower</b>	
Gustavo Abratti, María Eugenia Bazzalo, Alberto León.....	617
<b>Verification of positive BAC clones near the <i>Rf1</i> gene restoring pollen fertility in the presence of the PET1 cytoplasm in sunflower (<i>Helianthus annuus</i> L.) and direct isolation of BAC ends</b>	
Sonia Hamrit, Barbara Kusterer, Wolfgang Friedt, Renate Horn.....	623

- 629 The efficiency of different molecular indices in sunflower breeding  
 Maria Duca, Ana Capatana.....
- 635 Differential gene expression in SuCMoV-tolerant and susceptible  
 sunflower lines  
 Daniel Mallo, Monica Poverene, Fabian Giolitti, Sergio Lenardon.....
- 641 Molecular mapping of a new induced gene for nuclear male sterility in  
 sunflower (*Helianthus annuus* L.)  
 Ana Capatana, Jihuan Feng, Brady A. Vick, Maria Duca, C.C. Jan.....
- 645 Fine mapping of the downy mildew resistance locus *PlARG* in sunflower  
 Silke Wieckhorst, Volker Hahn, Christina M. Duble, Steven J. Knapp, Chris Carolin  
 Schön, Eva Bauer.....
- 651 Identification of molecular markers linked to a new nuclear male-sterility  
 gene *ms7* in sunflower (*Helianthus annuus* L.)  
 Caifeng Li, Jihuan Feng, Fengming Ma, Brady A. Vick, C.C. Jan.....
- 655 Construction of a linkage map with TRAP markers and identification of  
 QTL for four morphological traits in sunflower (*Helianthus annuus* L.)  
 Bing Yue, Brady A. Vick, Jerry F. Miller, Xiwen Cai, Jinguo Hu.....
- 661 Candidate gene analysis and identification of TRAP and SSR markers  
 linked to the *Or5* gene, which confers sunflower resistance to race E of  
 broomrape (*Orobanchae cumana* Wallr.)  
 Angustias Marquez-Lema, Philippe Delavault, Patricia Letouzey, Jinguo Hu, Begoña  
 Pérez-Vich.....
- 667 Tribenuron-methyl resistance in accessions of annual wild sunflower  
 species from the Novi Sad germplasm collection  
 Stren Terzic, Jovanka Atlagic.....
- 673 Hybridization between cultivated sunflower *Helianthus annuus* L. and  
 wild perennial species *Helianthus pumilus* Nuttall  
 Miroslava M. Hristova-Cherbadzi, Michail Christov.....
- 679 Studies on some morphological characters of wild *Helianthus annuus* L.  
 accessions with different origin  
 Daniela Valkova, Miroslava Hristova-Cherbadzi, Michail Christov, Emil Penchev.....
- 685 Weedy sunflowers in France: Prevalence and first inferences on their  
 origin  
 Marie-Hélène Muller, Vincent Lecomte, Bernard Garric, Pierre Jouffret, Martine Letlon,  
 Florent Pourgeaux, Richard Ségura.....
- 691 Characterization of hybrids from crosses between cultivated *Helianthus  
 annuus* L. and subspecies *rydbergii* (Britton) Long of perennial diploid  
*Helianthus nuttallii*  
 Miroslava M. Hristova-Cherbadzi, Michail Christov.....
- 697 Preventing botanical contamination risk of sunflower hybrid seed in the  
 Valle Bonaerense del Río Colorado, Argentina  
 Miguel Cantamutto, Alejandro Presotto, Juan Pablo Renzi, Mónica Poverene.....



**Seed morphology and oil composition of wild *Helianthus ann.***

**Argentina**

Miguel Cantamutto, Daniel Alvarez, Alejandro Presotto, Ivana Fernandez-Moron, Seiler, Mónica Poverene.....

***Helianthus* species in breeding research on sunflower**

Michail Christov.....

**Wild sunflower species from the southeastern United States as potential sources for improving oil content and quality in cultivated sunflower**

Gerald J. Seiler, Tom J. Gulya, Gary Kong.....

**Cytogenetic study of an F<sub>1</sub> sunflower interspecific hybrid (*Helianthus annuus* x *Helianthus praecox*)**

Jovanka Atlagić, Sreten Terzić..... 721

**Sunflower nested core collections for association studies and phenomics**

Marie Coque, Sébastien Mesnildrey, Michel Romestant, Bruno Grezes-Beset, Félicity Vear, Nicolas B. Langlade, Patrick Vincourt..... 725

**Using interspecific hybrids with *Helianthus tuberosus* L. to transfer genes for quantitative traits into cultivated sunflower, *H. annuus* L.**

Brent S. Hulke, Donald L. Wyse..... 729

**2008 update: The USDA sunflower collection at the north central regional plant introduction station, Ames, IA, USA**

Laura Fredrick Marek, Charles C. Block, Candice C.A. Gardner..... 735

***Helianthus annuus* natural populations to increase the whole genetic diversity of domesticated sunflower: the concept of neodomestication**

Hervé Serieys, Hamidreza Nooryazdan, François Kaan, Roberto Bacilieri, Jacques David, Marie-France Ostrowski, Marie-Hélène Muller, André Bervillé..... 741

**VARIATION IN OIL AND MEAL QUALITY**

**Effect of the environment on the chemical composition and some other parameters of sunflower seed quality**

Velimir Radić, Siniša Jocić, Jelena Mrđa ..... 747

**Variability and genetic analysis of sterols content in sunflower seeds**

Marion Alignan, Jane Roche, Felicity Vear, Patrick Vincourt, Andrée Bouniols, Muriel Cerny, Zephirin Mouloungui, Othmane Merah..... 751

**Caractérisation par infra-rouge des teneurs en acides gras de la graine entière décortiquée de tournesol**

Alicia Ayerdi Gotor, Philippe Moreau, Antoine Gaillard, Anne Calmon..... 757

**Near infrared spectrometry (NIRS) prediction of minor components in sunflower seeds**

Alicia Ayerdi Gotor, Monique Berger, Françoise Labalette, Sylvie Centis, Jean Daydé, Anne Calmon..... 763

**CURRENT STATUS AND NEW USES OF THE CROP**

**Expansion of sunflower crop production in Brazil: a survey of future trends**

Nilza Patrícia Ramos, Cláudio César de A. Buschinelli, Ariovaldo Luchiar Junior, Adriana M. Moreno Pires..... 769

775	Breeding of sunflower as a biogas substrate Volker Hahn, Martin Ganssmann.....
779	Veinte años de ensayos de girasol en Andalucía: evolución del rendimiento de semilla y riqueza grasa J. R. García Ruiz, J. Domínguez Giménez, J. García López.....
785	The situation and future directions of sunflower production in the Black Sea Region Yalcin Kaya, Mukadder Ustun Kaya, Veyzel Kaya, Ibrahim Sahin.....
791	The future potential of oleic type sunflower in Turkey Yalcin Kaya, Veyzel Kaya, Ibrahim Sahin, Mukadder Ustun Kaya, Nesrin Citak.....
797	Oil type sunflower production in Turkey Yalcin Kaya, Veyzel Kaya, Goksel Evci, Ibrahim Sahin, Mukadder Ustun Kaya.....
803	Oilcake as a fuel alternative to wood pellets Yuichi Kobayashi, Hitoshi Kato, Genta Kanai.....



## Effect of the environment on the chemical composition and some other parameters of sunflower seed quality

Velimir Radić, Siniša Jocić, Jelena Mrđa

<sup>1</sup>Institute of Field and Vegetable Crops, Maksima Gorkog 30, Novi Sad 21000, Serbia,

E-mail: vanjar@ifvcns.ns.ac.yu

### ABSTRACT

Considering the area of sunflower cultivation in the world, the profit made by selling the seed varieties and sunflower hybrids, the number of companies whose business activity is related to production and selling of sunflower seed, the tendency of the struggle for taking the greatest market share is bigger and tougher. This type of competition conditions all seed companies to encourage as many producers as they can with their choice of hybrids and seed quality, all aimed at making bigger profits. The common argument of seed companies for attracting producers is the chemical composition of seed. Chemical composition of sunflower seed consists of great number of different organic compounds, the oil content and proteins being the most important. The objective this research has been to evaluate the differences appearing in seed quality parameters in two localities.

**Key words:** linoleic acid – oil content – oleic acid – seed germination – sunflower seed – tocopherols.

### INTRODUCTION

Seed is the beginning of a plant's new life, a complex biological system and as such it is the first and basic factor of successful plant production (Milošević et al., 1996).

Seed quality is a complex category, determined by great number of factors (purity, germination, moisture, weight of 1000 seeds and others), which are under the influence of different environmental factors (Karagić et al., 2001). One of the most important characteristics of a seed is the germination energy and germination of a seed itself (Radić, 2003). The same author further mentions that great number of factors influence the germination such as climate conditions in time of production, pollination, fertility, harvest, factors that appear during the seed drying and seed cleaning (transport from a plot, method of drying, storage of natural seed, cleaning, seed treatment) as well as factors of storage of the seed (damage possibility by various warehouse insects as well as damage during the seed usage). Škorić et al., (1996) conclude that environmental factors can influence the chemical composition of a seed as well.

Sunflower is produced in more than 40 countries of the world (Putt, 1997). Considering the area that sunflower covers in the world, the profit made from selling the seed varieties and sunflower hybrids, the number of seed companies whose business activity is related to the production and selling of sunflower seed, the tendency of a struggle for taking the greatest market share is becoming bigger and tougher. This type of competition conditions all seed companies to encourage as many producers as they can in their choice of hybrids and seed quality, all aimed at making bigger profits.

The common argument of seed companies for encouraging producers is the chemical composition of seed and its use. Chemical composition of sunflower seed consists of great number of different organic compounds, but for sunflower, oil content and composition and protein content in seed are the most important (Marinković et al., 2003).

The same authors further mention that beside fatty acids, sterols, carotenoids, phosphates and some other compounds, the quality of oil is determined by the total amount of tocopherols, as well as the content of its respective forms.

During the process of breeding, breeders are helped by information about correlations that appear between the oil content in the seed and other characteristics of a plant and seed, since they make it possible to find connections that help or prevent the successful selection for these characteristics. Most authors have dealt with studying the relation of oil content and seed yield, plant height, weight of 1000 seeds, head diameter, number of leaves, dry matter obtained during the vegetation and others .

However, some authors have dealt with problems related to the influence of the environment on oil composition and content in sunflower seed. According to Škorić (1988) and Krizmanić et al. (1992), who take into account the environmental factors that have an effect on oil content, the average daily temperatures and the amount of moisture in the soil are the factors that have the biggest effect. Marinković et al., (2003) believe that the content of oil in seed is lower if the lack of soil moisture

appears in the period of flowering-maturation. Dušanić (1994; 1998) and De la Vega and Hall (2002) believe that the main cause of variations in content and composition of oil in sunflower seed, as well as for oil yield is the influence of the production year, locality and the sowing deadline.

The aim of the research described in this paper was to determine if the content of oil,  $\alpha$ -tocopherol, linoleic and oleic acid in sunflower seed depends on the place of production, and if there are factors that can affect the sunflower seed germination.

### MATERIALS AND METHODS

The research was made on seeds produced in two localities, Argentina and Serbia, in quite similar conditions of production. Two genotypes were used: HA-26-OL (high oleic type) and HA-48 (standard type). Complete research was carried out at the Institute of Field and Vegetable Crops, Oil Crops Department.

During the research, the following parameters were used:

1. Germination – Examination of seed germination of both genotypes was repeated 6 times. Each time 50 seeds were used. Germination was determined after 10 days. Only naturally formed germinated seeds were used for determination of this parameter. Germination was expressed in percentage.
2. Oil content – Determined by classical method and expressed in percentage.
3. Tocopherols content – The content of  $\alpha$ -tocopherol was determined by liquid chromatography method (HPLC) and expressed in mg/kg oil.
4. Linoleic and oleic acid content – Determined by gas chromatography method. The content of these two acids is expressed percentage (% of the total fatty acids).

Computer programme GENSTAT was used for the analysis of variance of two factor experiment and interdependence of the observed parameters.

### RESULTS AND DISCUSSION

The results of the research show that the percentage of seed germination of the two genotypes produced in a locality in Serbia was 64% and 71% respectively (Table 1), while for those produced in a locality in Argentina germination percentage of HA-26-OL was 97% and that of HA-48, 96% (Table 2).

As for seed germination, lower values were determined for oil content for seed produced in a locality in Serbia (Table 1) than for seed produced in a locality in Argentina (Table 2). A percentage of 31.34% of oil content was determined for HA-26-OL genotype for seed produced in a locality in Serbia and 36.42% for seed produced in a locality in Argentina. For HA-48 genotype those values were 36.82% in Serbia and 48.90% in Argentina.

**Table 1.** Results on a locality in Serbia

Genotype	Seed germination (%)	Oil content and composition			
		Oil content (%)	$\alpha$ -tocopherol (mg/kg oil)	Linoleic acid (%)	Oleic acid (%)
HA-26-OL	64	31.34	425.67	19.63	72.73
HA-48	71	36.82	393.40	62.23	26.57

Similar situation was found during the observation of tocopherol content. For the seed obtained in a locality in Serbia a total tocopherol content of 425.67mg/kg oil was determined in HA-26-OL and 393.0 mg/kg oil in HA-48 genotype (Table 1). These values were higher than those observed for the seed obtained in Argentina for HA-26-OL (585.30 mg/kg oil) and for HA-48 genotype (569.45 mg/kg) (Table 2).

Considering that for this study a high oleic acid and a standard type hybrid were used, the content of linoleic and oleic acid was different between observed genotypes. Nevertheless, HA-26-OL had lower linoleic acid content (19.63%) in a locality in Serbia (Table 1) in comparison to the value observed in a locality in Argentina (35.83%) (Table 2). The same situation was observed for the HA-48 (standard type) genotype. In Serbia the result was 62.33% while in Argentina it was 67.23%.

In contrast to all observed characteristics, the content of oleic acid had a higher value in a locality in Serbia than in a locality in Argentina, in both genotypes observed (Tables 1 and 2).



Oleic acid content was 72.73% for HA-26-OL in a locality in Serbia, in comparison to 57.30% in Argentina. HA-48 had a oleic acid content of 26.57% in Serbia was 26.57% while in Argentina that value was 30.73%.

**Table 2.** Results in a locality in Argentina

Genotype	Seed	Oil content and composition			
	germination (%)	Oil content (%)	$\alpha$ -tocopherol (mg/kg oil)	Linoleic acid (%)	Oleic acid (%)
HA-26-OL	97	36.42	585.43	35.83	57.20
HA-48	96	48.90	569.45	67.23	30.73

The calculation of simple coefficients of correlation, showed a highly significant positive correlation between seed germination and tocopherol content (Table 3). A highly significant positive correlation between oil content and linoleic acid content was also observed (Table 3). Škorić (1982) and Petakov et al. (1983) reached similar conclusions. A highly significant negative correlation between linoleic and oleic acid content was identified (Table 3). These values are in agreement with the results of Seiler (1994), while Gonzales et al. (2000) determined the existence of negative correlation between both fatty acids, even though the authors found that the correlation was not statistically significant.

A significant positive correlation was observed between seed germination and oil content. Correlation between oil content and oleic acid was significantly negative (Table 3). In their research, Álvarez et al. (1992) detected the existence of a highly significant positive correlation of oleic acid with and seed oil content and weight of 1000 seeds, while the correlation coefficient between linoleic acid content and seed yield was negative but not statistically significant.

A positive but not significant correlation was detected between seed germination and linoleic acid content, between oil content and tocopherol content, as well as between linoleic acid content and tocopherol content. A weak and negative correlation was observed between seed germination and oleic acid content, and between oleic acid content and tocopherol content (Table 3).

**Table 3.** Coefficients of correlation of the observed parameters

Parameters	Oil content	Linoleic acid	Oleic acid	Tocopherols
Germination	0.675*	0.392	-0.289	0.766**
Oil content		0.791**	-0.681*	0.433
Linoleic acid			-0.985**	0.136
Oleic acid				-0.025

Studying the relationship between oil content, linoleic and oleic acid with tocopherol content, Demurin (1986) determined that there is no significant correlation between these parameters.

The results of seed germination point to a great possibility of utilizing other localities for the purpose of getting high quality seed. It is obvious that Serbia, in the case of these two genotypes, is characterized as a locality that gives weak results. The results obtained showed that the influence of the environmental factors is important, even highly significant for certain observed parameters. In contrast to this statement, Balalić et al. (2006) considered that for the interaction between the hybrids and the year there is no significant difference, while in all variation sources (relation of oil content and yield, locality, year, sowing deadline and plant density) there are significant differences. Miklič (2001) concluded that the external factors do not have a great effect on oil content in the seeds, but the influence of the genotype prevails.

Seed produced in Serbia showed poorer results than the seed produced in Argentina. On the basis of given results it can be concluded that different environmental factors affect the content and chemical composition of oil in sunflower seed with certain genotypes.

In conclusion, oleic acid content was higher in seed produced in Serbia in comparison with the same seed produced in Argentina. Differences between both locations were highly significant, except for tocopherol content. Considering the great difference in seed germination, this experiment provided information about the possibility of successful seed production in other localities. Correlations were in agreement with previous results. Considering the results obtained in this research, further research should be directed towards observation of the relationships between seed germination, oil content, tocopherol content and oleic acid content in sunflower seeds.

### ACKNOWLEDGEMENTS

This work was supported by Ministry of Science of Republic of Serbia.

### REFERENCES

- Álvarez, D., P. Ludueña, and E. Frutos. 1992. Correlation and causation among sunflower traits. p. 957-962. In: Proc. 13<sup>th</sup> Int. Sunflower Conf., Pisa, Italy.
- Balalić, I., J. Crnobarac, and N. Dušanić. 2006. Efekat rokova setve na sadržaj i prinos ulja kod suncokreta. Zbornik radova, 47, savetovanje industrije ulja, Herceg Novi, 49-51.
- Demurin, J.N. 1986. Fenotipska izmenčivost i korrjacija sadržanja tokoferola s nekim biokemijskim priznakama semjan podsolnečnika. Naučno tehnič. Buletin VNIIMK-a, vip. II (93):21-24.
- De la Vega, A.J., and A.J. Hall. 2002. Effects of planting date, genotype and their interactions on sunflower yield: I. Determinants of oil-corrected grain yield. Crop Sci. 42:1191-1201.
- Dušanić, N. 1994. Dinamika mineralnog azota u zemljištu i njegov uticaj na prinos, kvalitet zrna i iznošenje azota usevom suncokreta. Magistarska teza. Univerzitet u Novom Sadu. Poljoprivredni fakultet.
- Dušanić, N. 1998. Uticaj gustine useva na dinamiku rastezanja i prinos hibrida suncokreta, kao i neke mikroklimatske činioce. Doktorska disertacija. Univerzitet u Novom Sadu. Poljoprivredni fakultet.
- Gonzales, J., N. Mancuso, and C. Oliva. 2000. Factores geneticos, ambientales y correlaciones entre rendimiento y calidad de girasol. p. E105-E110 (Vol. II). In: Proc. 15<sup>th</sup> Int. Sunflower Conf., Toulouse, France.
- Karagić, Đ., S. Katić, V. Mihajlović, and M. Vujaković. 2001. Semenski kvaliteti domaćih sorti lucerke. Zbornik radova, 35:367-379.
- Krizmanić, M., V. Jukić, M. Bilandzić, and M. Vratarić. 1992. Značaj oplemenjivanja nekih kvantitativnih svojstava suncokreta i njihov utjecaj na urod ulja po hektaru. Sjemenarstvo, Zagreb, 4-5:241-252.
- Marinković, R., B. Dozet, and D. Vasić. 2003. Oplemenjivanje suncokreta. Monografija. Školska knjiga.
- Miklić, V. 2001. Uticaj momenta desikacije na semenski kvalitet i prinos suncokreta. Doktorska disertacija. Univerzitet u Novom Sadu. Poljoprivredni fakultet.
- Milošević, M., M. Čirović, I. Mihaljev, and P. Dokić. 1996. Opšte Semearstvo. Institut za ratarstvo i povrtarstvo Novi Sad.
- Petakov, D., P. Ivanov, and V. Nikolova. 1983. Correlations of oil content and fatty acid composition in high oleic lines and hybrids of sunflower (*Helianthus annuus* L.). Biotechnol. Biotech. Eq. 7:139-138.
- Putt, E.D. 1997. Early history of sunflower. p. 1-10. In: Schneiter AA (Ed.). Sunflower technology and production. American Society of Agronomy, Madison, WI, USA.
- Radić, V. 2003. Uticaj nepovoljnih činilaca na klijavost pojedinih genotipova kukuruza (*Zea mays* L.). Magistarska teza. Univerzitet u Novom Sadu. Poljoprivredni fakultet.
- Seiler, G., J. 1994. Oil concentration and fatty acid composition of achenes of North American *Helianthus* (Asteraceae) species. Econ. Bot. 48:271-279.
- Škorić, D. 1982. Correlations for important agronomic characters between parent lines and F<sub>1</sub> hybrids of sunflower. In: Proc. 10<sup>th</sup> Int. Sunflower Conf., Surfers Paradise, Australia.
- Škorić, D. 1988. Sunflower breeding. Uljarstvo, 25:1-90.
- Škorić, D., M. Mihaljević, S. Jocić, R. Marinković, B. Dozet, J. Atlagić, and N. Hladni. 1996. Najnovija dostignuća u oplemenjivanju suncokreta. Zbornik radova sa 37. savetovanja o proizvodnji i preradi uljarica, Budva, 18-25.