

# Soybean Research for Sustainable Development



Abstracts

**World Soybean Research Conference 11**

18-23 June 2023, Vienna, Austria

# GOLD SPONSORS



# SILVER SPONSORS



# BRONZE SPONSORS

# SPECIAL SPONSORS



food cluster of lower austria



Kofinanziert von der Europäischen Union



# EXHIBITORS



Immer einen Schritt voraus!



Johann Vollmann · Marjana Vasiljević · Leopold Rittler ·  
Jegor Miladinović · Donal Murphy-Bokern

Editors

# **Soybean Research for Sustainable Development**

Abstracts of the World Soybean Research Conference 11 (WSRC 11)  
18-23 June 2023  
Vienna, Austria



University of Natural Resources and Life Sciences, Vienna, Austria

### *Editors*

Johann Vollmann, Marjana Vasiljević, Leopold Rittler, Jegor Miladinović, Donal Murphy-Bokern

### *Title*

**Soybean Research for Sustainable Development. Abstracts of the World Soybean Research Conference 11 (WSRC 11), 18-23 June 2023, Vienna, Austria**

### *Publisher*

University of Natural Resources and Life Sciences, Vienna, Austria

ISBN: 978-3-900397-09-8. doi: 10.5281/zenodo.7974681

Corresponding editor: Dr. Johann Vollmann, ✉ [johann.vollmann@boku.ac.at](mailto:johann.vollmann@boku.ac.at)

### *Edition 1.1*

The abstracts submitted by the authors have not undergone a rigorous editorial review. Thus, full responsibility for text and scientific content of each abstract is with the respective authors.

### **Correct citation**

Vollmann J., Vasiljević M., Rittler L., Miladinović J., Murphy-Bokern D. (eds.) 2023: Soybean Research for Sustainable Development. Abstracts of the World Soybean Research Conference 11 (WSRC 11), 18-23 June 2023, Vienna, Austria. University of Natural Resources and Life Sciences, Vienna, Austria. doi: 10.5281/zenodo.7974681

Pdf e-book published through

ZENODO repository: <https://doi.org/10.5281/zenodo.7974681> and LegumeHub: [www.legumehub.eu](http://www.legumehub.eu)

### **Front page image source**

Nikolaus Joseph v. Jacquin (1781-1786): *Icones Plantarum Rariorum*. Vol. I, C.F. Wappler Publ., Vienna. Plate 145. For detail information see poster abstract on page 518.

**Conference venue:** Austria Center Vienna, Bruno-Kreisky-Platz 1, 1220 Vienna, Austria

---

## Acknowledgements

The editors thank Mrs. Theresa Schauppenlehner (Columbus Congress & Events, Vienna, Austria) for technical support during the preparation of this document. The editors are also grateful to Mrs. Marcella Gross-Varga (Donau Soja, Vienna, Austria) for handling author communications. Mr. Martin Pachner (University of Natural Resources and Life Sciences Vienna, Tulln an der Donau, Austria) contributed to abstract formatting. Mr. Xindong Yao (Harbin, Heilongjiang, China) contributed to author communications and translation of Chinese abstracts. Mr. Matthias Svojtka (University of Vienna, Botany Section of Biology and Botany Library, Vienna, Austria) provided access to N.J. Jacquin's 1781-1786 publication containing the first colored image of a soybean plant (see front cover page). The editors are also grateful to members of the scientific committee who reviewed all submitted contributions for their scientific content.

# World Soybean Research Conference 11 (2023)

## International Scientific Committee

Alvadi Balbinot | Embrapa, Brazil  
Kristin Bilyeu | University of Missouri, USA  
Adeney de Freitas Bueno | Embrapa, Brazil  
Marina Čeran | Institute of Field and Vegetable Crops, Serbia  
Vuk Djordjević | Institute of Field and Vegetable Crops, Serbia  
Tomislav Duvnjak | Agricultural Institute Osijek, Croatia  
Julio Ferrarotti | Haziak, Argentina  
Volker Hahn | University of Hohenheim, Germany  
Tianfu Han | Chinese Academy of Agricultural Sciences, China  
Fernando Henning | Embrapa, Brazil  
Jeong-Dong Lee | Kyungpook National University, South Korea  
Zenglu Li | University of Georgia, USA  
Keshun Liu | United States Department of Agriculture, USA  
Febina Mathew | North Dakota State University, USA  
Mark Messina | Loma Linda University, USA  
Jegor Miladinović | Institute of Field and Vegetable Crops, Serbia  
Alexandre Nepomuceno | Embrapa, Brazil  
Henry Nguyen | University of Missouri, USA  
Leonardo Daniel Ploper EEAOC, Tucumán, Argentina  
Vara Prasad | Kansas State University, USA  
Istvan Rajcan | University of Guelph, Canada  
Milind Ratnaparkhe | Indian Council of Agriculture Research (ICAR), India  
Asheesh Singh | Iowa State University, USA  
Johann Vollmann (chair) | University of Natural Resources and Life Sciences Vienna, Austria  
Deyue Yu | National Key Laboratory of Crop Genetics and Germplasm Enhancements, China

## Local Organizing Committee

### Institute of Field and Vegetable Crops (Novi Sad)

Vuk Djordjević  
Dragana Latković  
Jegor Miladinović  
Marjana Vasiljević

### Donau Soja Association (Vienna)

Elisabeth Berger  
Marcella Gross-Varga  
Matthias Kroen (chair)  
Donal Murphy-Bokern  
Leopold Rittler

### Columbus Congress & Events (Vienna)

Renate Androsch-Holzer  
Sissy Aschenbach  
Karin Fussi  
Theresa Schauppenlehner

# WSRC Continuing Committee

The Continuing Committee for the World Soybean Research Conference (WSRC) is comprised of four soybean specialists from each of four regions of the world for a two-conference term plus the past chair-person. Two members from each region are elected at each conference.

## **Chair of the WSRC Continuing Committee**

Ricardo Vilela Abdelnoor | Brazilian Agriculture Research Corporation (Embrapa Soja), Londrina, PR, Brazil

## **REGION I - USA and Canada**

Randall Nelson | USDA-ARS, Urbana, IL, USA

Zenglu Li | University of Georgia, Athens, GA, USA

Istvan Rajcan | University of Guelph, Guelph, Canada

Lorna Woodrow | Agriculture and Agri-Food Canada, Harrow, Canada

## **REGION II - México, Central America, South America, and the Caribbean Islands**

Ricardo Vilela Abdelnoor | Brazilian Agriculture Research Corporation (Embrapa Soja), Londrina, PR, Brazil

Alexandre Jose Cattelan | Brazilian Agriculture Research Corporation (Embrapa Soja), Londrina, PR, Brazil

Eligio N. Morandi | Universidad Nacional de Rosario, Cordoba, Argentina

Julio Ferrarotti | Haziak, Rosario, Argentina

## **REGION III - Continental Asia (Bangladesh, Burma, Cambodia, India, Nepal, Laos, Malaysia, Pakistan, Thailand, Vietnam, China, North and South Korea)**

Philips Varghese | Agharkar Research Institute (MACS), Maharashtra, India)

Wenbin Li | Northeast Agricultural University, Harbin, China

Xiaobing Liu | Northeast Institute of Geography and Agroecology, Harbin, China

Deyue Yu | Nanjing Agricultural University/National Center for Soybean Improvement, Nanjing, China

## **REGION IV – Africa and adjacent Island countries, Indonesia, Europe, Middle East, Oceania, Philippines, Sri Lanka, Australia, New Zealand, Japan, Russia and East European countries**

Makita Hajika | NARO Institute of Crop Science, Tsukuba, Japan

Monique Berger | Université de Toulouse, Toulouse, France

Gerhard Jan Hendrik Scholtemeijer | Protein Research Foundation, Rivonia, South Africa

Willem Engelbrecht | K2 Seed Marketing, South Africa

## **Past chair of the WSRC Continuing Committee**

Leonardo Daniel Ploper | Estación Experimental Agroindustrial Obispo Colombres (EEAOC), Tucumán, Argentina

# Foreword

At present, the world community is facing substantial changes of different nature which have a considerable impact on the global food and nutrition situation: Man-made climate change is becoming clearly visible and is negatively affecting food security, while disruptions and changes in global supply chains also contribute to increased food insecurity as well as economic instability. The soybean stands in the middle of these processes, as it is and continues to be the most important protein and oilseed crop of the world, which is utilized for the production of livestock feeds, numerous human food components, and non-food products. Due to a global rise in the consumption of meat and dairy products during previous decades, the massive expansion of soybean production was partly associated with increased greenhouse gas emission, deforestation, and loss of biodiversity. Because of this, present soybean production is experiencing harsh criticism from science and civil society.

Therefore, the timely motto of the World Soybean Research Conference 11 is “*Soybean research for sustainable development*”. As research and development are the major engine behind any progress in crop production, agricultural research is urgently needed to mitigate the negative effects of climate change and to minimize the impact of soybean on the environment towards a fully sustainable soybean production in the near future. This includes a reduction of production inputs through improvement of agronomic and genetic input efficiency, new pathways in livestock feed uses, and finally an enhanced direct utilization of soybeans in human foods. The five major themes of the conference are (i.) breeding-genetics-genomics-biotechnology, (ii.) food-feed-nutrition, (iii.) agronomy-physiology-agrotechnology, (iv.) weeds-diseases-pests, and (v.) business-policy-market issues, which will be elaborated in over 30 scientific parallel sessions, 4 plenaries and 2 poster sessions. The versatility of approaches to sustainable soybean production is well reflected by miscellaneous contributions from fields such as genetic diversity, genomics-assisted soybean breeding, genome editing, stress and climate change mitigation, novel developments in soy-foods and feeds including chemistry, nutrition and health, agro-technology and data-driven crop and soil management, organic soybean production, seed science, the soybean rhizosphere, progress in the management of weeds, diseases and pests, as well as policy and environmental sciences.

The WSRC 11 organizers have received over 500 scientific contributions, and the most outstanding scientists from all major soybean producing countries including China, the United States, Argentina, Canada, Brazil, India, Ukraine, France, Serbia, Japan and South Korea will share and discuss their latest results. In addition, reports from all world regions with significant soybean production will provide unique insights into specific issues of soybean cropping and utilization. Dedicated workshops have been organized to address specific needs such as soybeans for Africa, or soybean breeding for organic farming. In the time since the last World Soybean Research Conference, huge progress has been achieved in the utilization of genomic information, in data-driven agronomic decision-making as well as in various other aspects of soybean biology including the interaction with stress factors. Thus, the conference aims at

fostering the exchange of new knowledge and stimulating collaborative research among the world scientific community. Apart from oral presentations, poster sessions and an industry exhibition provide unique opportunities for personal interaction between the conference participants. Posters in particular are covering a very wide range of specific topics and research questions, and therefore poster sessions are a major source of exchange and scientific inspiration.

I am grateful to all conference chairpersons, members of the international scientific committee, members of the WSRC continuing committee and to all scientific contributors for their excellent inputs to compile an outstanding program, which delivers both a comprehensive overview as well as numerous specific highlights and new landmarks of soybean research. I also thank all sponsors, industry representatives and exhibitors for their valuable support of this conference. And most particularly, I acknowledge the help of the two WSRC organizing teams of the earlier conferences to be held in Savannah, GA, United States and Novi Sad, Serbia which had to be cancelled due to natural disasters; both teams were generously sharing their experience and preparation work they had carried out before.

During the historical Vienna World Exposition of 1873, soybean made its first legendary appearance on the international stage. In 2023 - 150 years later - the soybean community is back in Vienna. I wish all participants of the World Soybean Research Conference 11 many inspiring encounters and helpful insights for meeting the challenges ahead of us.

Johann Vollmann

Chair, International Scientific Committee, WSRC 11

Vienna, June 18, 2023



# Table of contents

Soybean lifetime achievement awards .....	1
Plenary presentations .....	11
Thematic presentations	
A. Breeding, genetics, genomics, and biotechnology .....	27
B. Food, feed, and nutrition .....	247
C. Agronomy, physiology, and agrotechnology .....	301
D. Weeds, diseases, pests, and their management .....	425
E. Business, legal, and corporate social responsibility (CSR) .....	503
F. Other themes .....	509
Sponsored pages .....	529
Author index .....	535

## Influence of two stink bug species (Heteroptera: *Pentatomidae*) on soybean morphological traits

Željko Milovac<sup>1</sup>, Filip Franeta, Marjana Vasiljević, Vuk Đorđević

<sup>1</sup>*Institute of Field and Vegetable Crops, Novi Sad, Novi Sad, Serbia*

Abundance of *Halyomorpha halys* (Stål), the brown marmorated stink bug (BMSB), and *Nezara viridula* (L.), the southern green stink bug (SGSB), after their introduction on the territory of Europe, has been constantly increasing. In order to assess potential damage, a set of trials was conducted in semi-controlled, field conditions in isolation cages, 1 m in diameter and 1.8 m in height, during 2020 and 2021. Testing was done on five different soybean varieties that belong to 00-III maturity groups (Fortuna, Romansa, Princeza, Sava and Senka). There were three types of treatments/cages, a control treatment without stink bugs (C), a treatment with BMSB specimens and a third treatment with SGSB specimens. In each isolation cage with stink bugs five specimens per plant were introduced. The results showed significant effects of stink bugs feeding on soybean plants. In 2020 the number of seeds per plant, average for all five varieties, was the highest in the C (118.9) while BMSB and SGSB treatments were with much less seeds (10.7 and 6.5 respectively). The number of completely damaged seeds was the lowest in the C (4), 11.1 in the SGSB and 26.8 in the BMSB treatment. The seed weight per plant was the highest in the C 27.6 g, in BMSB 3.5 g and the lowest in the SGSB treatment 1.7 g. In 2021 the average number of seeds per plant was the highest in the C (102.2), while 66.5 in BMSB and 58.22 in the SGSB treatment. The number of completely damaged seeds was the lowest in C (1.2), 4.6 in BMSB and 5.4 in the SGSB treatment. The seed weight per plant was the highest in C (19.7 g), in BMSB 11.1 g and the lowest in the SGSB treatment (9.6 g). Significant variations between soybean varieties and years have been observed.

**Keywords:** *Nezara viridula*, *Halyomorpha halys*, soybean yield decrease

**Acknowledgment:** This research was supported by the European Union's Horizon 2020 Project *ECOBREED -Increasing the efficiency and competitiveness of organic crop breeding* under grant agreement number 771367.

**W | S | R | C | 11**



World Soybean  
Research Conference

18 – 23 June 2023

**Vienna | Austria**

University of Natural Resources  
and Life Sciences, Vienna  
doi: 10.5281/zenodo.7974681