







Host-specific competitiveness to form nodules in *Rhizobium leguminosarum* symbiovar *viciae*

Stéphane Boivin¹, Nassima Ait Lahmidi¹, David Sherlock², Maxime Bonhomme³, Doriane Dijon¹, Karine Heulin-Gotty¹, Antoine Le-Queré¹ , Marjorie Pervent¹, Marc Tauzin¹, Georg Carlsson⁴ , Erik Jensen⁴, Etienne-Pascal Journet^{5,6} , Raphael Lopez-Bellido⁷, Marek Seidenglanz⁸, Jelena Marinkovic⁹, Stefano Colella¹ , Brigitte Brunel¹, Peter Young²  and Marc Lepetit¹ 

¹Laboratoire des Symbioses Tropicales et Méditerranéennes (LSTM) INRAE, IRD, CIRAD, University of Montpellier, Montpellier SupAgro 34398, Montpellier cedex 5, France; ²Department of Biology, University of York, York, YO10 5DD, UK; ³Laboratoire de Recherche en Sciences Végétales, CNRS, UPS, Université de Toulouse, 31326 Castanet-Tolosan, France; ⁴Department of Biosystems and Technology, Swedish University of Agricultural Sciences, SE-230 53, Alnarp, Sweden; ⁵AGroécologie, Innovation et teRritoires (AGIR) INRAE, ENSAT, 31326 Castanet-Tolosan, France; ⁶Laboratoire des Interactions Plantes Microorganismes, Université de Toulouse, INRAE, CNRS, 31326 Castanet-Tolosan, France; ⁷Departamento de Ciencias y Recursos Agrícolas y Forestales, University of Córdoba, 14071 Córdoba, Spain; ⁸AGRITEC, 78701, Šumperk, Czech Republic; ⁹Institute of Field and Vegetable Crops (IFVC), 21000 Novi Sad, Serbia

Summary

Author for correspondence:

Marc Lepetit

Tel: +33 4 67 59 38 62

Email: marc.lepetit@inra.fr

Received: 22 November 2019

Accepted: 9 December 2019

New Phytologist (2020) 226: 555–568

doi: 10.1111/nph.16392

Key words: competitiveness, genospecies, nod genes, nodules, *Pisum sativum*, *Rhizobium leguminosarum* symbiovar *viciae*, symbiosis, *Vicia faba*.

- Fabaeae legumes such as pea and faba bean form symbiotic nodules with a large diversity of soil *Rhizobium leguminosarum* symbiovar *viciae* (*Rlv*) bacteria. However, bacteria competitive to form root nodules (CFN) are generally not the most efficient to fix dinitrogen, resulting in a decrease in legume crop yields. Here, we investigate differential selection by host plants on the diversity of *Rlv*.
- A large collection of *Rlv* was collected by nodule trapping with pea and faba bean from soils at five European sites. Representative genomes were sequenced. In parallel, diversity and abundance of *Rlv* were estimated directly in these soils using metabarcoding. The CFN of isolates was measured with both legume hosts. Pea/faba bean CFN were associated to *Rlv* genomic regions.
- Variations of bacterial pea and/or faba bean CFN explained the differential abundance of *Rlv* genotypes in pea and faba bean nodules. No evidence was found for genetic association between CFN and variations in the core genome, but variations in specific regions of the *nod* locus, as well as in other plasmid loci, were associated with differences in CFN.
- These findings shed light on the genetic control of CFN in *Rlv* and emphasise the importance of host plants in controlling *Rhizobium* diversity.

Introduction

Rhizobia are soil bacteria that have the ability to form root nodules with legumes. These symbiotic organs fix atmospheric dinitrogen (N₂) into organic forms that relieve the plant of nitrogen limitation. The importance of biological nitrogen fixation (BNF) for agriculture and the nitrogen cycle in the biosphere is well recognised. *Rhizobium leguminosarum* symbiovar *viciae* (*Rlv*) is the specific symbiont of legumes of the tribe Fabaeae (formerly Viciaeae) that include crops of agronomic interest such as pea (*Pisum sativum* L.) and faba bean (*Vicia faba* L.). *Rlv* bacteria belong to the same species complex as *Rhizobium leguminosarum* symbiovars *trifolii* (*Rlt*) and *phaseoli* (*Rlp*), which nodulate clover (*Trifolium* species) and *Phaseolus* bean (*Phaseolus vulgaris* L.), respectively. *Rlv* genome sequences belong to at least five distinct genospecies that are not symbiovar specific (Kumar *et al.*, 2015). The genetic determinants of the symbiosis are plasmid encoded (Young, 2016; Andrews *et al.*, 2018), so symbiovars reflect the

symbiosis plasmid rather than the chromosome diversity. The *nodD* gene, encoding a regulator of symbiosis genes, has frequently been used as a marker to discriminate *Rlv* bacteria based on their symbiotic capacities (Zézé *et al.*, 2001; Laguerre *et al.*, 2003). Indigenous populations of *Rlv* are frequent in soils (Laguerre *et al.*, 2003; Mutch & Young, 2004). Plant benefits rely on both legume and *Rhizobium* traits that operate at different stages of interaction. Early stages of symbiotic association require *Rhizobium* and host plant compatibility to form root nodules together (ability to form nodules, AFN) that is generally assessed by inoculating the bacteria alone on its host. Nevertheless, in soil there are generally several compatible *Rlv* bacteria for a host plant and the bacteria which finally form nodules vary among multiple possible associations. Compatible combinations display various levels of competitive to form root nodules (CFN). Pioneer studies showed that CFN was not associated with the ability to fix nitrogen in *Sinorhizobium/Medicago sativa* associations (Amarger, 1981). Co-inoculation of a mixture of diverse pea-nodulating *Rlv*

strains on a panel of 104 pea genotypes, representative of the variability of the genus *Pisum*, revealed that the CFN varied greatly depending on both pea and *Rlv* genotypes and it was not associated with BNF efficiency (Bourion *et al.*, 2018). Plants may sanction BNF-inefficient partners by reducing the number of cultivable cells present in nodules (Kiers *et al.*, 2003) and preferentially stimulate the growth of symbiotic organs formed with most BNF-efficient rhizobia (Laguerre *et al.*, 2012). Nevertheless, BNF may be suboptimal because of the presence of poorly effective but highly competitive *Rlv* that outcompete BNF-efficient compatible bacteria (Laguerre *et al.*, 2003). Despite the potential interest for inoculation of pea with effective *Rlv* strains (Bremer *et al.*, 1988; Fesenko *et al.*, 1995; McKenzie *et al.*, 2001), inoculant strains are frequently outcompeted by naturally occurring ineffective rhizobia (Meade *et al.*, 1985). Understanding the determinants of CFN will allow the selection of bacteria for improved inoculation strategies (Triplett & Sadowsky, 1992; Laguerre *et al.*, 2003).

The mechanisms controlling CFN have not been fully elucidated. There is evidence for microbe–microbe interactions such as antibiosis and quorum sensing that are potentially involved in CFN (Robledo *et al.*, 1998; McAnulla *et al.*, 2007; Naamala *et al.*, 2016). It is likely that differential proliferation of competitive bacterial genotypes in the host rhizosphere contributes to CFN, but the association between symbiosis plasmid diversity and partner choice supports the hypothesis that plant–*Rhizobium* interaction mechanisms are major drivers of CFN (Moawad, Ellis, & Schmidt, 1984; Laguerre *et al.*, 2003). They involve the synthesis and secretion by rhizobia of lipo-oligosaccharides called Nod Factors (NFs). NFs are recognised by plasma membrane localised plant receptors and trigger pathways that activate nodule formation. This interaction can confer species specificity (Dénarié *et al.*, 1992; Mergaert *et al.*, 1997; Broughton *et al.*, 2000; Radutoiu *et al.*, 2007). In rhizobia transcription of the *nod* genes is controlled by NodD, a transcriptional regulator, activated by flavonoids secreted by the plant (Broughton *et al.*, 2000). The common rhizobial genes *nodABC* are responsible for biosynthesis of the core NF, and *nodIJ* are involved in their secretion from the bacteria (Mergaert *et al.*, 1997). NFs from different rhizobia share the same chitin-like *N*-acetyl glucosamine backbone with a fatty acyl chain at the nonreducing end, but differ in the backbone length, the size and the saturation of the fatty acyl chain, as well as substitutions such as glycosylation, acetylation and sulfation (Dénarié *et al.*, 1992). These differences are encoded by accessory *nod* genes and may result in variations of the interactions among rhizobial and plant species (Debellé *et al.*, 1988; Surin & Downie, 1988; Downie & Surin, 1990; Surin *et al.*, 1990; Lewis-Henderson & Djordjevic, 1991a,b; Spaink *et al.*, 1991; Bloemberg *et al.*, 1995). The *nodX* gene of some *Rlv*, such as the strain TOM, is responsible for the acetylation of the NFs at the reducing terminus, and this modification allows nodulation with specific pea genotypes from the Middle East which have a *SYM2* allele encoding a LysM-RLK able to recognise this modified NF (Firmin *et al.*, 1993; Hogg *et al.*, 2002; Sulima *et al.*, 2017). In addition, bacterial surface polysaccharide recognition by specific LysM receptors modulates plant-bacterial

recognition and potentially CFN (Kawaharada *et al.*, 2015). Secretion of bacterial effectors by the type I and type III bacterial secretion systems have been implicated in modulating the partner choice (Devine *et al.*, 1980; Sutton *et al.*, 1994; Deakin & Broughton, 2009; Linhartová *et al.*, 2010; Yang *et al.*, 2010).

In this study, we focus on early partner choice and investigate the genetic basis of CFN of *Rlv* with pea and faba bean (the most cultivated Fabaceae crops in Europe) in natural populations of bacteria. The strategy was to characterise extensively the large diversity of *Rlv* from various geographic sites (directly in soil or using trapping in pea and faba bean nodules), to measure CFN with both hosts, and to associate *Rlv* genomic variation with pea/faba bean CFN phenotype in order to identify candidate genes and/or markers controlling this early symbiotic trait.

Materials and Methods

Bacterial collection

Rhizobia were isolated from an agricultural site in each of five European countries (France, Spain, Sweden, Serbia and the Czech Republic; Supporting Information Table S1). Soils were collected from each site at the beginning of the growing season by collecting five subsamples of top soil (1–10 cm) from 6 to 15 plots belonging to diverse culture systems (Table S2). Pooled samples were formed by combining soils from the different plots at each site. Pool samples have relatively homogeneous chemical compositions, except the Spanish soil that was more clayey than others. *Rlv* bacteria were trapped by growing pea or faba bean (Table S3) in 11 pots filled with a mixture of soil and sterile siliceous sand. For each pool, 100 nodules were sampled individually from 32-d-old plants, surface sterilised in 3% sodium hypochlorite for 3 min, washed four times in sterile water, crushed in 12.5% glycerol and plated on yeast-extract–mannitol (YEM) broth. Bacteria were isolated from all pooled soil samples and both hosts except the Spanish pool (unsuccessful trapping with pea). After three cycles of single colony purification, 210 isolates were collected (130 from faba bean and 80 from pea nodules). A part of the coding sequence of the *nodD* gene was amplified by PCR using specific primers (Table S4) and sequenced (Genoscreen, Lille, France; <https://www.genoscreen.fr/>).

The diversity and abundance of *nodD* sequences in soil

Soil subsamples were taken from each of the plots at each European site, and of the pooled soil from each site that was used for plant nodulation trapping. Soil was well mixed and 100 000 copies of the QQstd-*nodD* artificial template (Table S4) were added to 250 mg soil. This template was based on the *nodD* sequence of *R. leguminosarum* sv. *viciae* strain 3841, retaining matches to the primers, but with each base of the sequence between them replaced with its complement, in order to ensure that the base composition of the amplicon was not altered but the actual sequence resembled no natural sequence. Soil DNA was extracted with the MoBio Powerlyzer Powersoil extraction

kit (Quiagen, Manchester, UK). The relative abundance of symbiovar *viciae* *nodD* genes in soil samples was estimated by PCR amplification and high-throughput DNA sequencing using the MAUI-seq protocol (Fields *et al.*, 2019). The NodD136fwd and NodD136rev primers (Table S4) were designed to amplify an informative sequence 136 bp long (excluding the primers) at the start of the coding sequences of all known *nodD* genes of symbiovar *viciae* (based on published and our unpublished data), but not of the related symbiovar *trifolii* or any other published DNA sequences. These sequences were incorporated in the extended primers (including linkers and a 12-nt random unique molecular identifier) used in the MAUI-seq method. After Illumina sequencing, the reads were processed according to the MAUI-seq protocol (Fields *et al.*, 2019) to estimate the relative abundance of NodD sequence variants, and their absolute abundance was estimated by comparison with the counts for the spiked artificial template.

Effectiveness and competitiveness measurement

Plant varieties regularly cultivated at the five European sites were used (Table S3). Seeds were surface sterilised in 3% Ca-hypochlorite solution for 10 min. Plants were grown in a glasshouse (16 h : 8 h, 22°C: 18°C, day : night cycle) in 2 l pots filled with sterilised siliceous sand. High-pressure sodium lamps with a mean photosynthetically active radiation of 250 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ were used to complement natural light at morning and evening. Pots were supplied with N-free nutrient solution (K_2HPO_4 0.8 mM, MgSO_4 1 mM, K_2SO_4 0.6 mM, CaCl_2 2 mM, NaCl 0.2 mM adjusted to pH 6.5) twice a week. Plants were inoculated at sowing with 400 ml suspension of bacteria of interest (10^7 CFU ml^{-1}) grown in YEM broth. For effectiveness experiments, shoots were collected from 6-wk-old plants inoculated with single *Rlv* isolates or well characterised control strains (P221 for pea and 3841 for faba bean). Relative normalised biomass was calculated by dividing shoot biomass by the shoot biomass obtained with the control strain in a parallel experiment on the same plant variety. For competitiveness experiments, the strain of interest was co-inoculated in equal proportion with a reference strain carrying antibiotic resistance. P1NP3CSt (Laguerre *et al.*, 2007) and P1NP2HSp (spontaneous mutant of P1NP2H; Laguerre *et al.*, 2003), resistant respectively to streptomycin ($100 \mu\text{g ml}^{-1}$) and spectinomycin ($100 \mu\text{g ml}^{-1}$), were used as reference with, respectively, pea and faba bean. Nodules (50–96 from three individual plants) were surface sterilised as mentioned above, crushed individually in 12.5% glycerol and the suspension streaked onto YEM agar plates. The bacteria present in each nodule are generally clonal and multiple infections (mixed nodules) are below 1% (Amarger, 1981). The number of nodules occupied by the reference strain was scored using antibiotic resistance. The CFN was defined as the ratio of nodules formed with the strain of interest to the reference strain. For relative comparison on a given host, CFN distribution was then centred on the median of the host dataset (i.e. 24 CFN tests for faba bean, 26 for pea).

Genome sequencing, genomic and genetic association analysis

Bacterial genomes were sequenced by MicrobesNG (Birmingham, UK, <https://microbesng.uk/>) on an Illumina HiSeq platform using a 250 bp paired-end protocol. Genomic DNA libraries were prepared using the Nextera XT Library Prep Kit (Illumina, San Diego, CA, USA). High-quality paired-end reads were assembled using the Galaxy/BBRIC pipeline (<https://bbri-pipelines.toulouse.inra.fr/>) and genome annotations were performed using EUGENE-PP (Sallet *et al.*, 2014) and RAST (<http://rast.nmpdr.org/>; Tables S8–S10). Pairwise average nucleotide identity (ANI) values were calculated using JSPECIES software (<http://jspecies.ribohost.com/jspeciesws/>), and dDDH values were calculated using the GGDC2.1 webtool (Meier-Kolthoff *et al.*, 2013). Heatmaps were built using the pheatmap R package. Core and accessory genomes of *Rlv* bacteria were generated using the SPINE/AGENT webtools (http://vfmspineagent.fsm.northwestern.edu/index_age.html; Ozer *et al.*, 2014) using the default parameters (except minimum core genome segment size to output = 200 bp). Genomic regions specific to strains of interest were selected and their presence/absence checked by a BLAST search in their genomes. The nucleotide sequences of the *nod* genes were aligned using CLUSTALOMEGA webtool (<https://www.ebi.ac.uk/services>) and biallelic single nucleotide polymorphisms (SNP) were selected. A Mixed Linear Model implemented in the EMMAX software (Kang *et al.*, 2010) was used to detect associations between CFN values and SNPs of the *nod* region. Briefly, the model, which is routinely used in plant genome-wide association studies, estimates allelic effect at each SNP accounting for the genetic relationships between individuals which are described by a kinship matrix calculated using the whole SNP dataset. Each allelic effect is then tested for significance using an *F*-test, producing then a *P*-value for each SNP tested.

Results

Diversity of *nodD* gene sequences

A *nodD* sequence was successfully amplified from each of the 210 strains that were isolated from root nodules on pea and faba bean plants grown in soils from five European countries (Table S5). The 31 distinct sequences (alleles) were more similar to symbiovar *viciae* than to *trifolii* (strain WSM1689) or *phaseoli* (strain 4292; Fig. 1). They fall into seven clades or '*nodD* groups' (bootstraps > 82%; Figs 1, S1; Table S5). The two main *nodD* groups are B1 (89 isolates) and A1 (91 isolates). The other five *nodD* groups (31 isolates) are closely related to these two groups. The strains isolated in the present study represent all known *nodD* groups and most of the known *nodD* alleles (23/31). They covered and enriched the *Rlv* diversity previously characterised on a limited number of bacteria. Although there were differences in their relative abundance, the two main *nodD* groups were found at all the European locations (Fig. 2; Tables S5, S6).




Fig. 1 Phylogenetic tree based on *nodD* gene sequences of the 210 pea and faba bean *Rlv* isolates. Blue, red and green colour boxes define the A, B and C *nodD* groups respectively. Horizontal scale indicate the number of base substitutions per site. Vertical scale is related to the number of isolates present in *nodD* groups. Detailed phylogenetic tree is provided in the Supporting Information Fig. S1.

Plant hosts select different *nodD* genotypes from the soil population

The relative frequency of *nodD* groups among rhizobia isolated from pea and faba bean root nodules was very different, even though the plants were exposed to the same set of soil populations. From all soils, the frequency of isolates belonging to the group B1 was higher in faba bean (72%) than in pea nodules (24%). An opposite result was observed for isolates belonging to the group A1 (Fig. 2; Table S6). To understand whether pea or faba bean, or both, are being selective in their choice of rhizobia, the composition of the soil populations from which they trapped symbionts is required. We used specific PCR primers for symbiovar *viciae* to amplify a part of the *nodD* sequence from DNA extracted directly from soil samples. The most abundant sequences recovered from the soil samples were in the *nodD* group A1, which was also the most abundant in the isolates from these soils (Fig. 3). The second group in abundance was B1, which was the second most abundant type among all isolates. Together, these two *nodD* groups represent 89.5% of all *nodD* sequences recovered from the soil. Only three other sequences exceeded 1% in overall frequency, and all of these were also found among the isolates. This demonstrates that direct amplification from the soil does recover

relevant sequences, and that isolates from nodules include most of the *Rlv* diversity available in the soil. Similar sequences were found in all European soils, but their relative abundance varied widely even among soils sampled from different plots at the same location (Fig. 3a), although replicate analyses of individual soil samples were reproducible (Fig. 3b). Quantitation using an artificial template added to soils immediately before DNA extraction did not work perfectly, as this sequence was not recovered from a few of the samples, perhaps because the DNA was rapidly immobilised or lost. However, results for the remaining samples (Fig. S2a) indicated that soils with a high abundance of *nodD* sequences were overwhelmingly dominated by group A1 (Fig. S2b), which is consistent with the observation that the soil mixes all had a high frequency of A1 (Fig. 3c). Group B sequences were in the minority in most soils, indicating that the high frequency of group B among faba bean nodule isolates (Fig. 2b) was the result of strong discrimination by this host.

Symbiotic effectiveness is not related to *nodD* genotype

Symbiotic effectiveness was judged by plant shoot biomass production on N-free growth medium for a representative subset of strains inoculated onto their host of origin. Large variations of




Fig. 2 Diversity of the *Rlv* isolates as a function of their host and their geographic origin. (a) Isolates from pea. (b) Isolates from faba bean. Colours represent the *nodD* groups. A1–A4: dark to pale blue; B1 and B2: dark and pale red; C: green (see Fig. 1). Origins are Sweden (SE), France (FR), Spain (SP), Czech Republic (CZ) and Serbia (RS). More details are available in Supporting Information Tables S5 and S6.

biomass were observed between isolates on both pea and faba bean but this variation was not significantly explained either by *nodD* groups or by geographic origins (ANOVA $P < 0.05$, Table S7).

Genome diversity of the *Rlv* isolates

Fifty isolates representative of the *nodD* sequence diversity as well as geographical origins were sequenced (Table S8). Based on ANI (Fig. 4) and digital DNA–DNA hybridisation (dDDH; Fig. S3) comparisons, these isolates are mainly related to three genospecies of *R. leguminosarum*, gsE (27 strains), gsC (10 strains) and gsB (3 strains), that have been previously identified in European isolates (Kumar *et al.*, 2015; Jain *et al.*, 2018; Cavassim *et al.*, 2019). Eight strains belonged to two other genospecies that were not observed in these previous studies. These two new genospecies gsF-1 and gsF-2 include respectively the *R. leguminosarum* strain TOM (Firmin *et al.*, 1993; 5 strains) and the strain *R. laguerreae* FB206 (Saïdi *et al.*, 2014; 3 strains). The last two strains (CZP3G4 and SEF4G12) display only low genomic similarities with *R. leguminosarum* bacteria ($88\% < \text{ANI} < 89\%$) but are further distinct from other *Rhizobium* species (González *et al.*, 2019).

All the *Rlv* genomes had the canonical cluster of nodulation genes, arranged in several different operons: *nodABCIJ(X)*, *nodD*, *nodFEL nodMN(T)* and *nodO*. A phylogeny based on the concatenated sequences of these genes was very similar to that based on *nodD* alone, and defined similar clusters, except that group C is split (Fig. S4). It is notable that there is no association between the *nodD* groups and the genospecies (Table S8).

Competitiveness for nodulation is host specific and associated with *nod* genotype

Although it is generally expected that *Rlv* bacteria form nodules with all legumes within the Fabaeae tribe (including pea and faba




Fig. 3 Frequencies of *nodD* sequences amplified directly from soils. (a) Samples from individual plots of five sites located in Sweden (SE), France (FR), Spain (SP), Czech Republic (CZ) and Serbia (RS). In each site, soils samples were collected from plots belonging to diverse culture systems (Supporting Information Table S2). (b) Triplicate analyses of three additional plots. (c) Duplicate samples of the mixed soils (pooled samples of the different plots of each site) used for nodulation of host plants. Group D is a clade of sequences that have not been found in nodule isolates. Ambiguous sequences are close to either A or B groups but cannot be placed in a specific group.

and for A1 strains it was 0.55 (Wilcoxon test; P -value = 0.0378). By contrast, the median CFN of B1 strains on pea was only 0.14, whereas for A1 strains it was 1.28 (Wilcoxon test; P -value = 0.0218). The results were consistent with the higher proportion of B1 strains isolated from faba bean nodules than from pea nodules (Figs 1, 2). The trade-off between competitive performance on pea and on faba bean was confirmed by focusing on 16 of these *Rlv* isolates that were tested on both host species. None of these was highly competitive on both hosts. A negative relationship between CFN on pea versus on faba bean was systematically found (Table 1; Fig. 5). Nevertheless, there was considerable variation in the CFN of strains within a *nodD* group, and even those with identical *nodD* alleles (Table 1). Some strains were uncompetitive on pea (or faba bean), as they formed no nodules in competition with the reference strain (CFN = 0). However, in these cases, their ability to form nodules when inoculated alone was confirmed (except SPFP2A11, which did not nodulate *Pisum sativum* cv. Lucy, although it was very competitive on faba bean; Fig. S5). Globally, the data indicated an association of pea and faba bean CFN with the *nodD* groups A1 and B1 respectively (Table 1). However, although all *Rlv* competitive with faba bean were found in B1 and most of the *Rlv* competitive with pea belong to A1, some A1 strains were more competitive on faba bean.

Genetic variation associated with differences in the CFN

In order to identify genetic differences that might be responsible for the CFN variation, we first examined structural variation within the *nod* gene region between bacteria of *nodD* groups A1 and B1. All the strains had the equivalent set of *nod* genes, arranged in the same order, except those in *nodD* group B1 that have the additional gene *nodT* together with a small gene of unknown function inserted downstream of *nodT*. Five strains of the group A1 have the *nodX* gene located downstream of the *nodABCII* operon but this insertion was not associated with pea/faba bean CFN in our experimental conditions (data not shown). The N-terminal end of NodO in B1 bacteria is also substantially divergent compared with A1 bacteria (only 57% identity is observed in the first 100 amino acids). Both the presence of the *nodT* gene and the *nodO* allele variation in B1 strains, compared with A1, is significantly associated with low CFN on pea, but there is no effect on faba bean ($P=0.00553$ and $P=0.3694$ respectively, Kruskal–Wallis test). It is possible that these structural differences may contribute to the contrasted CFN phenotypes on pea.

The second approach aimed to identify SNPs within the conserved *nod* genes region, which may be associated with pea/faba bean CFN phenotypes. The two hosts were first considered independently; 24 isolates have been tested on faba bean and 26 on pea. The *nod* regions of these isolates were aligned to identify the SNPs associated with pea or faba bean CFN values. In total, 1449 and 1785 biallelic SNPs (with faba bean or pea respectively) were identified. Following association tests with faba bean and pea CFN phenotypes, two clusters of 53 and 76 significant SNPs, respectively, were detected (P -value threshold of 5%; Fig. 6). In order to account for multiple tests, we calculated a chromosome-wide 5% P -value threshold for each dataset by using

permutations (see Methods S1). The empirical P -value thresholds were 0.004 and 0.0015 for the faba bean and pea dataset, respectively. Using these thresholds, no SNP was detected as significant, although the smallest SNP P -values were 0.01 and 0.0068 in the faba bean and pea association analyses, respectively, thus close to the thresholds. However, 92% (49/53) and 76% (58/76) of the top significant SNPs (i.e. with P -value < 0.05) associated with CFNs in faba bean and pea respectively, were clustered within the *nodMNO* genes zone. Using a procedure of spatial permutations of each P -value dataset (see Methods S1), the probabilities to observe in the permuted datasets >92% (faba bean) or 76% (pea) of SNPs with a P -value threshold < 0.05 and lying within the 3300 bp *nodMNO* zone, were null. This indicated that the physical clustering of these significant SNPs within the *nodMNO* zone is unlikely to be due to random chance.

To identify loci outside of the *nod* region potentially associated with CFN phenotypes, we compared bacteria with contrasted CFN phenotypes sharing close or contrasted *nod* genotypes (i.e. A vs A or A vs B). In the first approach, *nod* type A bacteria displaying contrasted pea/faba bean CFN phenotypes were compared. Five significant genomic sequences were identified and annotated by reference to the genome of the strain 3841 (Tables 2, S9). Three of these regions were found on pRL9 and pRL12, but it was not possible to assign the two others regions to a replicon because they were absent from the reference. Presence/absence of these specific genomic regions was then characterised in all sequenced genomes of Nod type A isolates, and the association with the CFN phenotype was estimated (Table S9). These five regions were significantly associated with faba bean CFN but only two of these with pea CFN (Tables 2, S9). In a second approach, the genomes of eight bacteria displaying the most contrasted CFN phenotypes (i.e. a high CFN with pea and a low CFN with faba bean, or vice versa), and belonging to the Nod type A or B were compared using the same strategy. This identified 32 specific genomic sequences (Tables 2, S10). Except for one region located on the chromosome and five regions absent in the reference strain 3841, all others clustered within 14 regions of the plasmids pRL8, pRL9, pRL10 (symbiosis plasmid) and pRL12. One of these regions included the *nodO* gene and the insertion of the *nodT* gene in the pRL10, as expected. The presence/absence of these specific genomic sequences was characterised in all genomes of isolates, and the association with the CFN phenotype was estimated (Table S10). Finally, 14 regions were significantly associated with both pea and faba bean CFN phenotypes, whereas six were solely associated with pea CFN (Table 2). Among these genomic regions, only one (encoding an acetyl-CoA hydrolase transferase) was identified in both analyses (i.e. A vs A and A vs B).

Discussion

A widespread *nod* gene polymorphism in *Rlv* is associated with host preference

The diversity of *R. leguminosarum* sv. *viciae* populations was characterised either by trapping bacteria in pea or faba bean root

Table 1 Competitive to form root nodules (CFN) of diverse *Rlv* isolates.

Strain	Isolated from	Genospecies	<i>nodD</i> group	<i>nodD</i> allele	Normalised CFN			
					Faba bean	Median	Pea	Median
SEF5G12	faba	1	A1	18	0.31	0.55	nd	1.28*
SPF4F3	faba	B	A1	18	0.6		2.41	
RSF2G1	faba	B	A1	17	0.00		2.17	
FRF1H7	faba	C	A1	21	0.2		1.78	
CZF5G8	faba	C	A1	26	1.38		0.38	
CZF5B4	faba	C	A1	18	1.03		0.16	
SEF4H7	faba	C	A1	21	0.69		nd	
RSP1E11	pea	E	A1	26	0.49		3.37	
SEP3F1	pea	E	A1	26	0.51		3.02	
FRP5H7	pea	E	A1	26	1.4		0.78	
RSP1F2	pea	F-1	A1	21	nd		0.77	
CZP2D1	pea	nd	A1	22	nd		0.06	
RSP1E6	pea	B	A4	15	nd		0.43	
CZP3C9	pea	E	A2	15	1.29		0.89	
CZP1G4	pea	E	A2	14	1.69		0.5	
CZP1G9	pea	E	A3	10	nd		2.31	
SEP2G2	pea	E	A4	11	nd		1.88	
FRP5D3	pea	E	A4	10	nd		0.00	
RSP1F10	pea	F-1	A4	10	nd		1.34	
SEF4G12	faba	1	B1	6	1.45	1.06*	0.79	0.14
CZF1B5	faba	C	B1	4	0.74		nd	
SEF4B9	faba	C	B1	4	0.71		nd	
CZF1F8	faba	E	B1	6	1.16		nd	
FRF1A1	faba	E	B1	6	0.96		nd	
FRF3E2	faba	E	B1	4	0.42		nd	
FRP3E5	pea	E	B1	6	nd		0.23	
FRP5C5	pea	E	B1	6	1.48		0.05	
FRP5A4	pea	E	B1	6	1.33		0.00	
SPF2F4	faba	F-2	B1	4	1.67		0.00	
SPF2A11 ²	faba	F-2	B1	5	1.25		0.00	
RSF1B9	faba	F-1	B1	6	0.96		0.27	
FRF1D12	faba	F-1	B1	4	0.82		0.36	
FRP3E11	pea	E	C	1	nd		0.00	

Isolates were co-inoculated with reference strains P1NP3CSt (pea) and P1NP2HSp (faba bean) onto host plants. CFN is expressed as the ration of nodules formed with the strain of interest to the reference strain. CFN values in bold are for strains on their original host species. nd, not determined.

¹Genospecies displaying <90% ANI with all known *R. leguminosarum* symbiovar *viciae* (*Rlv*) sequences.

²SPF2A11 was unable to nodulate *Pisum sativum* cultivar Lucy.

* $P < 0.05$.

nodules or by quantifying them directly in the soils. This study confirmed that bacteria of the same symbiovar may belong to different genospecies and a genospecies may include bacteria from different symbiavars (Kumar *et al.*, 2015). The phylogeny of *nod* genes located on the symbiosis plasmid is independent of chromosome variations. Lateral gene and plasmid transfers are the major drivers of the evolution of the symbiotic phenotype (at least at the level of *R. leguminosarum*) and therefore genospecies are not ecologically relevant for symbiotic traits (Kumar *et al.*, 2015; Andrews *et al.*, 2018). However this does not exclude that some genes present on the chromosome may also contribute to the symbiotic phenotype and its variation.

Most of the *Rlv* nodule isolates (87%) and sequences identified directly in the soil (90%) belong to the two main *nodD* groups A1 and B1. These groups include and extend those already defined by RFLP analysis in previous studies on bacteria trapped from European soils with pea and faba bean (Laguerre *et al.*,

2003; Mutch & Young, 2004) or from Chinese soils with faba bean (Tian *et al.*, 2010). Few examples of specific *nodD* alleles in particular geographic location were found (even when bacteria isolated outside of Europe and present in GenBank were included in the analysis), giving little support for strong phylogeographic structuring of the *Rlv* diversity. This study confirmed previous observations, made on a limited number of soils, that pea and faba bean select differentially A1 and B1 nod groups of *Rlv* from the same soil populations to form nodules (Laguerre *et al.*, 2003; Mutch & Young, 2004; Jorin & Imperial, 2015).

Characterising the diversity of the 'hidden' soil population of rhizobia

As evaluated either by the number of collected isolates or by the level of *nodD* sequences estimated in the soil, the relative sizes of *nodD* groups may vary greatly between geographic sites. Indeed,




Fig. 5 Competitive to form root nodules (CFN) of the *Rlv* isolates tested with both pea and faba bean. Data were extracted from the Table 1.

local variation of the relative composition of *Rlv* communities between soil plots in each site is large and in the same range as the variation between geographic locations. Systematic analyses have already revealed high local variation of *Rlt* population sizes and their symbiotic potentials (Wakelin *et al.*, 2018). Whether these variations are the consequences of independent evolution trends of the bacteria, agronomic practices, the presence of different hosts, environmental parameters or soil composition favouring specific groups of isolates remain to be investigated.

Nevertheless, it is noteworthy that the main *nodD* groups are maintained in all sites. This suggests that *nodD* groups of *Rlv* bacteria are nonsubstitutable. They are likely under positive selection in the various sites probably because of specific ecological functions. Whether these functions are related to their interactions with Fabaceae legumes, other plant species or to factors related to their free-living lifestyle remains to be discovered.

We have no explanation: (1) why pea trapped many *nodD* group B1 bacteria on the French pool of soil, whereas it generally trapped *nodD* group A1 bacteria on other pools; and (2) why we were repeatedly unable to trap bacteria with pea from the Spanish pool despite the presence of *nodD* group A1 bacteria in these soils. We can only speculate that CFN is interacting with abiotic and biotic environmental factors such as soil composition, pH or culture system that may contribute to modulate nodulation and may result in CFN variation. Response of CFN to environment is an open question that requires further investigation beyond the objective of this study.

The amplification of *nodD* sequences directly from soil DNA, followed by high-throughput sequencing, is a novel approach to address the challenge of describing the diversity of potential symbionts in the soil. On the whole, the sequences found and their relative abundance are consistent with the genotypes of the nodule isolates, but there is one clear anomaly. Most of the isolates from the French soil were *nodD* group B1 (Fig. 2), but all the sequences amplified from the pooled French soil were *nodD* group A1 (Fig. 3c). We can speculate that this was the result of inadequate mixing of the soil before the small subsample was taken for DNA amplification. The various French soil samples that were pooled for the nodulation tests showed extreme variation in allele frequency (Fig. 3a,b) and, in retrospect, we can see that extremely thorough mixing is needed before subsampling of pooled soils.




Fig. 6 Association between pea or faba bean competitive to form root nodules (CFN) and single nucleotide polymorphism (SNP) within the *nod* genes sequence. Black circles represent biallelic SNPs along *nod* gene cluster. The location of the *nod* genes is indicated. Red arrow correspond to the 3 kb region between *nodN* and *nodO* genes (containing *nodT* in some strains), too variable to be included in this analysis.

Table 2 Specific genomic regions associated with CFN with *Pisum sativum* or *Vicia faba*.

Genomic comparison (nod types)	Replicons ¹	Region/CDS positions ²	Annotation ³	Association with pea CFN			Association with faba bean CFN		
				Total (n = 20)	A1 (n = 12)	B1 (n = 8)	Total (n = 23)	A1 (n = 11)	B1 (n = 12)
A vs B	chromosome	833874–838499	Hypothetical protein	*	0.10	(*)	*	*	0.35
A vs B	pRL7	51839–52045	Hypothetical protein	*	1.00	1.00	0.12	1.00	(*)
A vs B	pRL8	77684–78604	LysR transcriptional regulator	*	*	0.26	*	0.22	*
A vs B		78784–79167	Endoribonuclease L-PSP	*	*	0.26	*	0.22	*
A vs B		79231–80253	Nitrilase	*	*	0.26	*	0.22	*
A vs B		19902–20633	Transcriptional regulator Crp-Fnr family	*	0.37	1.00	*	*	*
A vs B	pRL9	20661–22586	<i>fixL</i>	*	0.37	1.00	*	*	*
A vs B		24138–25490	Coproporphyrinogen III oxidase, oxygen-independent	*	0.37	1.00	*	*	(*)
A vs B		26312–27064	Transcriptional regulator Crp-Fnr family	*	0.37	1.00	*	*	(*)
A vs B		27191–27916	Truncated response regulator	*	0.37	1.00	*	*	(*)
A vs B		28190–29263	Alcohol dehydrogenase	*	0.37	1.00	*	*	(*)
A vs B, A vs A		30916–32436	Acetyl-CoA hydrolase	*	*	1.00	*	*	(*)
A vs B		32449–33273	Universal stress protein family, tandem domain	*	*	1.00	*	*	(*)
A vs B		33432–33740	hypothetical protein, miscellaneous	*	*	1.00	*	*	(*)
A vs B	pRL9	49741–50607	Universal stress protein family	*	0.37	1.00	*	*	(*)
A vs B	pRL9	8920–9131	no annotated gene	*	*	1.00	*	*	1.00
A vs B	pRL10	174948–175802	<i>nodO</i>	*	1.00	1.00	0.27	1.00	1.00
A vs B		176905–177522	Hypothetical protein	*	1.00	1.00	0.27	1.00	1.00
A vs B		177479–178927	<i>nodT</i>	*	1.00	1.00	0.27	1.00	1.00
A vs B	pRL10	205375–205767	Transposase	*	1.00	1.00	0.12	1.00	(*)
A vs B	pRL10	217295–216726	Transposase	*	1.00	1.00	0.12	1.00	(*)
A vs B	pRL10	153549–153184	Hypothetical protein	*	1.00	1.00	0.12	1.00	(*)
A vs B	pRL10	215237–215476	IS481 transposase	*	1.00	1.00	0.32	1.00	0.31
A vs B	pRL10	182899–182656	no annotated gene	*	1.00	1.00	0.27	1.00	1.00
A vs A	pRL12	57481–57729	VapC toxin protein antagonist	1.00	*	0.45	*	*	0.25
A vs A	pRL12	58177–58587	Hypothetical protein	1.00	*	0.45	*	*	0.25
A vs B	pRL12	99541–100443	Transcriptional regulator	*	0.31	0.17	*	*	*
A vs B		100563–101459	UDP-glucose 4-epimerase	*	0.31	0.17	*	*	*
A vs B		101512–102480	Hypothetical protein	*	0.31	0.17	*	*	*
A vs B		102545–103369	Aldo/keto reductase family	*	0.31	0.17	*	*	*
A vs B	pRL12	371983–372822	Hypothetical protein	*	0.17	0.12	(*)	0.53	*
A vs B		372947–373888	LysR transcriptional regulator	*	0.17	0.12	(*)	0.53	*
A vs B	pRL12	598629–598466	No annotated gene	*	*	1.00	*	*	1.00
A vs B	pRL12	868438–869751	Hypothetical protein	*	0.28	*	*	*	0.19
A vs B	scaffold#4 ⁴	255964–256491	ECF family sigma factor	*	0.66	*	*	0.21	*
A vs A	scaffold#12 ⁴	37450–40707	DNA polymerase III subunit alpha	1.00	*	0.82	*	*	0.52
A vs B	scaffold#14 ⁴	157714–158751	Aspartate-semialdehyde dehydrogenase	*	0.39	1.00	*	0.10	(*)
A vs B	scaffold#32 ⁴	52588–528333	Hypothetical protein	*	0.39	1.00	*	0.10	(*)
A vs A	scaffold#39 ⁴	21998–22116	no annotated gene	1.00	*	1.00	*	*	1.00
A vs A	scaffold#39 ⁴	28925–29324	no annotated gene	1.00	*	1.00	*	*	(*)
A vs B	scaffold#42 ⁴	16775–17494	Methyl-accepting chemotaxis protein	*	1.00	1.00	0.12	1.00	(*)
A vs B	scaffold#42 ⁴	19264–20550	DegT/DnrJ/EryC1/StrS aminotransferase	*	1.00	1.00	0.12	1.00	(*)
A vs B		20691–21362	HAD family hydrolase	*	1.00	1.00	0.12	1.00	(*)
A vs B		21391–22038	GnaT-family acetyltransferase	*	1.00	1.00	0.12	1.00	(*)
A vs B		22343–23551	Glycerophosphoryl diester phosphodiesterase	*	1.00	1.00	0.12	1.00	(*)
A vs B		23588–24670	Hypothetical protein	*	1.00	1.00	0.12	1.00	(*)

Genomic regions are listed. Associations of presence/absence of these genomic regions with pea or faba bean CFN were estimated by a Kruskal–Wallis test using the data of the Table 1. Association was tested in global populations of bacterial isolates as well as subpopulations belonging to the *nod* type A or B (the size of the population used for the test is indicated). *, $P < 0.05$; (*), $P < 0.1$; no label, P -value > 0.1 .

¹Based on sequence homology to the *Rlv* 3841 reference genome.

²Positions of the homologous sequences in the *Rlv* 3841 reference genome.

³Annotations of RAST databases.

⁴Positions of the homologous sequences in the *Rlv* RSP1E11 genome.

Competitiveness for nodulation is host specific and associated with the *nod* genotypes

Both microbe–microbe and plant–microbe interactions are likely to affect competitiveness for nodulation. As our main interest was in plant–microbe interactions, we chose to minimise the opportunity for microbe–microbe interactions by inoculating with a mixture of *Rlv* at high densities. Although most *Rlv* strains can nodulate both pea and faba bean when inoculated alone, in mixture they display highly contrasted CFN with these hosts. As this trait operates only in mixture, it explains possible discrepancies between the capacity of individual strains tested alone and their performance in competition with large *Rlv* communities in the soil. We found that Nod types A and B are associated with pea and faba bean CFN, respectively, although additional genetic determinants not linked to *nod* genes can be also involved. Whether equivalent associations of CFN with specific nod types are also found with other Fabaceae legumes (lentil, for example) remains to be investigated. The negative relationship between pea and faba bean CFN in natural isolates suggests a trade-off between these two traits. More generally, this implies that the diversity and evolution of rhizobial populations cannot be understood without considering interactions with all the potential hosts that it has encountered.

Genetic factors underlying differences in CFN

Comparative genomic analyses and genetic association studies are a powerful approach for identifying genetic determinants of important traits in symbiotic bacteria (Lipuma *et al.*, 2014; Porter *et al.*, 2017). In this study, we identified genetic markers (specific genomic regions or SNPs) directly associated with pea/faba bean CFN. As expected, most of these genomic markers belong to *Rlv* plasmids and only a few to the chromosome. The association between specific SNP clusters and the pea/faba bean CFN phenotype revealed that the *nodMNO* zone within the *nod* gene island is likely to be one genetic component of CFN. A pool-seq analysis of rhizobia populations trapped with pea, faba bean, lentil or *Vicia sativa* in a single soil has already identified this region as highly polymorphic (Jorriin & Imperial, 2015), but the potential association with CFN was not directly documented. Although global *nodMNO* sequence variation (including the insertion of *nodT* within this zone) is associated with the *nod* group, SNPs specifically related to pea and faba bean CFN were identified. The *nodM*, *N*, *T* and *O* genes encode, respectively, glucosamine synthases (likely involved in the synthesis of the GlcNAc Nod factor moiety), a putative outer membrane translocator, and a cation-specific channel. These are not essential for nodulation but may modulate the process (Economou *et al.*, 1990; Baev *et al.*, 1992; Wakelin *et al.*, 2018). Consistent with a role in modulating nodulation capacities in a host-specific manner, the *nodM* and *nodT* genes have been implicated in cultivar-specific nodulation of *Rlt* (Lewis-Henderson & Djordjevic, 1991a,b), and *nodO* was found to complement loss of *nodE*, an essential determinant of host specificity in *Rlv*, which is required for the synthesis of

NFs with unsaturated acyl moieties (Downie & Surin, 1990; Demont *et al.*, 1993).

Presence/absence of additional genomic regions that potentially contribute to the pea or/and faba bean CFN phenotypes was identified, suggesting that the control of CFN may involve other genetic determinants in addition to *nod* genes. Again, these genomic regions are mainly on plasmids and therefore are potentially horizontally transferred. Interestingly, these regions are associated with both pea and faba bean CFN, with several exceptions associated only with pea CFN (including *nodT*). The association of the presence/absence of these genomic regions with pea or faba bean CFN may vary depending on the Nod type context, arguing for interactions between *nod* genes and additional genetic loci. Altogether, the data suggest that pea and faba bean CFN may be associated with a relatively limited number of specific genetic loci. Functions of the candidate genes are consistent with a role in competitiveness for nodulation. For example a putative methyl-accepting chemotaxis protein (Rivilla *et al.*, 1995; Yost *et al.*, 1998), or a putative protein involved in a type II toxin–antitoxin system (Yost *et al.*, 2004; Lipuma *et al.*, 2014) may be implicated either in plant–microbe or microbe–microbe interactions and possibly modulate CFN. However, it would be too speculative at this stage to propose specific roles for these loci in pea/faba bean CFN. Indeed, even if the corresponding sequences are statistically associated with the CFN phenotype, this could be the result of genetic associations with variations in neighbouring genes of interest and not to direct roles in CFN. Reverse genetics studies with several combinations of various alleles will be required to confirm the biological impact of candidate genes on pea/faba bean CFN. Nevertheless, these loci represent valuable markers to select *Rlv* strains competitive for nodulation with pea and/or faba bean and open the way to the identification of genes controlling competitiveness for nodulation.







Acknowledgements

This paper is dedicated to our colleague Gisèle Laguerre who opened the way for this work. This work was supported by the FP7 European Community's Seventh Framework Programme 'LEGumes for the Agriculture of TOMorrow' under the grant agreement no. FP7-613551 and by the GrasP grant of the Agence Nationale de la Recherche.

Author contributions

ML and PY designed research. GC, EJ, E-PJ, RL-B, MS and JM collected soils. NAL, KH-G and BB performed trapping experiments. SB, MP, DD, AL-Q, SC and MT performed phylogenetic analysis and genome sequencing. DS and PY performed metabarcoding experiments from soils. SB and MB performed genetic association and comparative genomic studies. ML, PY and SB wrote the manuscript with assistance from all co-authors.

ORCID

Georg Carlsson  <https://orcid.org/0000-0002-5503-5660>
 Stefano Colella  <https://orcid.org/0000-0003-3139-6055>
 Etienne-Pascal Journet  <https://orcid.org/0000-0003-1704-2998>
 Marc Lepetit  <https://orcid.org/0000-0002-0372-5985>
 Antoine Le-Queré  <https://orcid.org/0000-0001-7470-5730>
 Peter Young  <https://orcid.org/0000-0001-5259-4830>

References

- Amarger N. 1981. Competition for nodule formation between effective and ineffective strains of *Rhizobium meliloti*. *Soil Biology and Biochemistry* 13: 475–480.
- Andrews M, De Meyer S, James EK, Stepkowski T, Hodge S, Simon MF, Young JPW. 2018. Horizontal transfer of symbiosis genes within and between rhizobial genera: occurrence and importance. *Genes* 9: 321.
- Baev N, Schultze M, Barlier I, Ha DC, Virelizier H, Kondorosi E, Kondorosi A. 1992. *Rhizobium nodM* and *nodN* genes are common *nod* genes: *nodM* encodes functions for efficiency of nod signal production and bacteroid maturation. *Journal of Bacteriology* 174: 7555–7565.
- Bloemberg GV, Kamst E, Hartevelde M, van der Drift KM, Haverkamp J, Thomas-Oates JE, Lugtenberg BJ, Spaik HP. 1995. A central domain of *Rhizobium* NodE protein mediates host specificity by determining the hydrophobicity of fatty acyl moieties of nodulation factors. *Molecular Microbiology* 16: 1123–1136.
- Bourion V, Heulin-Gotty K, Aubert V, Tisseyre P, Chabert-Martinello M, Pervent M, Delaitre C, Vile D, Siol M, Duc G *et al.* 2018. Co-inoculation of a pea core-collection with diverse rhizobial strains shows competitiveness for nodulation and efficiency of nitrogen fixation are distinct traits in the interaction. *Frontiers in Plant Science* 8: 2249.
- Bremer E, Rennie DA, Rennie RJ. 1988. Dinitrogen fixation of lentil, field pea and fababean under dryland conditions. *Canadian Journal of Soil Science* 68: 553–562.
- Broughton WJ, Jabbouri S, Perret X. 2000. Keys to symbiotic harmony. *Journal of Bacteriology* 182: 5641–5652.
- Cavassim MIA, Moeskjær S, Moslemi C, Fields B, Bachmann A, Vilhjalmsson B, Schierup MH, Young JPW, Andersen SU. 2019. The genomic architecture of introgression among sibling species of bacteria. *bioRxiv*: 526707. doi: 10.1101/526707.
- Deakin WJ, Broughton WJ. 2009. Symbiotic use of pathogenic strategies: rhizobial protein secretion systems. *Nature Reviews. Microbiology* 7: 312–320.
- Debellé F, Maillat F, Vasse J, Rosenberg C, de Billy F, Truchet G, Dénarié J, Ausubel FM. 1988. Interference between *Rhizobium meliloti* and *Rhizobium trifolii* nodulation genes: genetic basis of *R. meliloti* dominance. *Journal of Bacteriology* 170: 5718–5727.
- Demont N, Debellé F, Aurelle H, Dénarié J, Promé JC. 1993. Role of the *Rhizobium meliloti nodF* and *nodE* genes in the biosynthesis of lipo-oligosaccharidic nodulation factors. *Journal of Biological Chemistry* 268: 20134–20142.
- Dénarié J, Debellé F, Rosenberg C. 1992. Signaling and host range variation in nodulation. *Annual Review of Microbiology* 46: 497–531.
- Devine TE, Kuykendall LD, Breithaupt BH. 1980. Nodulation of soybeans carrying the nodulation-restrictive gene, *rj1*, by an incompatible *Rhizobium japonicum* strain upon mixed inoculation with a compatible strain. *Canadian Journal of Microbiology* 26: 179–182.
- Downie JA, Surin BP. 1990. Either of two *nod* gene loci can complement the nodulation defect of a nod deletion mutant of *Rhizobium leguminosarum* *bv. viciae*. *Molecular & general genetics: MGG* 222: 81–86.
- Economou A, Hamilton WD, Johnston AW, Downie JA. 1990. The *Rhizobium* nodulation gene *nodO* encodes a Ca²⁺(+)-binding protein that is exported without N-terminal cleavage and is homologous to haemolysin and related proteins. *EMBO Journal* 9: 349–354.
- Fesenko AN, Provorov NA, Orlova IF, Orlov VP, Simarov BV. 1995. Selection of *Rhizobium leguminosarum* *bv. viciae* strains for inoculation of *Pisum sativum* L. cultivars: analysis of symbiotic efficiency and nodulation competitiveness. *Plant and Soil* 172: 189–198.
- Fields B, Moeskjær S, Friman V-P, Andersen SU, Young JPW. 2019. MAUI-seq: multiplexed, high-throughput amplicon diversity profiling using unique molecular identifiers. *bioRxiv*: 538587. doi: 10.1101/538587.
- Firmin JL, Wilson KE, Carlson RW, Davies AE, Downie JA. 1993. Resistance to nodulation of cv. Afghanistan peas is overcome by *nodX*, which mediates an *O*-acetylation of the *Rhizobium leguminosarum* lipo-oligosaccharide nodulation factor. *Molecular Microbiology* 10: 351–360.
- González V, Santamaría RI, Bustos P, Pérez-Carrascal OM, Vinuesa P, Juárez S, Martínez-Flores I, Cevallos MÁ, Brom S, Martínez-Romero E *et al.* 2019. Phylogenomic *rhizobium* species are structured by a continuum of diversity and genomic clusters. *Frontiers in Microbiology* 10: 910.
- Hogg B, Davies AE, Wilson KE, Bisseling T, Downie JA. 2002. Competitive nodulation blocking of cv. Afghanistan pea is related to high levels of nodulation factors made by some strains of *Rhizobium leguminosarum* *bv. viciae*. *Molecular Plant–Microbe Interactions* 15: 60–68.
- Jain C, Rodriguez-R LM, Phillippy AM, Konstantinidis KT, Aluru S. 2018. High throughput ANI analysis of 90K prokaryotic genomes reveals clear species boundaries. *Nature Communications* 9: 5114.
- Jorin B, Imperial J. 2015. Population genomics analysis of legume host preference for specific rhizobial genotypes in the *Rhizobium leguminosarum* *bv. viciae* symbioses. *Molecular Plant–Microbe Interactions* 28: 310–318.
- Kang HM, Sul JH, Service SK, Zaitlen NA, Kong S-Y, Freimer NB, Sabatti C, Eskin E. 2010. Variance component model to account for sample structure in genome-wide association studies. *Nature Genetics* 42: 348–354.
- Kawaharada Y, Kelly S, Nielsen MW, Hjuler CT, Gysel K, Muszyński A, Carlson RW, Thygesen MB, Sandal N, Asmussen MH *et al.* 2015. Receptor-mediated exopolysaccharide perception controls bacterial infection. *Nature* 523: 308–312.
- Kiers ET, Rousseau RA, West SA, Denison RF. 2003. Host sanctions and the legume–rhizobium mutualism. *Nature* 425: 78–81.
- Kumar N, Lad G, Giuntini E, Kaye ME, Udomwong P, Shamsani NJ, Young JPW, Bailly X. 2015. Bacterial genospecies that are not ecologically coherent: population genomics of *Rhizobium leguminosarum*. *Open Biology* 5: 140133.
- Laguette G, Depret G, Bourion V, Duc G. 2007. *Rhizobium leguminosarum* *bv. viciae* genotypes interact with pea plants in developmental responses of nodules, roots and shoots. *New Phytologist* 176: 680–690.
- Laguette G, Heulin-Gotty K, Brunel B, Klonowska A, Le Queré A, Tillard P, Prin Y, Cleyet-Marel J-C, Lepetit M. 2012. Local and systemic N signaling are involved in *Medicago truncatula* preference for the most efficient *Sinorhizobium* symbiotic partners. *New Phytologist* 195: 437–449.
- Laguette G, Louvrier P, Allard M-R, Amarger N. 2003. Compatibility of rhizobial genotypes within natural populations of *Rhizobium leguminosarum* *biovar viciae* for nodulation of host legumes. *Applied and Environmental Microbiology* 69: 2276–2283.
- Lewis-Henderson WR, Djordjevic MA. 1991a. *nodT*, a positively-acting cultivar specificity determinant controlling nodulation of *Trifolium subterraneum* by *Rhizobium leguminosarum* *biovar trifolii*. *Plant Molecular Biology* 16: 515–526.
- Lewis-Henderson WR, Djordjevic MA. 1991b. A cultivar-specific interaction between *Rhizobium leguminosarum* *bv. trifolii* and subterranean clover is controlled by *nodM*, other bacterial cultivar specificity genes, and a single recessive host gene. *Journal of Bacteriology* 173: 2791–2799.
- Linhartová I, Bumba L, Mašín J, Basler M, Osička R, Kamanová J, Procházková K, Adkins I, Hejnová-Holubová J, Sadílková L *et al.* 2010. RTX proteins: a highly diverse family secreted by a common mechanism. *Federation of European Microbiology Societies* 34: 1076–1112.
- Lipuma J, Cinege G, Bodogai M, Oláh B, Kiers A, Endre G, Dupont L, Dusha I. 2014. A vapBC-type toxin-antitoxin module of *Sinorhizobium meliloti* influences symbiotic efficiency and nodule senescence of *Medicago sativa*. *Environmental Microbiology* 16: 3714–3729.
- McAnulla C, Edwards A, Sanchez-Contreras M, Sawers RG, Downie JA. 2007. Quorum-sensing-regulated transcriptional initiation of plasmid transfer and replication genes in *Rhizobium leguminosarum* *biovar viciae*. *Microbiology* 153: 2074–2082.

- McKenzie RH, Middleton AB, Solberg ED, DeMulder J, Flore N, Clayton GW, Bremer E. 2001. Response of pea to rhizobia inoculation and starter nitrogen in Alberta. *Canadian Journal of Plant Science* 81: 637–643.
- Meade J, Higgins P, O'Gara F. 1985. Studies on the inoculation and competitiveness of a *Rhizobium leguminosarum* strain in soils containing indigenous rhizobia. *Applied and Environmental Microbiology* 49: 899–903.
- Meier-Kolthoff JP, Auch AF, Klenk H-P, Göker M. 2013. Genome sequence-based species delimitation with confidence intervals and improved distance functions. *BMC Bioinformatics* 14: 60.
- Mergaert P, Montagu MV, Holsters M. 1997. Molecular mechanisms of Nod factor diversity. *Molecular Microbiology* 25: 811–817.
- Moawad HA, Ellis WR, Schmidt EL. 1984. Rhizosphere response as a factor in competition among three serogroups of indigenous *Rhizobium japonicum* for nodulation of field-grown soybeans. *Applied and Environmental Microbiology* 47: 607–612.
- Mutch LA, Young JPW. 2004. Diversity and specificity of *Rhizobium leguminosarum biovar viciae* on wild and cultivated legumes. *Molecular Ecology* 13: 2435–2444.
- Naamala J, Jaiswal SK, Dakora FD. 2016. Antibiotics resistance in Rhizobium: type, process, mechanism and benefit for agriculture. *Current Microbiology* 72: 804–816.
- Ozer EA, Allen JP, Hauser AR. 2014. Characterization of the core and accessory genomes of *Pseudomonas aeruginosa* using bioinformatic tools Spine and AGEnt. *BMC genomics* 15: 737.
- Porter SS, Chang PL, Conow CA, Dunham JP, Friesen ML. 2017. Association mapping reveals novel serpentine adaptation gene clusters in a population of symbiotic *Mesorhizobium*. *The ISME Journal* 11: 248–262.
- Radutoiu S, Madsen LH, Madsen EB, Jurkiewicz A, Fukai E, Quistgaard EM, Albrektsen AS, James EK, Thirup S, Stougaard J. 2007. LysM domains mediate lipochitin–oligosaccharide recognition and Nfr genes extend the symbiotic host range. *EMBO Journal* 26: 3923–3935.
- Rivilla R, Sutton JM, Downie JA. 1995. *Rhizobium leguminosarum* NodT is related to a family of outer-membrane transport proteins that includes TolC, PrtF, CyaE and AprF. *Gene* 161: 27–31.
- Robledo EA, Kmiecik K, Oplinger ES, Nienhuis J, Triplett EW. 1998. Trifolixin production increases nodulation competitiveness of *Rhizobium etli* CE3 under agricultural conditions. *Applied and Environmental Microbiology* 64: 2630–2633.
- Saïdi S, Ramírez-Bahena M-H, Santillana N, Zúñiga D, Álvarez-Martínez E, Peix A, Mhamdi R, Velázquez E. 2014. *Rhizobium laguerreae* sp. nov. nodulates *Vicia faba* on several continents. *International Journal of Systematic and Evolutionary Microbiology* 64: 242–247.
- Sallet E, Gouzy J, Schiex T. 2014. EuGene-PP: a next-generation automated annotation pipeline for prokaryotic genomes. *Bioinformatics* 30: 2659–2661.
- Spaink HP, Sheeley DM, van Brussel AA, Glushka J, York WS, Tak T, Geiger O, Kennedy EP, Reinhold VN, Lugtenberg BJ. 1991. A novel highly unsaturated fatty acid moiety of lipo-oligosaccharide signals determines host specificity of *Rhizobium*. *Nature* 354: 125–130.
- Sulima AS, Zhukov VA, Afonin AA, Zhernakov AI, Tikhonovich IA, Lutova LA. 2017. Selection signatures in the first exon of paralogous receptor kinase genes from the Sym2 region of the *Pisum sativum* L. Genome. *Frontiers in Plant Science* 8: 1967.
- Surin BP, Downie JA. 1988. Characterization of the *Rhizobium leguminosarum* genes *nodLMN* involved in efficient host-specific nodulation. *Molecular Microbiology* 2: 173–183.
- Surin BP, Watson JM, Hamilton WDO, Economou A, Downie JA. 1990. Molecular characterization of the nodulation gene, *nodT*, from two biovars of *Rhizobium leguminosarum*. *Molecular Microbiology* 4: 245–252.
- Sutton JM, Lea EJ, Downie JA. 1994. The nodulation-signaling protein NodO from *Rhizobium leguminosarum biovar viciae* forms ion channels in membranes. *Proceedings of the National Academy of Sciences, USA* 91: 9990–9994.
- Tian CF, Young JPW, Wang ET, Tamimi SM, Chen WX. 2010. Population mixing of *Rhizobium leguminosarum* bv. *viciae* nodulating *Vicia faba*: the role of recombination and lateral gene transfer. *FEMS Microbiology Ecology* 73: 563–576.
- Triplett EW, Sadowsky MJ. 1992. Genetics of competition for nodulation of legumes. *Annual Review of Microbiology* 46: 399–428.
- Wakelin S, Tillard G, van Ham R, Ballard R, Farquharson E, Gerard E, Geurts R, Brown M, Ridgway H, O'Callaghan M. 2018. High spatial variation in population size and symbiotic performance of *Rhizobium leguminosarum* bv. *trifolii* with white clover in New Zealand pasture soils. *PLoS ONE* 13: e0192607.
- Yang S, Tang F, Gao M, Krishnan HB, Zhu H. 2010. R gene-controlled host specificity in the legume-rhizobia symbiosis. *Proceedings of the National Academy of Sciences, USA* 107: 18735–18740.
- Yost CK, Del Bel KL, Quandt J, Hynes MF. 2004. *Rhizobium leguminosarum* methyl-accepting chemotaxis protein genes are down-regulated in the pea nodule. *Archives of Microbiology* 182: 505–513.
- Yost CK, Rochepeau P, Hynes MF. 1998. *Rhizobium leguminosarum* contains a group of genes that appear to code for methyl-accepting chemotaxis proteins. *Microbiology* 144: 1945–1956.
- Young JPW. 2016. Bacteria are smartphones and mobile genes are apps. *Trends in Microbiology* 24: 931–932.
- Zézé A, Mutch LA, Young JP. 2001. Direct amplification of *nodD* from community DNA reveals the genetic diversity of *Rhizobium leguminosarum* in soil. *Environmental Microbiology* 3: 363–370.

Supporting Information

Additional Supporting Information may be found online in the Supporting Information section at the end of the article.

Fig. S1 Phylogenetic tree based on *nodD* gene sequences of the 210 pea and faba bean *Rlv* isolates.

Fig. S2 Quantitative estimates of *nodD* sequence abundance in soils.

Fig. S3 Genomic diversity of *Rhizobium leguminosarum* sv. *viciae* isolates using dDDH analysis.

Fig. S4 Phylogenetic tree based on concatenated *nodABCDEFGHIJLMN* gene sequences.

Fig. S5 Monoinoculation experiments with uncompetitive strains (CFN = 0).

Methods S1 Procedure to estimate a chromosome-wide *P*-value threshold in the association analysis of pea/faba bean CFN phenotype with SNP data; procedure to test for the random spatial clustering of significant SNPs within a given genomic region.

Table S1 Composition of soil pools used for trapping experiments.

Table S2 Agronomic history of plots used for sampling in the five European sites.

Table S3 Pea and faba bean cultivars.

Table S4 Primers and DNA spike used in this study.

Table S5 *nodD* sequences, *nodD* groups and allele numbers of isolates and references strains used in this work.

Table S6 Distribution and enrichment of *nodD* alleles and groups.

Table S7 Pea and faba bean symbiotic efficiency of selected isolates as the function of their phylogenetic classes and their geographic origins.

Table S8 Genomic data of *Rhizobium* bacteria sequenced in this study.

Table S9 Presence/absence of candidate genes correlated with the CFN with *P. sativum* or *V. faba*.

Table S10 Presence/absence of candidate genes correlated with the CFN with *P. sativum* or *V. faba*.

Please note: Wiley-Blackwell are not responsible for the content or functionality of any supporting information supplied by the authors. Any queries (other than missing material) should be directed to the *New Phytologist* Central Office.



About New Phytologist

- *New Phytologist* is an electronic (online-only) journal owned by the New Phytologist Foundation, a **not-for-profit organization** dedicated to the promotion of plant science, facilitating projects from symposia to free access for our Tansley reviews and Tansley insights.
- Regular papers, Letters, Research reviews, Rapid reports and both Modelling/Theory and Methods papers are encouraged. We are committed to rapid processing, from online submission through to publication 'as ready' via *Early View* – our average time to decision is <26 days. There are **no page or colour charges** and a PDF version will be provided for each article.
- The journal is available online at Wiley Online Library. Visit **www.newphytologist.com** to search the articles and register for table of contents email alerts.
- If you have any questions, do get in touch with Central Office (np-centraloffice@lancaster.ac.uk) or, if it is more convenient, our USA Office (np-usaoffice@lancaster.ac.uk)
- For submission instructions, subscription and all the latest information visit **www.newphytologist.com**