

IRC | 2019 | Berlin
INTERNATIONAL RAPESEED CONGRESS

Flowering for the Future



ufop

With support from



by decision of the
German Bundestag




Book of Abstracts

15th International Rapeseed Congress

16.–19.06.2019 in Berlin

Thank you to our Sponsors!



PLATINUM



TITANIUM



DIAMOND



GOLD



SILVER



BRONZE



MEDIA PARTNER

ADDRESS

PLENARY TALKS

ORALS

POSTERS

WORKSHOPS

Table of Contents

DETAILED INFORMATION ABOUT ALL TOPICS

ADDRESS #000

PLENARY TALKS #001 – #009

1. GENETICS, GENOMICS AND BREEDING

- Orals #010 – #063
- Posters #200 – #318

2. DISEASES AND PESTS, PLANT PROTECTION AND WEEDS

- Orals #064 – #181
- Posters #319 – #410

3. AGRONOMY AND CROP SCIENCE

- Orals #114 – #140
- Posters #411 – #464

4. ANALYSIS, USE OF PRODUCTS

- Orals #141 – #147
- Posters #465 – #483

5. RAPESEED/CANOLA FOR HUMAN NUTRITION

- Orals #148 – #153
- Posters #484 – #498

6. RAPESEED/CANOLA FOR ANIMAL NUTRITION

- Orals #154 – #158
- Posters #499 – #514

7. ECONOMY AND MARKET

- Orals #159 – #163
- Posters #515 – #520

8. MUSTARD AND OTHER CRUCIFEROUS OILSEED CROPS

- Orals #164 – #172
- Posters #522 – #537

9. OTHER TOPICS

- Orals #173 – #180

WORKSHOP: BLACKLEG DISEASE: RESISTANCE AND MANAGEMENT #550 – #563

WORKSHOP: CLUBROOT IN OILSEED RAPE – FROM MINOR DISEASE TO MAJOR CHALLENGE #564 – #573

WORKSHOP: AGRONOMY – MANAGING ENVIRONMENT STRESS #574 – #578

WORKSHOP: RAPESEED/CANOLA PROTEIN FOR HUMAN NUTRITION #579 – #584

WORKSHOP: FUTURE-PROOFING INSECT PEST CONTROL IN A WORLD WITH DECLINING INSECTICIDAL OPTIONS #585 – #592

WORKSHOP: SCLEROTINIA – CURRENT AND FUTURE BREEDING METHODS #593 – #598

Detailed Information About All Topics

The IRC 2019 especially springs to life with the contributions and insights given by its participants. We are looking forward to fascinating speeches, lively discussions, and valuable poster contributions. Following, you will find eight different topics in which contributions will be presented.

1. GENETICS, GENOMICS AND BREEDING

- Pan-genomic revolution in crucifer genetics and breeding (genome organisation, structural variation, plasticity)
- New diversity, interspecific hybridization, wide crosses
- Improving plant development: plant architecture, phenology
- Genetics, physiological basis and improvement of resource use efficiency
- Genetics and breeding for improved seed composition for human and animal nutrition (oil, protein, minor components)
- Breeding for higher heterosis and hybrid yield in OSR/canola
- Transgenics and New Breeding Techniques (NBT) – applications in OSR/canola research and breeding
- Genomic selection in OSR/canola
- Breeding for abiotic stress tolerance in OSR/canola (cold, heat, drought, etc.)

2. DISEASES AND PESTS, PLANT PROTECTION AND WEEDS

- Major fungal and viral diseases, regional impact and measures of control (e.g. Blackleg, Clubroot, Sclerotinia, Verticillium, Alternaria, TuYV)
- Breeding for disease resistance
- Chemical protection against insect pests, safeguarding beneficials and non-target organisms (e.g. bees)
- Breeding for insect resistance or tolerance in OSR/canola
- Weed control in OSR/canola incl. herbicide resistance

3. AGRONOMY AND CROP SCIENCE

- International comparison of OSR/canola cultivation
- Optimizing crop rotations for/with OSR/canola
- NUE – Nutrient use efficiency (N, P, other)
- Requirements of OSR/canola cultivation in temperate regions
- Identifying suitable variety types adapted to adverse conditions

4. ANALYSIS, USE OF PRODUCTS

- Economy in gross quality of OSR/canola commodities (long-time trend)
- Seed chemistry and seed composition
- Oil quality (low sats, omega-3, HOLLI, HEAR)
- Meal quality – protein and antinutritives (fibre, glucosinolates, phytate, sinapin): Genetic vs technological approach
- OSR/canola oil as biofuel

5. RAPESEED/CANOLA FOR HUMAN NUTRITION

- OSR/canola oil for human nutrition
- Oil composition vs. stability and functionality – Quality requirements for oil from OSR/canola (minor components, sensoric aspects)
- “Fish oil” (EPA, DHA) from crucifers (OSR/canola)
- Protein for human nutrition
- Politics, markets, consumer affairs (e.g. GMO)

6. RAPESEED/CANOLA FOR ANIMAL NUTRITION

- Requirements for the use of OSR/canola cake and extraction meal: breeders’ and nutritionists’ view
- Improvement of meal/protein quality for ruminants, pigs, poultry, and aquaculture
- Politics, markets, environment, acceptance (e.g. GMO)

7. ECONOMY AND MARKET

- Global comparison of OSR/canola farm economy
- Optimizing farm economy with OSR/canola: Australia, Canada, China and Europe
- Global status of genetically modified or genome edited OSR/canola
- Global markets of OSR/canola oil (incl. biodiesel), meal and protein
- Sustainability of OSR/canola production

8. MUSTARD AND OTHER CRUCIFEROUS OILSEED CROPS

9. OTHER TOPICS

Hubertus Paetow

President of DLG,
Frankfurt, Germany

The DLG was founded in 1885 with the goal to implement scientific and economic knowledge, technical novelties and organizational competence into practice. Above 130 years later, nearly 3,000 voluntary and 250 full-time employees just as more than 30,000 members still have the same goal: Impulses for progress. Despite the long persistence, progress is not a rigid ritual for the DLG, but rather a goal that has to be redefined to survive.

In keeping with this tradition, we did that again in 2017. With the ten theses "Agriculture 2030", we have faced the challenges of feeding a growing world population while protecting natural resources and respecting social goals. At the same time, for agriculture, this means increasing productivity, reducing environmental damage, and organizing arable and livestock farming in ways that are supported by a broad social consensus.

For arable farming, this implies that the focus must be return on a holistic approach and moving away from the falsely emphasis on short-term optimization of the profit margin. This requires site-adapted designing crop rotation using a well-filled toolbox.

In the area of rapeseed cultivation, Germany is second in the EU ranking, behind France, with 1.225 million ha in 2018 (AMI 2018), which corresponds to about one-ten of the arable land. Rapeseed is an all-rounder in use, as it fills the plate, tank and trough, almost without competition, because main and side products complement each other. Of the 3.7 million harvested tonnes, around 1.5 million tonnes of rapeseed oil, 2.2 million tonnes of rapeseed cake and extraction meal are produced. 0.5 million t of rapeseed oil are annually used in human nutrition.

But even in terms of arable farming, rape has hardly any other crop, despite or precisely because he is not self-sufficient and thus can be only cultivate in a wide crop rotation. He has highest demands on the soil and the supply of nutrients. Rape is particularly important in crop rotation with a high cereal content, as it promotes the structure and biological activity of the soil as well as the humus content through roots and remaining straw. Winter rape, for example, can take up released amounts of nitrogen in the autumn.

However, the high standards also include a high level of occupational and crop protection intensity. Especially here, the toolbox is restricted by unilateral prohibitions in the field of neonicotinoids and thus repealed the balance, especially since there are no alternatives. At the same time, rape cultivation within the EU is characterized by very volatile prices. Reasons for this are strong international competitors, including other oil-producing plants. In addition, extreme weather conditions, as in the past year, make cultivation even more difficult. According to the German Federal Statistics Office, only 0.917 million ha were drilled with rap in 2018, which corresponds to a decline of 25% compared to autumn 2017.

Possibilities that counteract the described trend are the expansion of the toolbox for example with mechanical weed control, which involves spaced planting and precision farming technology, thus combining old-fashioned and innovative methods. An opportunity in the marketing is to replace imported soya by rapeseed cake as an energy and protein supplier.

In order to stop the decline in cultivation due to restrictions in fertilization and crop protection as well as the increase in crop rotation diseases, emphasis is needed on research, breeding and testing for alternatives in crop protection. Furthermore it takes also new editions of pest damage thresholds, breeding against diseases and fertilization experiments in combination with growth models based on weather forecasts. A wide range of measures is the

Željko Milovac

Filip Franeta, Snežana Pešić, Petar Mitrović, Velimir Radić, Ana Marjanović Jeromela

Institute of Field and Vegetable Crops, Novi Sad, Serbia

Background: The abundance of rape stem weevil, *Ceutorhynchus napi* Gyll., and cabbage stem weevil, *C. pallidactylus* Marsh., in northern Serbia vary considerably depending on the production season and locality. The damage these two species cause on oilseed rape depends on various factors, among which mostly on their abundance.

Methods: The monitoring of these pests was conducted in northern Serbia, in the vicinity of Novi Sad, at Rimski Šančevi (N 45° 19' 51.03" ' E 19° 50' 07.16" ' , 80 m a.s.l.) using yellow water traps (Moericke dishes). Chemical control against stem weevils was performed regularly at the monitored site. Assessments were done every seven days from the beginning of the year until May during three consecutive years, from 2016 to 2018.

Results: The highest number of weevil specimens (153) was collected in 2017, while a total of 40 and 34 specimens were collected in 2016 and 2018 respectively. During the monitoring period, 2016 to 2018, the rape stem weevil was more abundant than the cabbage stem weevil (128:99), which is not in accordance with the data obtained for the same site for the 2011-2013 period, when an approximately equal species proportion was recorded. The sex ratio for both stem weevil species was approximately 1:1. The first adult specimens of rape and cabbage stem weevils were trapped at the beginning of March in 2016, in the last week of February in 2017. In 2018 the first adults of the cabbage stem weevil were collected on the 12th of January and on the 12th of February for the rape stem weevil.

Conclusions: During the monitoring period (2016-2018), although the total number of collected specimens was not high, it was observed that the rape stem weevil was more abundant than the cabbage stem weevil (1.29:1). The likely dominance of the rape stem weevil has been known for the area 100 km north of the experimental site but not for this locality. It seems plausible to assume that a shift in species dominance is occurring, however this idea should be verified. A significant influence of weather conditions on the beginning of the early year activity of both species and even their abundance is obvious.

Imprint

Please excuse any misspelling or grammatical errors that may occur in the book of abstracts. The book of abstracts contains data from diverse sources. The IRC-Team has requested clearance for all presentations.

Publication date

07|06|2019

IRC 2019

c/o Union zur Förderung von Oel-
und Proteinpflanzen e. V. (UFOP)

Claire-Waldoff-Straße 7

10117 Berlin (Germany)

Phone +49 30 31904-202

Fax +49 30 31904-485

Email info@irc2019-berlin.com

Picture credits:

- Cover © CCat82/Shutterstock.com