



PROCEEDINGS OF THE

23rd International Symposium on Analytical and Environmental Problems

October 9-10, 2017

**University of Szeged, Department of Inorganic and
Analytical Chemistry**



***Szeged
Hungary***

Edited by:
Tünde Alapi
István Ilisz

Publisher:
University of Szeged, Department of Inorganic and Analytical
Chemistry, H-6720 Szeged, Dóm tér 7, Hungary

ISBN 978-963-306-563-1

2017.
Szeged, Hungary

***The 23rd International Symposium
on Analytical and Environmental Problems***

Organized by:

SZAB Kémiai Szakbizottság Analitikai és Környezetvédelmi Munkabizottsága

Supporting Organizations

*University of Szeged, Department of Inorganic and Analytical Chemistry
Hungarian Academy of Sciences*

Symposium Chairman:

István Ilisz, PhD

Honorary Chairman:

Zoltán Galbács, PhD

Organizing Committee:

István Ilisz, PhD

associate professor

University of Szeged Department of Inorganic and Analytical Chemistry

ilisz@chem.u-szeged.hu

Tünde Alapi, PhD

assistant professor

University of Szeged Department of Inorganic and Analytical Chemistry

alapi@chem.u-szeged.hu

FIBER HEMP RESPONSE TO FOLIAR APPLICATION OF GROWTH REGULATORS

Biljana Kiprovski¹, Anamarija Stojanović¹, Vladimir Sikora¹, Tijana Zeremski¹, Bojan Konstantinović², Dragana Latković^{2,3}, Senka Vidović³

¹*Institute of Field and Vegetable Crops, Makisma Gorkog 30, 21000 Novi Sad, Serbia*

²*University of Novi Sad, Faculty of Agriculture, Trg Dositeja Obradovića 8, 21000 Novi Sad, Serbia*

³*University of Novi Sad, Faculty of Technology, Bulevar Cara Lazara 1, 21000 Novi Sad, Serbia*

e-mail: bkiprovski@gmail.com

Abstract

The aim of this paper was to test the effect of three plant growth regulators (PGR) on fiber content of hemp cultivar Helena. Applied PGRs, Cycocel[®], Regalis[®] and Moddus[®] had impact on fiber content in hemp cultivar Helena in medium concentrations: 1-2, 1.25-2.5 and 0.4-0.5 kg/ha, respectively. Further analysis of fiber quality, after application of various PGRs concentrations, should explain which PGR should be used in hemp fiber production.

Introduction

Plant growth regulators (PGR) are synthetic compounds used to regulate/retard the shoot length of plants. This is achieved by reducing cell elongation and by lowering the rate of cell division [1]. Most plant growth retardants inhibit the formation of active gibberellins and can be used to reduce shoot elongation [2]. Cycocel[®] (chlormequat chloride) is a synthetic growth retardant that is extensively used for dwarfing of plants. Studies have indicated that Cycocel[®] is effective in stimulating the production of secondary metabolites [3]. Regalis[®] (prohexadione-Ca or 3-oxido-4-propionyl-5-oxo-cyclohexenecarboxylate), primarily used for the control of shoot growth in pome and other fruit trees, is a multifunctional plant bio-regulator that mimic of 2-oxoglutarate, inhibiting dioxygenases, which catalyse distinct steps in the biosynthesis of the growth hormone gibberellin [1]. Moddus[®] (trinexapac-ethyl) is effective anti-gibberellin PGR marketed primarily for the control of lodging.

Plants cells sense the environment and make appropriate adjustments to gene expression, metabolism and physiology using oxidative stress or oxidative signaling, which could be regarded as interactions between reactive oxygen species (ROS) and the antioxidative system, which set signal intensity, to relay the oxidative message and to determine its physiological outcome [4].

Hemp cultivar Helena is primarily grown for grain production, however it is rich in fiber and can reach up to 4 m height. To achieve complete use of the plant, it is important to understand in which way it is possible to enhance yield of its usable parts.

The purpose of this work was to determine response of hemp cultivar Helena to foliar application of different concentrations of mentioned PGRs with a view to test their impact on plant membrane integrity and the fiber yield.

Experimental

Plants were grown in the experimental field of the Institute of Field and Vegetable Crops (Novi Sad, Serbia). Four concentrations of growth regulators, previously tested on other plant

species, were applied: Cycocel[®] (C1: 0.5, C2: 1, C3: 2, C4: 3 kg/ha), Regalis[®] (R1: 1, R2: 1.25, R3: 2.5, R4: 3 kg/ha) and Moddus[®] (M1: 0.2, M2: 0.4, M3: 0.5, M4: 0.8 kg/ha). Plants were treated at the beginning of a period of intensive vegetative growth, after which samples for biochemical analysis were collected. Quantification of investigated biochemical parameters was performed using Perkin Elmer Lambda 25 UV/Visible spectrophotometer. One g of fresh leaves was ground with cooled mortar and pestle and then homogenized with 5 ml 20% trichloroacetic acid for lipid peroxidation determination (LP) or total amount of TBARS (TBA-reactive substances) and expressed as nmol malondialdehyde (MDA) equivalents/g fresh weight (fw). Malondialdehyde (MDA) is a secondary end-product of the oxidation of polyunsaturated fatty acids, and reacts with thiobarbituric acid (TBA) to yield a pinkish-red chromogen with maximal absorbance at 532 nm [5]. Correction for non-specific turbidity at 600 nm was measured due to interference of other compounds including sugars, anthocyanins and other phenolics. Fiber yield [6], as well as plant heights and width were recorded. All statistical analysis were performed by STATISTICA for Windows version 13 (Dell Software).

Results and discussion

There is no registered product that is used for hemp growth regulation, the reason of which three commonly applied PGRs are used in this study. As it is showed in the Figure 1, control represented 100% of lipid peroxidation intensity, and plants reacted differently to various PGRs concentrations. Lipid peroxidation products such as MDA (malondialdehyde) are considered useful and reliable indicator of membrane damage and oxidative signalling, due to the susceptibility of membranes to reactive oxygen species (ROS). Only higher concentrations of Cycocel[®] had impact on hemp leaves, while Regalis[®] and Moddus[®] induced oxidative stress in lower concentrations (Figure 1).




Figure 1. Lipid peroxidation intensity in leaves of hemp 'Helena' treated with growing concentrations of growth regulators (C-cycocel, R-regalis and M-modus, concentrations in Experimental section). Results expressed as percent of control, control represents 100%.

Shoot height (2.5-3 m) and width (approx. 2 cm) were unaffected by the treatment and however were the differences among treatments, there was no correlation between shoot

length and width parameters and fiber content. Higher fiber content was recorded for the second and third concentration of all tested PGRs.

Table 1. Fiber content of ‘Helena’ plants treated with growing concentrations of growth regulators (concentrations in Experimental section).

% fiber content	C - Cycocel [®]				R - Regalis [®]				M - Moddus [®]						
	contr ol	C 1	C 2	C 3	C 4	contr ol	R 1	R 2	R 3	R 4	contr ol	M 1	M 2	M 3	M 4
	34	33	35	35	33	32	33	35	35	33	33	31	32	33	32

Conclusion

Applied PGRs had impact on fiber content in hemp cultivar Helena. Further analysis of fiber quality after application of various PGRs concentration should explain which PGR should be used in hemp fiber production.

References

- [1] W. Rademacher, *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 51 (2000) 501.
- [2] A.K. Singh, *Indian J. Agr. Sci.* 74 (2004) 130.
- [3] A.H.A. Farooqi, S. Fatima, A. Khan, S. Sharma. *Plant Growth Regul.* 46 (2005) 277.
- [4] H.C. Foyer, G. Noctor, *Plant Cell Environ.* 28.8 (2005) 1056.
- [5] D.M. Hodges, J.M. DeLong, C.F. Forney, R.K. Prange, *Planta* 207(4) (1999) 604.
- [6] G. Bredeman, *Faserforschung* 16 (1942) 14-39.