

Variation in quality traits of newly developed Serbian wheat cultivars under different environmental conditions of Pannonian plain

Dragan Živančev,¹ Milan Mirosavljević,¹ Vladimir Aćin,¹ Vojislava Momčilović,¹ Sanja Mikić,¹ Aleksandra Torbica,² Bojan Jocković¹

¹Institute of Field and Vegetable Crops, Novi Sad; ²Institute of Food Technology, University of Novi Sad, Novi Sad, Serbia

Highlights

- The modern wheat cultivars showed improved grain yield but lower technological quality than widespread cultivars.
- The growing seasons had the highest effect on bread volume and extensograph properties.
- Grain yield was negatively related to protein content and bread volume.

Abstract

The introduction of new high-yielding wheat cultivars and the intensification of high year-to-year variability require an additional update of information about environmental effects on main wheat quality parameters lacking for newly released cultivars in Serbia. This study aimed to determine the effect of growing seasons on the technological properties of new Serbian wheat cultivars and examine the relationships among the quality traits in different growing seasons. The experiment was set up in three growing seasons (2016-17, 2017-18, and 2018-19). A total of 13 recent-

ly developed wheat cultivars in Serbia and the five most widespread local cultivars were evaluated to compare differences in end-use quality. Also, the influence of climatic conditions on technological quality was examined and relations among them. Modern Serbian cultivars have improved grain yield but lower technological quality than the widespread Serbian cultivars. Changes in farinograph resistance and degree of softening were related to the differences in growing seasons, cultivars, and their interaction, while growing seasons had the highest effect on bread volume and extensograph parameters energy and ratio resistance, and extensibility variation. Moreover, a strong positive correlation was found between protein content, sedimentation value, wet gluten content, water absorption with extensograph extensibility, and negative correlation with the second parameter of extensograph, resistance ratio, and extensibility.

Correspondence: Dragan Živančev, Institute of Field and Vegetable Crops, Maksima Gorkog 30, Novi Sad 21101, Serbia.
Tel.: +38.1214898209. E-mail: dragan.zivancev@ifvcns.ns.ac.rs

Key words: Wheat; cultivar; environment; quality traits.

Funding: this research was supported by the Ministry of Education, Science and Technological Development of the Republic of Serbia, grant numbers: 451-03-68/2021-14/200032 and based upon the work from COST Action 18101 SOURDOMICS - Sourdough biotechnology network towards novel, healthier and sustainable food, and bioprocesses.

Contributions: the authors contributed equally.

Conflict of interest: the authors declare no potential conflict of interest.
Received for publication:

Received for publication: 18 May 2021.
Revision received: 12 September 2021.
Accepted for publication: 19 September 2021.

©Copyright: the Author(s), 2022
Licensee PAGEPress, Italy
Italian Journal of Agronomy 2022; 17:1911
doi: 10.4081/ija.2021.1911

This article is distributed under the terms of the Creative Commons Attribution Noncommercial License (by-nc 4.0) which permits any non-commercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

Introduction

Wheat is the most important winter cereal crop in the world. Furthermore, it is one of the most widely cultivated field crops in Europe and Serbia, with average 2014-2018 production ranging from 254,716.6 to 2583.6 thousand t, respectively (<http://www.fao.org/faostat/en/#data/FBS>). However, in the last decade in Europe, a fluctuation in wheat yields has been recorded from 3.5 t ha⁻¹ to 4.4 t ha⁻¹ (<http://www.fao.org/faostat/en>). Similarly, in the same period, the average grain yield in Serbia varied notably from 3.4 t ha⁻¹ in 2010 to 4.4 t ha⁻¹ in 2019 (Statistical Office of the Republic of Serbia, <https://data.stat.gov.rs/Home/Result/130102?languageCode=en-US>), showing remarkable instability. Next to the high year-to-year grain yield variability, several studies (Bilgin *et al.*, 2016; Mastilović *et al.*, 2018; Liu *et al.*, 2019) also confirmed a significant fluctuation of bread-making quality of wheat cultivars among different growing seasons. Furthermore, Fleitas *et al.*, (2020) reported that one of the aims is to identify which cultivars from a broad set possess both high yielding potential and stable bread-making quality under heat stress. Since most of the produced Serbian wheat is intended for local milling, baking sector, and export, it is necessary to ensure stable production of high-quality wheat.

The grain quality of winter wheat is strongly influenced by environmental conditions as extreme weather events negatively

affect starch and storage protein metabolism (Tomás *et al.*, 2020). Different environmental factors, such as temperature, amount and distribution of precipitation, the occurrence of heat and drought stress periods during plant development and grain filling period, have significant effects on wheat grain yield and quality (Balla *et al.*, 2011; Malik *et al.*, 2012; Dencic *et al.*, 2013; Kondić-Špika *et al.*, 2019). During the anthesis and grain-filling period, heat and drought stress significantly decrease wheat grain yield by reducing grain number per plant and grain weight (Akter *et al.*, 2017). Also, heat stress, temperatures above 35°C during grain filling, harm gluten extensibility (Hernández-Espinosa *et al.*, 2018), although some genotypes retain high end-use quality. In addition, a large amount of precipitation before harvest can trigger pre-harvest sprouting, and even a small amount of germinated grains can cause a significant reduction of dough strength (Barbeau *et al.*, 2006).

Different climate scenarios predict a rise of the average temperature by 2.0-4.9°C until the end of this century due to climate changes (Asseng *et al.*, 2015), and more frequent dry periods, heat stress, and heavy rainfall (Trnka *et al.*, 2014). Among the main wheat production regions in Europe, the Pannonian Plain has been identified as one of the most sensitive regions to the influence of further climate changes (Olesen *et al.*, 2011). Therefore, predicted climate variability will further disturb quality wheat's high and stable production.

Apart from the environmental influence, wheat grain yield and quality are controlled by genotype, production management, and interaction (Rozbicki *et al.*, 2015; Laidig *et al.*, 2017). However, the influence of these factors on grain quality is inconsistent. Some traits, such as Zeleny sedimentation or flour yield, are more related to the genotypic effect (Carson and Edwards, 2009), while others, such as falling number or wet gluten, are under higher environmental control (Kong *et al.*, 2013). Moreover, genotype by environment interaction strongly influences wheat quality, and variable genotypic response to different environments further complicates the production of high-quality wheat (Hristov *et al.*, 2010; Rozbicki *et al.*, 2015). However, in some long-term studies (Laidig *et al.*, 2017), genotype by environment interaction was not so pronounced. Maintaining a high quality of wheat cultivars across different growing seasons is crucial for human nutrition and wheat breeders, producers, and the processing industry (Denčić *et al.*, 2012).

In addition, grain quality is defined by several factors, such as physical and rheological traits strictly defined by end-users from different markets. The protein content is a primary indicator of wheat quality, strongly affected by environmental factors (Malik *et al.*, 2013). Protein quality is highly related to the composition of gluten molecules and its main fractions, gliadins and glutenin subunits (GS) - high molecular weight (HMW) and low molecular weight (LMW), the specific composition of which is mainly determined by genotype (Johansson *et al.*, 2013). Among them, HMW-GS are mostly related to the dough rheological properties of all proteins and their specific composition has a significant role in the end-use quality of wheat. The difference in dough rheological properties is that the result of different compositions of the HMW-GS amino acid sequences possesses from four to seven cysteine residues at the N-terminal and C-terminal end (Shewry *et al.*, 1992). Also, subunit HMW-GS 5 encoded by the D loci possesses an extra cysteine residue at the N-terminal end absent in the HMW-GS 2 subunit encoded by the same loci (Gras *et al.*, 2001). Liang *et al.* (2010) showed that LMW-GS also affected the end-use quality and acted as polymer chain terminators (Schmid *et al.*, 2016).

Furthermore, Guo *et al.* (2018) recently showed that gliadins increase the viscosity of gluten molecules and decrease glutenin

elasticity resulting in actually softer gluten. In general, dough strength and elasticity are determined by glutenin. Furthermore, different important wheat rheological traits are closely related to the content and quality of wet gluten and sedimentation value (Yang *et al.*, 2014). Previous studies about the effects of genotype by environmental interaction on quality traits focused mainly on a limited set of generally outdated cultivars (Zečević *et al.*, 2009; Živančev *et al.*, 2016; Kondić *et al.*, 2020). The introduction of new high-yielding wheat cultivars and the intensification of high year-to-year variability require additional information about environmental effects on main wheat quality parameters. Therefore, the main objectives of this study were: i) to examine the effect of growing seasons on the leading quality traits of thirteen newly released wheat cultivars in Serbia and five of the most commercially widespread wheat cultivars; and ii) to analyse the relationships among the grain, flour, dough and baking traits across different growing seasons.

Materials and methods

Cultivars and experimental design

The study was conducted at the experimental field of the Institute of Field and Vegetable Crops, Novi Sad, Serbia (45°20'N and 19°51'E), during three growing seasons (2016-17, 2017-18 and 2018-19). The location is characterised by haplic chernozem aric soil (IUSS Working Group WRB, 2014) and a long-term three-year crop rotation of small grains, maize, and legume crops. A total of 18 winter wheat cultivars developed at the Institute of Field and Vegetable Crops were selected for this study (Table 1). The main criterion for cultivar selection was to include five of the most commercially widespread cultivars (Simonida, NS 40S, Zvezdana, NS Futura, and NS Ilina), as well as newly released high yielding cultivars (NS Mila, NS Obala, NS Javorka, NS Moma, NS Tajna, NS Jevrosima, NS Klica, NS Rani otkos, NS Vljajna, NS Epoha, NS Igra, NS Grivna, and NS Todorka). The allelic composition of the high molecular weight glutenin subunits (HMW-GS) in the studied cultivars was obtained by the procedure described in Živančev *et al.* (2016) using chip electrophoresis technique on an Agilent 2100 Bioanalyzer (Agilent Technologies, Santa Clara, CA, USA) with Protein 230 Lab-on-a-Chip kit for separation of proteins and presented in Table 1.

The experiment was set up as a completely randomised block design in three replications on plots 10 m long and 1 m wide with 10 rows per plot. Crops were sown on 17 October 2016, 23 October 2017, and 18 October 2018, providing the recommended sowing density for wheat production in southern Pannonian Plain of 500 plants per m². Basic soil properties were determined from the soil samples taken in each growing season from the topsoil layer (0-30 cm) before ploughing (Table 2). Based on these analyses, average doses of 60 kg ha⁻¹ of phosphorus (P) and 60 kg ha⁻¹ of potassium (K) were applied before sowing, along with 60, 45, and 50 kg of nitrogen (N) ha⁻¹ in 2016, 2017, and 2018, respectively. In addition, to avoid N deficiency during intensive plant growth, ammonium nitrate (34% N) was top-dressed based on the N-min analysis (Wehrmann and Scharpf, 1979) in early spring (Table 2). During the growing seasons, pests, diseases, and weeds were controlled by recommended application of chemicals when required.

Grain, flour and bread quality traits

Plots were combine-harvested at the grain ripening stage (late June), and grain yield (GY) was calculated at a 13% moisture level. After harvest, the samples were threshed and cleaned to remove the remaining parts of stalks and chaffs. The protein content (PC) in grain was determined according to the Kjeldahl method by Kejltec 2300 (Foss, Hillerød, Denmark) by AACC (2000) method 46-10 and expressed on a grain dry weight basis. Wheat samples were tempered for 24 h prior to milling to 13.5% moisture. Half an hour before milling, samples were prepared to the final tempering moisture of 15% and milled by a Bühler laboratory mill MLU-202 (Uzwil, Switzerland). Wet gluten content

(WG) was obtained according to the ICC methods 137/1 using Glutomatic 2200 (Perten Instruments, Huddinge, Sweden). Falling number (FN) was determined with a Falling Number 1800 instrument (Perten Instruments, Huddinge, Sweden), according to Hagberg, using the 107/1 method (ICC 2011). The sedimentation value (SV) of wheat samples was determined according to AACC (2000) method 56-62.01. Dough properties water absorption (WA), farinograph resistance (FR), and softening degree (SD) were obtained using a farinograph (Brabender OHG, Duisburg, Germany), according to the Hungarian Standard method 6369/653 (MSZ 1988). Extensograph energy (E), extensibility (EE), and resistance/extensibility ratio (R/EE) were measured on an extenso-

Table 1. The analysed cultivars, year of release, and allelic composition at the high molecular weight glutenin subunits with GLU score.

Cultivar	Year of release	HMW-GS			GLU score
		Glu-A1	Glu-B1	Glu-D1	
Commercially widespread cultivars					
Simonida	2003	N	7+9	2+12	5
NS 40S	2006	N	7+9	5+10	7
Zvezdana	2005	N	7+9	2+12	5
NS Futura	2010	2*	7+9	5+10	9
NS Ilina	2010	N	7+8	5+10	8
Newly released high yielding					
NS Mila	2014	N	7	5+10	6
NS Obala	2015	N	7+9	5+10	7
NS Javorka	2015	N	7+9	5+10	7
NS Moma	2016	N	7+9	2+12	5
NS Tajna	2016	N	7+9	5+10	7
NS Jevrosima	2016	2*	7+9	2+12	7
NS Klica	2016	2*	7+9	5+10	9
NS Rani otkos	2017	2*	7+9	2+12	7
NS Vlajna	2014	N	7+9	5+10	7
NS Epoha	2016	2*	7+8	5+10	10
NS Igra	2016	2*	7+9	3+12	7
NS Grivna	2016	2*	7+9	5+10	9
NS Todorka	2016	2*	7+9	2+12	7

HMW-GS, high molecular weight glutenin subunits.

Table 2. Basic soil properties and amounts of applied nutrients.

	2016/17	2017/18	2018/19
Soil properties			
pH (KCl)	7.33	7.17	7.41
CaCO ₃ (%)	5.17	5.08	4.85
P ₂ O ₅ (Egner-Riehm, AL method) (mg 100g ⁻¹)	24.5	28.4	20.8
K ₂ O (Egner-Riehm, AL method) (mg 100g ⁻¹)	18.3	19.1	17.6
Humus (Tyurin method) (%)	2.7	2.9	2.5
Available N (N-min method) (kg ha ⁻¹ ; 0-90 cm)	80	70	50
Fertilisation			
P ₂ O ₅ (kg ha ⁻¹)	60	50	70
K ₂ O (kg ha ⁻¹)	60	50	70
N (kg ha ⁻¹ prior to sowing)	60	45	50
N (kg ha ⁻¹) topdressing (in spring)	80	98	120

graph (Brabender OHG, Duisburg, Germany) according to the 114/1 method (ICC 2011), while Alveograph deformation energy (W) was obtained using the alveograph (Chopin, Paris, France) according to the 121 method (ICC 2011). The bread volume (BV) was measured by displacement method with millet, 24 h after baking, according to the internal procedure for evaluating wheat cultivars (Kaluderski and Filipović, 1998).

Weather conditions

The weather data for October to April in 2016-2017, 2017-2018, and 2018-2019 were obtained from the meteorological station located near the experimental field (Figure 1).

Weather conditions during flowering and grain-filling stages, which in Serbia occur in May and June, respectively, are crucial for good grain quality performance. For this reason, the average temperatures and rainfalls were recorded daily starting from the flowering time (Figure 2A-D).

Statistical analysis

The studied wheat quality parameters were subjected to analysis of variance (ANOVA) using Infostat (Di Rienzo *et al.*, 2011). ANOVA was performed using a generalized linear model (GLM), including environments (growing seasons - GS), replications within environments, blocks within environments, and their interactions. The variance components were calculated to estimate the share of the factors and their interaction in the total variation of studied parameters. Linear regression and Pearson's correlation were calculated by averaging three replicates in the software InfoStat (Di Rienzo *et al.*, 2011).

Results and discussion

Mid-daily air temperatures in October and November did not vary considerably among the growing seasons (Figure 1). On the other hand, precipitation amounts varied among growing seasons in autumn, when high, normal, and low rainfall were measured in 2016-17, 2017-18, and 2018-19, respectively. The winter period in

2016-17 was characterised by the lowest average monthly temperatures and low rainfalls, while 2018-19 and 2017-18 were mild with moderate rainfall levels. February and March coincided with the mid- and late-tillering period and were relatively drier in 2016-17 and 2018-19 than in 2017-18. Plants were in the stem elongation stage in April. All seasons were characterised by high rainfall and moderate temperatures, except 2017-18, which had a higher average monthly temperature. The season 2017-18 was characterised by relatively higher temperatures during the first half of the grain filling period, while the lowest daily temperatures were recorded in 2018-19 (Figure 2). The second half of the grain filling period was characterised by high day-to-day temperature variability in 2017-18, relatively high daily temperatures in 2017-18 and 2018-19, and moderate daily temperatures in 2016-17. Daily precipitation was unequally distributed during the grain-filling period. There were several rainy days during the mid-grain filling period in 2016-17 (Figure 2B), while the best rainfall distribution was observed in 2018-19 (Figure 2D).

The results showed that almost half of grain yield (GY) variation (45.7%) was under genotypic effect (C), while the influence of year by cultivar interaction ($Y \times C$) and year (Y) explained 32.4% and 21.9% of the total variation, respectively (Table 3). Although Serbian cultivars used in this study have been released in the last 15 years, a high percentage of total variation explained by genotypic effect indicated diverse pedigrees of the cultivars reflected in differences in yield performance. The first six high-yielding cultivars were newly released, whereas four of the five most commercially widespread wheat cultivars (Simonida, Zvezdana, NS Futura, and NS Ilina) ranked from 12 to 15. Across the growing seasons, Serbian wheat cultivars achieved the highest grain yield in 2017-18, while the lowest GY was reported in 2016-17 (Table 4). According to Figure 1, 2017-18 was characterised by a high amount of precipitation that was equally distributed in spring, favouring crop growth and development. In small grain cereals, weather conditions during the pre-anthesis and anthesis periods are crucial for grain number and grain yield determination, as previously described by Slafer *et al.* (2014).

Growing season (Y), C, and $Y \times C$ had a significant ($P < 0.01$) effect on PC (Table 3). More than half of the variation in PC was




Figure 1. Average monthly rainfall and temperature during pre-anthesis developmental phases in three growing seasons.

explained by the influence of Y, followed by the influence of C and $Y \times C$ interaction. PC ranged from 11.6% in 2017-18 to 13.8% in 2018-19, with the total average protein content of 12.8% (Table 4). Also, three of the five most commercially widespread wheat cultivars (Simonida, Zvezdana, and NS Futura) ranked among the first top-five based on PC. The analysed Serbian cultivars had higher PC on average than 20 most widely grown cultivars and advanced inbred lines from the International Winter Wheat Improvement Program in Turkey (Kaya and Akcura, 2014), and slightly lower PC than the average of 16 most widely grown cultivars in Spain (Sanchez-Garcia *et al.*, 2015). Moreover, a significant variation in PC across different growing seasons has been reported for other environments, such as the Mediterranean (Kaya and Akcura, 2014) or central Europe (Hlisenikovsky *et al.*, 2020). In our study, the lowest protein content was recorded in 2017-18, characterised by the

highest average grain yield. This is not surprising as a strong negative correlation between PC and GY is generally known (Da Silva *et al.*, 2014). During the grain filling period, the weather conditions had a significant effect on grain nitrogen accumulation and PC (Pan *et al.*, 2006). The second part of the grain filling period was characterised by higher temperatures in 2018-19 than in 2016-17 and 2017-18. High temperatures tend to shorten grain filling, thus decreasing grain carbohydrates accumulation in the developing grain and resulting in more nitrogen per grain weight (Stone and Nicolas, 1998). The highest average PC was recorded in the cultivar NS Todorka (13.9%, recently developed), followed by NS Jevrosima (13.7%, recently developed), both of which had the same combination of HMW-GS 2*, 7+9, and 2+12 with Glu score 7. On the other hand, the lowest PC was recorded in the cultivars NS Klica (11.6%, recently developed) and NS Grivna (11.9%,




Figure 2. Average daily temperatures during three growing seasons (A) and daily rainfall in 2016-17 (B), 2017-18 (C), and 2018-19 (D) after flowering.

Table 3. Relative contributions to the total sum of squares and significance level of source of variance for the studied traits of winter wheat.

Source	GY	PC	WG	SV	WA	FR	SD	FN	E	EE	R/EE	W	BV
Year	10.3**	69.7**	1090**	791**	134**	112**	27,613**	1,036,614**	26,502**	6296**	40.6**	42,850**	581,376**
Cultivar	2.5**	4.2**	146**	173**	29**	13**	1640**	24,488**	2761**	2195**	3.5**	7973**	50,413**
$Y \times C$	0.9**	0.8**	24**	31**	10 ^{ns}	11**	1366**	12,250**	464**	714**	1.3**	3567**	15,837**
Error	0.1	0.3	2	2	6	0.04	15	211	19	44	0.02	73	3436
% of SS													
Year	21.9	58.8	39.8	28.3	24.5	27.2	42.6	71.3	45.8	17.0	44.1	25.0	45.5
Cultivar	45.7	30.4	45.3	52.8	45.7	27.1	21.5	14.3	40.6	50.3	32.2	39.6	33.5
$Y \times C$	32.4	10.9	14.8	18.9	29.8	45.7	35.8	14.3	13.6	32.7	23.7	35.4	21.0

GY, grain yield; PC, protein content; WG, wet gluten content; SV, sedimentation value; WA, water absorption; FR, farinograph resistance; SD, softening degree; FN, falling number; E, extensograph energy; EE, extensibility; R/EE, resistance/extensibility ratio; W, alveograph deformation energy; BV, bread volume; Y, year; C, cultivar; $Y \times C$, year by cultivar interaction; SS, sum of squares. **Significant at the 0.01 probability level; ^{ns}, not significant.

recently developed), which were characterised by the same combination of HMW-GS 2*, 7+9, and 5+10 and with Glu score 9. This is in accordance with the results of Johansson *et al.* (2020) and Blumenthal *et al.* (1995), who showed that some wheat cultivars with HMW-GS 5+10 at the *Glu-D1* locus had lower values of technological parameters quality than cultivars with HMW-GS 2+12. Previous studies indicated higher variation in PC among different wheat cultivars, ranging between 10.8 and 15.7% in Romania (Liana *et al.*, 2012) or between 12 and 16% in Croatia (Horvat *et al.*, 2015). However, cultivars used in our study are mostly modern Serbian cultivars developed recently for intensive high-yielding production, showing less variability in PC. Generally, breeding activities in the past century resulted in the development of high-yield cultivars with decreased PC (Miroslavljević *et al.*, 2020), and such a trend was noted in our research.

Variations in WG and SV of the examined cultivars were mostly related to differences among genotypes and were followed by the influence of the growing season. This is in accordance with Laidig *et al.* (2017), who found that SV was highly influenced by cultivars (variation - 69%). Furthermore, Rozbicki *et al.* (2015) showed that in addition to the significant influence of genotype on WG, Y × C interaction significantly influenced the same parameter. The SV and WG content was higher in the years characterised by increased PC content. The highest SV was recorded in 2016-17, while WG was similar in 2016-17 and 2018-19. There was a high positive correlation of WG and SV with a PC (Tables 5 and 6). The SV ranged between 22.0 and 41.7 mL, while WG varied between

24.9 and 38.8% (Table 4). The highest SV and WG content were recorded in NS Todorka, while NS Mila was characterised by the lowest value of SV and WG, whereas Simonida and Zvezdana ranked among the first top six places based on these two parameters. Previously, Denčić *et al.* (2011) reported lower average WG content and its higher variation (21.3–40.6%) in winter wheat cultivars of different origins grown under conditions of the Pannonian Plain. On the other hand, Surma *et al.* (2012) reported that SV varied between 34.1–53.6 mL, mainly under environmental control in Poland.

Farinograph trait WA significantly varied due to the influence of Y and C, which was in accordance with the results of a few studies (Linina *et al.*, 2014; Rozbicki *et al.*, 2015). However, Y × C interaction did not significantly influence WA in the examined Serbian cultivars, which was not the case in the study Rozbicki *et al.* (2015). On the other hand, FR and SD were under the significant effect of Y, C, and their interaction, whereas in Rozbicki *et al.* (2015), this effect was not significant. Variation in WA of the examined Serbian cultivars was mostly related to the differences among genotypes, whereas SD was mostly related to the influence of growing season. On the other side, almost half of the variation of the FR parameter was explained by the influence of the Y × C interaction. During mixing, the best properties of dough were associated with the increased WA and FR and decreased SD. Among the examined Serbian cultivars, the WA ranged from 53.9 to 61.2%, FR varied between 2.3 and 7.3 min, and SD from 51.7 to 108.3 BU (Table 4). Two of the five most commercially

Table 4. Average values of grain yield (GY), protein content (PC), wet gluten (WG), sedimentation value (SV), water absorption (WA), farinograph resistance (FR), softening degree (SD), falling number (FN), extensograph energy (E), extensibility (EE), resistance/extensibility ratio (R/EE), alveograph deformation energy (W) and bread volume (BV) in 18 wheat cultivars during three growing seasons (2016-17, 2017-18 and 2018-19).

Cultivar	GY	PC	WG	SV	WA	FR	SD	FN	E	EE	R/EE	W	BV
Simonida	7.47 ^{c-e}	13.5 ^{ab}	35.2 ^a	36.2 ^b	58.8 ^{ab}	3.8 ^{fg}	78.3 ^{hi}	376 ^{ab}	120.7 ^a	173 ^a	2.1 ^{jk}	236 ^a	1433 ^a
NS 40S	7.75 ^{c-e}	12.1 ^{d-f}	23.3 ^g	30.7 ^{f-h}	56.7 ^{b-d}	2.3 ^k	108.3 ^a	260 ^g	133.2 ^a	140 ^{f-h}	4.1 ^a	134 ^b	1200 ^{gh}
Zvezdana	7.40 ^{c-e}	13.4 ^{ab}	34.8 ^{ac}	34.3 ^{b-d}	59.6 ^{ab}	5.3 ^a	75.0 ⁱ	353 ^{bc}	104.0 ^{c-e}	169 ^{aa}	1.9 ^{kl}	221 ^{b-d}	1413 ^{ab}
NS Futura	7.36 ^{c-e}	13.3 ^{ab}	32.6 ^{cd}	32.0 ^{d-f}	58.2 ^{a-c}	7.3 ^a	51.7 ^k	286 ^e	103.1 ^{c-e}	149 ^{c-f}	2.4 ^{hi}	224 ^{a-c}	1427 ^{aa}
NS Ilina	7.46 ^{c-e}	12.7 ^{b-e}	28.9 ^{ef}	31.2 ^{e-g}	56.9 ^{b-d}	2.7 ^{jk}	80.0 ^{g-i}	324 ^d	108.8 ^c	148 ^{c-g}	2.7 ^{fg}	181 ^{ef}	1353 ^{a-e}
NS Mila	7.30 ^{c-e}	12.0 ^{ef}	24.9 ^g	22.0 ^j	56.0 ^{b-d}	2.7 ^{jk}	83.3 ^{f-h}	332 ^{cd}	64.3 ^j	115 ^j	3.5 ^a	154 ^g	1247 ^{f-h}
NS Obala	7.54 ^{c-e}	12.6 ^{b-e}	28.9 ^{ef}	28.5 ^{hi}	56.2 ^{b-d}	3.5 ^{gh}	85.0 ^{e-g}	237 ^{sh}	102.4 ^{c-e}	138 ^{f-h}	3.2 ^{cd}	180 ^{ef}	1287 ^{c-g}
NS Javorka	7.19 ^e	12.6 ^{b-e}	30.3 ^{de}	28.3 ^{hi}	56.7 ^{b-d}	3.0 ^{ij}	83.3 ^{f-h}	380 ^a	90.9 ^{fg}	127 ⁱ	3.4 ^{ac}	207 ^d	1273 ^{d-h}
NS Moma	7.77 ^{cd}	13.1 ^{a-c}	33.8 ^{ac}	33.3 ^{c-e}	58.7 ^{ab}	4.3 ^{de}	90.0 ^{de}	340 ^{cd}	98.0 ^{ef}	157 ^c	2.6 ^{gh}	208 ^d	1360 ^{a-e}
NS Tajna	8.52 ^a	12.0 ^{ef}	27.6 ^f	27.5 ⁱ	53.9 ^d	2.8 ^{ij}	85.0 ^{e-g}	334 ^{cd}	85.3 ^{gh}	138 ^{g-i}	2.9 ^{ef}	176 ^f	1253 ^{f-h}
NS Jevrosima	7.48 ^{c-e}	13.7 ^a	35.0 ^{ac}	34.5 ^{bc}	58.4 ^{ab}	4.3 ^{de}	103.3 ^{aa}	229 ^h	99.9 ^e	159 ^{bc}	2.2 ^j	221 ^{b-d}	1380 ^{a-c}
NS Klica	8.37 ^{ab}	11.6 ^f	27.8 ^f	27.5 ⁱ	54.0 ^{cd}	4.0 ^{ef}	75.0 ⁱ	327 ^d	87.3 ^{gh}	131 ^{hi}	3.2 ^{cd}	155 ^g	1187 ^h
NS Rani otkos	7.28 ^{de}	13.2 ^{ab}	34.8 ^{ac}	36.0 ^b	58.0 ^{a-d}	4.9 ^c	97.1 ^{ac}	191 ⁱ	83.1 ^{hi}	154 ^{cd}	1.9 ^k	229 ^{ab}	1368 ^{a-d}
NS Vlajna	7.58 ^{c-e}	13.2 ^{ab}	33.5 ^{ac}	29.7 ^{f-i}	57.7 ^{a-d}	4.7 ^{cd}	87.4 ^{d-f}	284 ^{ef}	76.3 ⁱ	140 ^{f-h}	2.4 ^{ij}	176 ^{ef}	1333 ^{b-f}
NS Epoha	7.87 ^{bc}	13.0 ^{a-d}	32.9 ^{ac}	34.8 ^{bc}	58.0 ^{a-d}	4.2 ^{ef}	75.0 ⁱ	271 ^{ef}	126.0 ^{ab}	153 ^{c-e}	3.0 ^{de}	213 ^{cd}	1360 ^{a-f}
NS Igra	8.89 ^a	12.2 ^{def}	29.6 ^{ef}	30.7 ^{f-h}	56.3 ^{b-d}	4.3 ^{de}	61.7 ^j	328 ^d	107.8 ^{cd}	143 ^{e-g}	3.0 ^{de}	191 ^e	1280 ^{d-h}
NS Grivna	8.84 ^a	11.9 ^{ef}	29.0 ^{ef}	29.2 ^{g-i}	58.0 ^{a-d}	3.2 ^{hi}	93.3 ^{cd}	332 ^{cd}	83.2 ^{hi}	144 ^{d-g}	2.6 ^{g-i}	167 ^{fg}	1267 ^{e-h}
NS Todorka	7.70 ^{c-e}	13.9 ^a	38.8 ^a	41.7 ^a	61.2 ^a	4.8 ^c	85.0 ^{e-g}	275 ^{ef}	100.9 ^{de}	173 ^a	1.9 ^l	217 ^{b-d}	1367 ^{a-d}
Years													
2016/17	7.39 ^c	13.0 ^b	33.3 ^a	34.1 ^a	57.2 ^a	4.8 ^a	107.5 ^a	158 ^c	78.4 ^c	159 ^a	2.0 ^c	182 ^a	1441 ^a
2017/18	8.24 ^a	11.6 ^c	26.0 ^c	27.1 ^c	55.9 ^c	2.4 ^c	79.4 ^a	317 ^b	122.3 ^a	137 ^c	3.7 ^a	174 ^c	1255 ^b
2018/19	7.68 ^b	13.8 ^a	34.3 ^a	33.4 ^b	59.1 ^a	4.9 ^a	62.8 ^c	434 ^a	95.2 ^b	146 ^a	2.5 ^a	226 ^a	1269 ^b

Means with at least one common letter were not significantly different at P=0.05 by Tukey's test.

widespread wheat cultivars (Zvezdana and NS Futura) possessed the highest values of WA and FR. In addition, there were high positive relationships among protein parameters (PC and WG) with the following farinograph parameters WA and FR (Table 4). Also, the increase in WA with PC is in accordance with the one reported in Constantinescu *et al.* (2011). The results from other studies showed higher WA among different wheat cultivars with an average content of 61.8% (high level of nitrogen fertilisation) and 60.0% (low level of nitrogen fertilisation) in Croatia (Šarčević *et al.*, 2014) or between 58.2-65.1% in Lebanon (Sakr *et al.*, 2021), whereas SD was on a similar level.

Growing season (Y), cultivar (C), and $Y \times C$ interaction had a significant ($P < 0.01$) influence on FN (Table 3). This is similar to Rozbicki *et al.* (2015), who found that years and genotype significantly influenced the parameter, but not year \times genotype interaction. More than 70% of the variation in FN was explained by the influence of Y, followed by the influence of C and $Y \times C$ interaction. The second and third highest FN were recorded in Simonida and Zvezdana, belonging to the five most commercially widespread wheat cultivars. The lowest FN in the examined

Serbian cultivars was recorded in 2016-17 (158 s), which is relatively lower than 250 s minimum of optimal value. However, the period of grain ripening (forty days after anthesis) was not characterised by a high precipitation amount (Figure 1B). High fluctuations in day and night temperatures during grain maturation in 2016-17 could be a reason for premature increased production of the first isomorph of wheat α -amylase in a small percentage of the cells in the aleurone layer without visible germination (Mares and Mrva, 2008), the effect known as late maturity α -amylase activity (LMA) triggered by environmental extremes (Farrell and Kettwell, 2008). In the grain aleurone layer, the first isomorph of wheat α -amylase remains active, without any effect on kernel shape and size. On the other hand, the 2018-19 growing season was characterised by the highest FN (>400 s), indicating insufficient α -amylase activity in wheat grain. In baking formulations with high FN flour, it is necessary to add α -amylase for bread loaf volume improvement (Posner and Hibbs, 2005). The obtained FN values in 2017-18 and 2018-19 were similar to the results of Šarčević *et al.* (2014) and Sakr *et al.* (2021).

The extensograph traits E, EE, and R/EE and alveograph

Table 5. Correlation matrix of 18 wheat cultivars for studied quality traits in three growing seasons.

GS	2017	2018	2017	2018	2017	2018
	PC		SV		WG	
2018	0.675**	-	0.718**	-	0.597**	-
2019	0.592**	0.550**	0.599**	0.601**	0.763**	0.607**
	WA		FR		SD	
2018	0.278	-	0.033	-	-0.473*	-
2019	0.580*	0.359	0.041	0.356	0.230	0.563*
	E		EE		R/EE	
2018	0.671**	-	0.478*	-	0.283	-
2019	0.542*	0.657**	0.466	0.298	0.681**	0.214
	BV		FN		W	
2018	0.537*	-	0.336	-	0.065	-
2019	0.357	0.417	-0.058	0.411	0.150	0.576

GS, growing season; PC, protein content; SV, sedimentation value; WG, wet gluten content; WA, water absorption; FR, farinograph resistance; SD, softening degree; E, extensograph energy; EE, extensibility; R/EE, resistance/extensibility ratio; BV, bread volume; FN, falling number; W, alveograph deformation energy. *, **Significant at the 0.05 and 0.01 probability level, respectively.

Table 6. Correlation between studied quality traits of 18 wheat cultivars in averaged across growing seasons.

	PC	SV	WG	WA	FR	SD	E	EE	R/EE	BV	FN	W
SV	0.801**											
WG	0.895**	0.836**										
WA	0.841**	0.805**	0.796**									
FR	0.600**	0.482*	0.656**	0.492*								
SD	-0.014	0.049	-0.110	0.050	-0.531*							
E	0.240	0.492	0.082	0.265	0.022	-0.061						
EE	0.776**	0.923**	0.811**	0.794**	0.478*	-0.007	0.489*					
R/EE	-0.741**	-0.692**	-0.882**	-0.670**	-0.628**	0.114	0.121	-0.772**				
BV	0.889**	0.695**	0.824**	0.758**	0.661**	-0.269	0.246	0.784**	-0.796**			
FN	-0.275	-0.292	-0.154	-0.132	-0.232	-0.379	-0.072	-0.113	0.124	-0.085		
W	0.825**	0.723**	0.870**	0.655**	0.642**	-0.273	0.197	0.723**	-0.752**	0.885**	-0.063	
GY	-0.594**	-0.189	-0.298	-0.373	-0.194	-0.116	-0.023	-0.135	0.168	-0.481*	0.201	-0.397

PC, protein content; SV, sedimentation value; WG, wet gluten content; WA, water absorption; FR, farinograph resistance; SD, softening degree; E, extensograph energy; EE, extensibility; R/EE, resistance/extensibility ratio; BV, bread volume; FN, falling number; W, alveograph work; GY, grain yield; *, **Significant at the 0.05 and 0.01 probability level, respectively.

parameter *W*, also significantly varied due to the influence of *Y*, *C*, and their interaction. However, influence varied among these extensibility traits since *E*, *EE*, *R/EE*, and *W* were explained mainly by the influence of *C*, *Y*, and *Y × C* interaction, respectively. Variation in *E* and *R/EE* of the Serbian cultivars was mostly related to the differences between growing seasons, whereas *EE* and *W* were mostly related to the influence of cultivars. This is similar to Hernández-Espinosa *et al.* (2018), who reported that genotype variation had the highest contribution to *W* than the environment, year, and all interaction. Contrary to our findings, the genotype predominantly influenced the *E* and *R/EE* of Chinese cultivars (Kun *et al.*, 2020). The best extensibility dough properties were in 2018-19, since the value of *R/EE* was in the optimum range between 1.5 and 2.5 (Kovačević, 2011), and *W* was the highest. The *E* ranged from 78.4 to 122.3 cm², while *EE* varied between 137 and 159 min, and *R/EE* from 2.0 to 3.7. Also, *W* varied from 176 to 226 J $\dot{\text{I}} 10^{-4}$ (Table 4). Moreover, Simonida, Zvezdana, and NS Futura possessed a statistically higher value of *W* in comparison to the most 13 recently developed wheat cultivars except for NS Javorka, NS Moma, and NS Jevrosima. The average results from the study of Šarčević *et al.* (2014) showed higher *EE* and lower *R/EE* among different wheat cultivars, especially in case of a high level of nitrogen fertilization - 190 mm and 1.3, respectively. Also, the values of alveograph parameter *W* were somewhat lower than the Lebanese wheat cultivars (Sakr *et al.*, 2021).

Parameters *Y*, *C*, and *Y × C* had a significant ($P < 0.01$) influence on bread volume (*BV*) (Table 3). However, Rozbicki *et al.* (2015) found that only *Y* had a significant ($P < 0.05$) influence on bread volume *BV*. Almost half of the variation in *BV* of the Serbian cultivars was explained by the influence of *Y*, followed by the influence of *C* and *Y × C* interaction. Contrary to our findings, in Hernández-Espinosa *et al.* (2018), more than half of the variation in *BV* was explained by the influence of genotype, followed by the influence of year and year \times genotype interaction. The highest *BV* was recorded in 2016-17 (1441 cm³) and was statistically higher than in 2017-18 (1255 cm³) and 2018-19 (1269 cm³). According to Ral *et al.*, (2016), there were no results about LMA's detrimental effect on the end-product quality. Although the best rheological properties were recorded in 2018-2019, they could not substantially affect *BV* since our test baking method contained no sugar or addition of malt, regardless of the $\text{FN} > 400$ s high value. Furthermore, the three highest *BV* were recorded in Simonida, Zvezdana, and NS Futura.

In order to determine whether some traits are stable across the environments, correlation coefficients were made between all growing seasons (Table 5). There were strong positive correlations between *PC*, *SV*, *WG*, and *E* parameters among all three growing seasons. However, in the case of *WA*, *FR*, *SD*, *FN* parameters and correlations exist only between some growing seasons. This indicates that *WA*, *FR*, *SD*, *FN*, and *W* parameters depended on weather conditions in the growing seasons.

The two most important wheat parameters *GY* and *PC*, of the examined Serbian cultivars, were in a strong negative correlation (Table 6). This was in agreement with the results of Marinciu and Săulescu (2008) and Da Silva *et al.* (2014). Also, *PC*, *SV*, and *WG* were strongly positively correlated with *WA*, according to the results of a study by Rozbicki *et al.* (2015). Moreover, *PC*, *SV*, *WG*, and *WA* were strongly positively correlated with *EE* while negatively correlated with the second parameter of uniaxial dough extensibility *R/EE*. Contrary to our findings, the study of Kun *et al.* (2020) did not show any strong correlation among these parameters in individual years. The alveograph trait *W* of Serbian cultivars strongly correlated with *PC*, *SV*, *WG*, *WA*, and *FR*. Also, the *W*

strongly correlated uniaxial extensibility dough traits; positive with *EE* and negative with *R/EE*.

Furthermore, bread volume (*BV*) was strongly positively correlated with the most important technological properties *PC*, *SV*, *WG*, *WA*, and *FR*, while negatively correlated with *R/EE*. These results are in accordance with Wrigley and Bekes (2002) previous findings, who showed a positive relationship between *PC* and *BV*, but not in accordance with Rozbicki *et al.* (2015), who showed a negative relationship between *WA* and *BV*. This correlation is used in the industry to help predict the suitability of flour for different end products.

Conclusions

Our study showed that recently developed Serbian wheat cultivars possess higher yield potential but lower end-use quality than most widespread local cultivars, especially cultivars Simonida, Zvezdana, and NS Futura. Also, the significant influence of cultivar, year, and their interaction on the variation of farinograph parameters *FR* and *SD* were found. Furthermore, weather conditions that prevailed in the growing seasons mostly influenced the extensograph parameters *E* and *R/E*. The year had a statistically significant impact on the most important parameter of end-use quality *BV* and almost half of the variation in the *BV* was explained by the influence of year. Moreover, strong positive correlations were found among *PC*, *SV*, *WG*, and *WA* with *EE* and a negative correlation with the second parameter of uniaxial dough extensibility *R/EE* was found. Therefore, further investigation of the studied technological traits variation is necessary to enable stable production of quality wheat cultivars.

References

- AACC, 2000. Approved methods of the American Association of Cereal Chemists (10th ed.). The American Association of Cereal Chemists, Inc., St. Paul, MN, USA.
- Akter N, Islam MR, 2017. Heat stress effects and management in wheat. A review. *Agron. Sustain. Dev.* 37:37.
- Asseng S, Ewert F, Martre P, et al., 2015. Rising temperatures reduce global wheat production. *Nat. Clim. Change* 5:143-7.
- Balla K, Rakszegi M, Li Z, Bekes F, Bencze S, Veisz O, 2011. Quality of winter wheat in relation to heat and drought shock after anthesis. *Czech J. Food Sci.* 29:117-28.
- Barbeau WE, Griffey CA, Yan ZH, 2006. Evidence that minor sprout damage can lead to significant reductions in gluten strength of winter wheats. *Cereal Chem.* 83:306-10.
- Bilgin O, Guzmán Ca, Başer İ, Crossa J, Korkut KZ, 2016. Evaluation of grain yield and quality traits of bread wheat genotypes cultivated in northwest Turkey. *Crop Sci.* 56:73-84.
- Blumenthal C, Gras PW, Bekes F, Barlow EWR, Wrigley CW, 1995. Possible role for the Glu-D1 locus with respect to tolerance to dough - quality change after stress. *Cereal Chem.* 72:135-6.
- Branlard G, Dardevet M, Saccomano R, Lagoutte F, Gourdon J, 2001. Genetic diversity of wheat storage proteins and bread wheat quality. *Euphytica* 119:59-67.
- Carson GR, Edwards NM, 2009. Criteria of wheat and flour quality. In: K. Khan, P.R. Shewry (Ed.), *Wheat: chemistry and technology*, 4th ed. AACC International, Inc., St. Paul, MN,

- USA, pp. 97-118.
- Constantinescu G, Dabija A, Buculei A, Rebenciuc I, 2011. Evaluation of cereal cultivar impact on bread quality. *J. Agroaliment. Proc. Technol.* 17:473-6.
- Da Silva C, Benin G, Bornhofen E, Todeschini M, Dallo S, Sassi L, 2014. Characterization of Brazilian wheat cultivars in terms of nitrogen use efficiency. *Bragantia* 73:87-96.
- Dencic S, DePauw R, Kobiljski B, Momcilovic V, 2013. Hagberg falling number and rheological properties of wheat cultivars in wet and dry preharvest periods. *Plant Prod. Sci.* 16:342-51.
- Dhaka V, Khatkar BS, 2015. Effects of gliadin/glutenin and HMW-GS/LMW-GS ratio on dough rheological properties and bread-making potential of wheat varieties. *J. Food Qual.* 38:71-82.
- Farrell AD, Kettwell PS, 2008. The effect of temperature shock and grain morphology on alpha-amylase in developing wheat grain. *Ann. Bot.-London* 102:287-93.
- Fleitas MC, Mondal S, Gerard GS, Hernández-Espinosa N, Singh RP, Crossa J, Guzmán C, 2020. Identification of CIMMYT spring bread wheat germplasm maintaining superior grain yield and quality under heat-stress. *J Cereal Sci.* 93:102981.
- Gras PW, Anderssen RS, Keentock M, Bekes F, Appels R, 2001. Gluten protein functionality in wheat flour processing: a review. *Aust J Agric Res.* 52:1311-23.
- Guo X, Sun X, Zhang Y, Wang R, Yan X, 2018. Interactions between soy protein hydrolyzates and wheat proteins in noodle making dough. *Food Chem.* 245:500-7.
- Hernández-Espinosa N, Mondal S, Autrique E, Gonzalez-Santoyo H, Crossa J, Huerta-Espino J, Singh RP, Guzmán C, 2018. Milling, processing and end-use quality traits of CIMMYT spring bread wheat germplasm under drought and heat stress. *Field Crops Res.* 215:104-12.
- Hlisnikovský L, Menšík L, Kunzová E, 2020. The development of winter wheat yield and quality under different fertilizer regimes and soil-climatic conditions in the Czech Republic. *Agronomy* 10:1160.
- Horvat D, Drezner G, Sudar R, Šimić G, Dvojčević K, Španić V, Magdić D, 2015. Distribution of wheat protein components under different genetic backgrounds and environments. *Turk. J. Field Crops* 20:150-4.
- Hristov N, Mladenov N, Djuric V, Kondic-Spika, A, Marjanovic-Jeromela A, Simic D, 2010. Genotype by environment interactions in wheat quality breeding programs in southeast Europe. *Euphytica* 174:315-24.
- Hrušková M, Švec I, Jirsa O, 2006. Correlation between milling and baking parameters of wheat varieties. *J. Food Eng.* 77:439-44.
- Hungarian Standard, 1988. Lisztvizsgálási módszerek. A vízfelvétel képesség és a sütőipari érték vizsgálata [Flour testing methods. Determination of water absorption capacity and baking quality]. Method MSZ 6369/6, MSZT, Budapest, Hungary.
- ICC, 2011. Standard Methods of the International Association for Cereal Science and Technology. Vienna, Austria.
- Johansson E, Branlard G, Cuniberti M, Flagella Z, Hüsken A, Nurit E, Peña RJ, Sissons M, Vazquez D 2020. Genotypic and environmental effects on wheat quality, pp 171-204 in G. Igrejas, T. M. Ikeda and C. Guzmán (Eds.), *Wheat quality for improving processing and human health*. Springer Nature Switzerland AG, Cham, Switzerland.
- Johansson E, Malik AH, Hussain A, Rasheed F, Newson WR, Plivelic T, Hedenqvist MS, Gällstedt M, Kuktaite R, 2013. Wheat gluten polymer structures: The impact of genotype, environment and processing on their functionality in various applications. *Cereal Chem.* 90:367-76.
- Kaluđerski G, Filipović N, 1998. Metoda kojom se u Jugoslaviji ocenjuje pecivost i mešavinska vrednost sote pšenice [Method for estimation baking properties of wheat cultivars and their mixtures], pp 255-6 in M. Žeželj, R Vukobratović (Eds.), *Metode ispitivanja kvaliteta žita, brašna i gotovih proizvoda* [Methods for the investigation of cereals, flour and final product quality]. Faculty of Technology, Novi Sad, Serbia.
- Kaya Y, Akcura M, 2014. Effects of genotype and environment on grain yield and quality traits in bread wheat (*T. aestivum* L.). *Food Sci. Technol. Campinas* 34:386-93.
- Kaya Y, Sahin M, 2015. Non-parametric stability analyses of dough properties in wheat. *Food Sci. Technol. Campinas* 35:509-15.
- Kondić D, Oručević Žuljević S, Hajder Đ, Selimbegović E, 2020. Evaluation of grain characteristics of domestic wheat (*Triticum aestivum* L.) obsolete cultivars and landraces. *Ital. J. Agron.* 15:3-9.
- Kondić-Špika A, Mladenov N, Grahovac N, Zorić M, Mikić S, Trkulja D, Marjanović-Jeromela A, Miladinović D, Hristov N, 2019. Biometric analyses of yield, oil and protein contents of wheat (*Triticum aestivum* L.) genotypes in different environments. *Agronomy* 9:270.
- Kong L, Si J, Zhang B, Feng B, Li S, Wang F, 2013. Environmental modification of wheat grain protein accumulation and associated processing quality: a case study of China. *Aust. J. Crop Sci.* 7:173-81.
- Konopka I, Fornal Ł, Abramczyk D, Rothkaehl J, Rotkiewicz D, 2004. Statistical evaluation of different technological and rheological tests of Polish wheat varieties for bread volume prediction. *Int. J. Food Sci. Technol.* 39:11-20.
- Kovačević M, 2011. Tehnološki postupak proizvodnje hleba [Bread Processing], pp. 79-148. In R. Perušković (Ed.), *Praktično pekarstvo* [Practical Breadmaking]. Progres, Novi Sad, Serbia.
- Kun S, Lijuan Y, Xiaohang L, Yang G, Zhikai J, Aiwang D, 2020. Grain quality variations from year to year among the Chinese genotypes. *Cereal Res. Commun.* 48:499-505.
- Laidig F, Piepho HP, Rentel D, Drobek T, Meyer U, Huesken A, 2017. Breeding progress, environmental variation and correlation of winter wheat yield and quality traits in German official variety trials and on-farm during 1983-2014. *Theor Appl Genet.* 130:223-45.
- Liana A, Alda S, For C, Diana M, Gogoasa I, Bordean D, Cărciu G, Cristea T, 2012. Climatic conditions influence on the variation of quality indicators of some Romanian and foreign winter wheat cultivars. *J. Hortic. For. Biotechnol.* 16:68-72.
- Liang D, Tang J, Peña RJ, Singh R, He X, Shen X, Yao D, Xia X, He Z, 2010. Characterization of CIMMYT bread wheat for high- and low-molecular weight glutenin subunits and other quality-related genes with SDS-PAGE, RP-HPLC and molecular markers. *Euphytica* 172:235-50.
- Linina A, Ruza A, Kunkulberga D, Rakcejeva T, 2014. The influence of environmental conditions on winter wheat wholemeal protein content and rheological properties. pp 66-70 in *Proceedings of 9th Baltic conference on Food Science and Technology: Food for Consumer Well-Being*, Foodbalt, Jelgava, Latvia,
- Liu L, Song H, Shi K, Liu B, Zhang Y, Tang L, Cao W, Zhu Y, 2019. Response of wheat grain quality to low temperature during jointing and booting stages - On the importance of considering canopy temperature. *Agr Forest Meteor.* 278:107658.
- Malik AH, 2012. Governing grain protein concentration and com-

- position in wheat and barley. *Acta Universitatis Agriculturae Sueciae (Swedish University of Agricultural Sciences)*, 2012:55.
- Malik AH, Kuktaite R, Johansson E, 2013. Combined effect of genetic and environmental factors on the accumulation of proteins in the wheat grain and their relationship to bread-making quality. *J Cereal Sci.* 57:170-4.
- Mares D, Mrva K, 2008. Late-maturity alpha-amylase: low falling number in wheat in the absence of preharvest sprouting. *J. Cereal Sci.* 47:6-17.
- Marinciu C, Săulescu NN, 2008. Cultivar effects on the relationship between grain protein concentration and yield in winter wheat. *Rom. Agric. Res.* 25:19-27.
- Mastilović J, Živančev D, Lončar E, Malbaša R, Hristov N, Kevrešan Ž, 2018. Effects of high temperatures and drought during anthesis and grain filling period on wheat processing quality and underlying gluten structural changes. *J. Sci. Food Agric.* 98:2898-907.
- Mirosavljević M, Momčilović V, Živančev D, Aćin V, Jocković B, Mikić S, Takac V, Denčić S, 2020. Genetic improvement of grain yield and bread-making quality of winter wheat over the past 90 years under the Pannonian Plain conditions. *Euphytica* 216:1-16.
- Olesen JE, Trnka M, Kersebaum KC, Skjelvag AO, Seguin B, Peltonen-Sainio P, Rossi F, Kozyra J, Micale F, 2011. Impacts and adaptation of European crop production systems to climate change. *Eur. J. Agron.* 34:96-112.
- Pan J, Zhu Y, Jiang D, Dai T, Li Y, Cao W, 2006. Modeling plant nitrogen uptake and grain nitrogen accumulation in wheat. *Field Crop. Res.* 97:322-36.
- Posner ES, Hibbs AN, 2005. *Wheat Flour Milling*. 1th ed. American Association of Cereal Chemists Inc, St Paul, MN, USA.
- Ral JP, Whan A, Larroque O, Leyne E, Pritchard J, Dielen AS, Howitt CA, Morell MK, Newberry M, 2016. Engineering high α -amylase levels in wheat grain lowers Falling Number but improves baking properties. *Plant Biotechnol. J.* 14:364-76.
- Rozbicki J, Ceglińska A, Gozdowski D, Jakubczak M, Cacak-Pietrzak G, Mądry W, Golba J, Piechociński M, Sobczyński G, Studnicki M, Drzazga T, 2015. Influence of the cultivar, environment and management on the grain yield and bread-making quality in winter wheat. *J. Cereal Sci.* 61:126-32.
- Sakr N, Rhazi L, Aussenac T, 2021. Bread wheat quality under limiting environmental conditions: II - Rheological properties of Lebanese wheat genotypes. *J. Saudi Soc. Agric. Sci.* 20:235-42.
- Sanchez-Garcia M, Álvaro F, Peremarti A, Martín-Sánchez JA, Royo C, 2015. Changes in bread-making quality attributes of bread wheat varieties cultivated in Spain during the 20th century. *Eur. J. Agron.* 63:79-88.
- Šarčević H, Jukić K, Ikić I, Lovrić A, 2014. Estimation of quantitative genetic parameters for grain yield and quality in winter wheat under high and low nitrogen fertilization. *Euphytica* 199:57-67.
- Schmid M, Wieser H, Koehler P, 2016. Isolation and characterization of high-molecular-weight (HMW) gliadins from wheat flour. *Cereal Chem.* 93:536-42.
- Shewry PR, Halford NG, Tatham AS, 1992. High molecular weight subunits of wheat glutenin. *J Cereal Sci.* 15:105-20.
- Tomás D, Rodrigues JC, Viegas W, Silva M, 2020. Assessment of high temperature effects on grain yield and composition in bread wheat commercial varieties. *Agronomy* 10:499.
- Trnka M, Rötter RP, Ruiz-Ramos M, Kersebaum KC, Olesen JE, Žalud Z, Semenov MA, 2014. Adverse weather conditions for European wheat production will become more frequent with climate change. *Nat. Clim. Change* 4:637-43.
- Wehrmann J, Scharpf HC 1979. Mineral nitrogen concentration of soil as a gauge of need for nitrogen-fertilizer (N-Min method). *Plant Soil* 52:109-26.
- Wrigley CW, Bekes F, 2002. Wheat quality elucidation the Bushuk Legacy. pp 65-86 in P.K.W Ng, C.W. Wrigley (Eds.), *Grain-protein composition as a document of wheat-quality type: new approaches to varietal identification*. American Association of Cereal Chemists Inc, St Paul, MN, USA.
- Yang X, Wu L, Zhu Z, Ren G, Liu S, 2014. Variation and trends in dough rheological properties and flour quality in 330 Chinese wheat varieties. *Crop J.* 2:195-200.
- Zečević V, Knežević D, Bošković J, Madić M, 2009. Effect of genotype and environment on wheat quality. *Genetika* 41:247-53.
- Živančev DR, Torbica AM, Tomić JM, Janić Hajnal EP, Belović MM, Mastilović JS, Kevrešan ŽS, 2016. Effect of climate change on wheat quality and HMW glutenin subunit composition in the Pannonian plain. *Cereal Chem.* 93:90-9.