

**XXVI INTERNATIONAL
ECO-CONFERENCE® 2022
21–23th SEPTEMBER**

XII SAFE FOOD



PROCEEDINGS

NOVI SAD, SERBIA

XXVI INTERNATIONAL ECO-CONFERENCE® 2022

XII SAFE FOOD

21nd – 23th SEPTEMBER 2022.

NOVI SAD, SERBIA

**XXVI INTERNATIONAL ECO-CONFERENCE
XII SAFE FOOD**
21nd – 23th SEPTEMBER 2022. NOVI SAD, SERBIA

Publisher

ECOLOGICAL MOVEMENT OF NOVI SAD
21102 Novi Sad, str. Cara Lazara 83/1
Phone: (+381 21) 6372 940
Mob: (+381 69) 304 73 38
E-mail: ekopokretns@gmail.com
www.ekopokret.org.rs

Editorial Board

Prof. Dr. Miroslav Malesevic, President
Nikola Aleksic
Prof. Dr. Desanka Bozidarevic
Prof. Dr. Vladan Joldzic
Prof. Dr. Velibor Spalevic
Prof. Dr. Viktor Zakrevski
Prof. Dr. Vera Popović

Project Editor

Nikola Aleksic

Copy Editor

Vesna Karajovic

For the Publisher

Nikola Aleksic

Print

Red Copy, Novi Sad

Cirkulation

100 copies

Publication year: 2022-09-18
THE AUTORS ARE RESPONSIBLE FOR THE QUALITY
OF ENGLISH TRANSLATION

XXVI INTERNATIONAL ECO-CONFERENCE® 2022

XII SAFE FOOD

21nd – 23th SEPTEMBER 2022

NOVI SAD, SERBIA

SAFE FOOD

PROCEEDINGS

2022

Organizer:



Ecological Movement of Novi Sad

Co-organizers:



University
Novi Sad



Russian State Agrarian University
– MTAA, Moskow, Russian Federation



International Independent
Ecological-Politicology
University in Moscow, RF



Institute for Field
and Vegetable Crops
Novi Sad, Serbia



Pasteur Institute
of Novi Sad, Serbia



Scientific Veterinary
Institute "Novi Sad"
Serbia



Legambiente, d' Itali
(National environmental
Organisation)

Patronage:



Matica srpska,
Novi Sad

Host:



INSTITUTE FOR NATURE CONSERVATION
OF VOJVODINA PROVINCE

Institute for Nature Conservation
of Vojvodina Province, Novi Sad, Serbia

Honorary Committee:

President:

– **Dragan Stanic, Ph.D**, President, Matica srpska, Serbia

Vice-Presidents:

- **Dejan Madic, Ph.D**, Rector, University of Novi Sad, Serbia
- **V. I. Trukhachev, Ph.D**, Rector, Russian University of Agriculture, Moscow, Russian Federation
- **Stanislav A. Stepanov, Ph.D**, Rector, International Independent Ecological-Politicology University, Moscow, Russian Federation
- **Jegor Miladinovic, Ph.D**, Head, Institute of Field and Vegetable Crops, Novi Sad, Serbia
- **Nenad Vranjes, Ph.D**, Head, Pasteur Institute, Novi Sad, Serbia
- **Sava Lazic, Ph.D**, Head, Scientific Veterinary Institute Novi Sad, Novi Sad, Serbia
- **Vittorio Cogliati Dezza, Ph.D**, President, Legambiente, Rome, Italy
- **Natasa Saric**, Acting Head, Institute for Nature Conservation of Serbia, Department of Geodiversity, Novi Sad, Serbia

Scientific Committee

President:

– **Miroslav Malesevic, Ph.D**, Faculty of Agriculture, University of Novi Sad, Serbia

Vice Presidents:

- **I. I. Vasenev, Ph.D**, Vice Rector, Russian State Agrarian University, MTAA, Moscow, Russian Federation
- **Marfenin N. N., Ph.D**, Vice Chancellor, International Independent Ecological-Politicology, University in Moscow, Russian Federation
- **Moreno Baccichet, Ph.D**, Faculty of Philology and Philosophy, University of Ferrara, Italy
- **Nikola Jorgovanovic, Ph.D**, Vice Rector of Science and Art, University of Novi Sad, Serbia
- **Ana Jeromela Marjanovic, Ph.D**, Assistant Head of the Science and Research, Institute of Field and Vegetable Corps, Novi Sad, Serbia
- **Dusan Lalosevic, Ph.D**, Head, Pasteur Institute of Novi Sad, Serbia
- **Tamas Petrovic, Ph.D**, Assistant Head of Science, Scientific Veterinary Institute, Novi Sad, Serbia

Secretary:

- **Ljubica Trbojevic**, Assistant Director, Environmental Movement of Novi Sad, Serbia

Members

- **Academician Serbislav Dencic**, Institute of Field and Vegetable Crops, Novi Sad, Serbia
- **Academician Branka Lazic**, The Faculty of Agriculture, University of Novi Sad, Serbia
- **Academician Vukadin Leovac**, Faculty of Science, Department of Chemistry, University of Novi Sad, Serbia
- **Radmila Bojovic, Ph.D**, Faculty of Agriculture, University of Belgrade, Zemun, Serbia
- **Desanka Bozidarevic, Ph.D**, Resident Professor, Faculty of Agriculture, University of Novi Sad, Serbia
- **Vesna Gantner, Ph.D**, Faculty of Agrobiotechnical Sciences Osijek, J. J. Strossmayer University of Osijek, Croatia
- **Ales Golja, Ph.D**, Faculty of Civil and Geodetic Engineering, University of Ljubljana, Slovenia
- **Jasna Grabic, Ph.D**, Faculty of Agriculture, University of Novi Sad, Serbia
- **Ivana Djujic, Ph.D**, Resident Professor, Faculty of Biology University of Belgrade, Serbia
- **Jela Ikanovic, Ph.D**, Faculty of Agriculture, University of Belgrade, Serbia
- **Gabrijela Jovcev, Ph. D**, Institute of Biodiversity and Ecosystem Research, Sofia, Bulgaria
- **Vladan Joldzic, Ph.D**, Resident Professor, Faculty of Biology and Chemistry, University of Belgrade, Serbia, Institute for Criminological and Sociological Research, Belgrade, Serbia
- **Ana Jovicic Vukovic, Ph.D**, High Business School, Novi Sad, Serbia
- **Ksenija Mackic, Ph.D**, Faculty of Agriculture, University of Novi Sad, Serbia
- **Ivana Maksimovic, Ph. D**, Resident Professor, Faculty of Agriculture, University of Novi Sad, Serbia
- **Andreja Mihailović**, Faculty of Law, University of Montenegro, Podgorica, Montenegro
- **Zeljko Mihaljev, Ph.D**, Higher Research Associate, Science Veterinary Institute „Novi Sad”, Novi Sad, Serbia
- **Aleksandar Milovanovic, Ph.D**, Higher Research Associate, Science Veterinary Institute, Novi Sad, Serbia
- **Rodoljub Oljaca, Ph.D**, Resident Professor, Faculty of Forestry, University of Banja Luka, Bosnia and Herzegovina
- **Vera Popovic, Ph.D**, Principal Research Fellow, Institute of Field and Vegetable Corps, Novi Sad, Serbia

- **Aleh Rodzkin, Ph.D**, International Sakharov Environmental University, Minsk, Belarus
- **Atila Salvai, Ph.D**, Faculty of Agriculture, University of Novi Sad, Serbia
- **Ivan Simunic, Ph.D**, Resident Professor, Faculty of Agriculture, University of Zagreb, Croatia
- **Velibor Spalevic, Ph.D**, Biotechnical Faculty, University of Montenegro, Montenegro
- **Mohammad Taghi Razavian, Ph.D**, Associate Professor, Department of Geography, Sahid Beheshti University, Evin, Tehran
- **Ion C. Ungureanu, Ph.D**, Professor, Faculty of Agricultural Sciences and Veterinary Medicine, University of Bucharest, Romania
- **Lu Zhongmei, Ph.D**, University of Beijing, University of Wuhan, Vice President of the High People’s Court, Hubei, Wuhan, China, Member of the National People’s Congress of the Republic of China

Organizational Committee

President:

- **Nikola Aleksic**, President, Environmental Movement of Novi Sad, Serbia

Vice President:

- **Angelo Mancone**, Legambiente, Verona, Veneto, Italy

Secretary:

- **Zoran Kovacevic**, Organizer, Environmental Movement of Novi Sad, Serbia

Members:

- **Vera Rajičić, Ph.D**, Faculty of Agriculture, University of Niš, Serbia
- **Djordje Okanovic, Ph.D**, Research Associate, Institute of Field and Vegetable Corps, Novi Sad, Serbia
- **Vera Popovic, Ph.D**, Principal Research Fellow, Institute of Field and Vegetable Corps, Novi Sad, Serbia
- **Dragoslav Stoilkovic, Ph.D**, Faculty of Technology, University of Novi Sad, Serbia
- **Milan Vurdelja, Mr.Sc**, Rectorate, University of Novi Sad, Serbia
- **Bratimir Nesic**, Graduate Engineer, Assistant President, Environmental Movement of Novi Sad, Serbia
- **Luka Vujasinovic**, Web Administrator, Environmental Movement of Novi Sad, Serbia

Official host of the XXVI International Eco-conference 2022 – Institute for Nature Conservation of Vojvodina Province in Novi Sad

CONTENT

THE ECOLOGICAL MOVEMENT OF NOVI SAD: AN IMPORTANT DECISION OF ITS PROGRAMME COUNCIL	9
FOREWORD	21

INTRODUCTORY PRESENTATION

<i>Vesna Gantner, Vera Popović, Djoko Bunevski, Muhamed Brka, Denis Kučević, Ranko Gantner</i> CHALLENGES AND SOLUTIONS FOR AGRICULTURAL PRODUCTION IN THE UPCOMING ERA	25
<i>Slavica Vesković Moračanin, Dragutin Đukić</i> FOOD CONTROL SYSTEMS AND PROTECTION OF HUMAN HEALTH	37

SOIL AND WATER AS THE BASIS FOR AGRICULTURAL PRODUCTION OF HEALTHY SAFE FOOD

<i>Dejan Zejak, Velibor Spalevic, Vera Popovic, Mile Markoski, Branislav Dudic, Abdessalam Ouallali, Artan Hysa, Ronaldo Luiz Mincato, Shuraik Kader</i> ANALYSIS OF THE PRESENCE OF HEAVY METALS IN THE SOILS OF THE HILLY-MOUNTAINOUS AREAS OF BALKAN PENINSULA WITH THE ASSESSMENT OF ITS POTENTIAL FOR THE FRUIT GROWING: CASE STUDY OF THE LJUBOVIDJA RIVER BASIN, POLIMLJE, MONTENEGRO	51
<i>Gordan Vrbanec, B.Sc.</i> AUTOMATED ECOMONITORING OF WATER IN THE FUNCTION OF SOIL AND WATER PROTECTION.....	63

<i>Ksenija Mačkić, Borivoj Pejić, Ljiljana Nešić, Milivoj Belić, Vladimir Ćirić, Jovica Vasin, Vera Popović, Dragan Radovanović</i> THE STRUCTURE OF VERTISOL SOIL IN THE AREA OF THE MUNICIPALITY OF SMEDEREVO	69
---	----

GENETICS, GENETIC RESOURCES, BREEDING AND GENETIC ENGINEERING IN THE FUNCTION OF HEALTHY SAFE FOOD PRODUCTION

<i>Prof. Dr. Sc. Aleksandar Markovski, Assoc. Prof. Dr. Sc. Lenche Velkoska-Markovska</i> GREEK STRAWBERRY TREE (<i>Arbutus andrachne L.</i>) GENOTYPES PROPAGATION	79
---	----

<i>Vera Popović, Vesna Gantner, Ljubica Šarčević Todosijević, Elizabetha Miskoska Milevska, Marija Mujović, Milica Vujović, Radmila Bojović</i> PRODUCTIVITY OF GLUTEN-FREE PSEUDOCEREAL <i>Fagopyrum esculentum</i> Moench – NOVOSADSKA BUCKWHEAT ON THE CHERNOZEM	89
--	----

<i>PhD student, Bojan Laloš, PhD student, Stanka Pešić, Doctor of Science, Violeta Mickovski Stefanović, Mr Predrag Brković, PhD student, Miloš Pavlović, Doctor of Science, Jasmina Bačić, Doctor of Science, Dragana Stanisavljević, Master of Arts, Predrag Ilić</i> BANKUT WHEAT	97
---	----

<i>Vladimir M. Filipović, Ljubica Šarčević-Todosijević, Vladan Ugrenović, Slobodan Popović, Milan Ugrinović, Snežana Dimitrijević, Dragana Popović</i> THE INFLUENCE OF DIFFERENT SUBSTRATE MIXTURES ON SEEDLINGS QUALITY AND ECONOMICALLY HIGH YIELD OF TOMATO, PEPPER AND BASIL	103
--	-----

INTEGRATED PLANTS PROTECTION, PESTICIDES APPLICATION AND EFFECTS OF RESIDUES

<i>Assoc. Prof. Dr. sc. Lenche Velkoska-Markovska, Prof. Dr. sc. Biljana Petanovska-Ilievska, Prof. Dr. sc. Aleksandar Markovski</i> APPLICATION OF NORMAL-PHASE HIGH PERFORMANCE LIQUID CHROMATOGRAPHY TO THE ANALYSIS OF SOME PESTICIDE RESIDUES IN APPLE JUICE	115
--	-----

<i>Vesna Perišić, Vladimir Perišić, Vera Rajičić, Kristina Luković, Filip Vukajlović</i> SPINOSAD APPLICATION IN PROCESS OF INTEGRATED PEST MANAGEMENT AGAINST <i>RHYZOPERTHA DOMINICA</i> F. IN STORED SMALL GRAINS	125
---	-----

*Ljubica Šarčević-Todosijević, Snežana Dorđević, Vera Popović,
Dragutin Đukić, Marija Perić, Nikola Dorđević, Ljubiša Živanović,
Ksenija Mačkić, Jelena Bošković, Aleksandar Stevanović*
THE INFLUENCE OF PESTICIDES ON PLANTS,
SOIL MICROORGANISMS AND FOOD SAFETY
IN PLANT PRODUCTION..... 133

*A. Georgieva, J. Miteva-Staleva, E. Krumova, A. Dobрева, G. Jovtchev,
S. Gateva, Ts. Gerasimova, and Milka Mileva*
ESSENTIAL OILS FROM BULGARIAN *ROSA ALBA* L.
AND *ROSA DAMASCENA* MILL. AMPLIFY THE ANTIFUNGAL
EFFECT OF NYSTATIN..... 141

AGRICULTURAL PRODUCTION IN VIEW OF SUSTAINABLE DEVELOPMENT ASPECT

Éva Erdélyi, Judit Lovasné Avató
IMPACT OF FOOD PRODUCTION AND CONSUMPTION
ON THE ENVIRONMENT, GEOGRAPHICAL APPROACH 151

Zsolt Törösvári, Éva Erdélyi
THE ROLE OF FOOD AND DRINK HUNGARIKUMS
IN TOURISM AND HOSPITALITY..... 161

PRODUCTION OF FIELD AND VEGETABLE CROPS FROM THE POINT OF HEALTHY SAFE FOOD

Bojana Kovačević Berleković, Ana Jovičić Vuković, Tatjana Bošković
THE SIGNIFICANCE OF SAFE FOOD IN TRAVEL AND TOURISM
DEVELOPMENT IN THE POST-COVID-19 ERA..... 173

Jovana Bajkanović, Vesna Vujasinović, Nemanja Lakić
STUDENTS 'ATTITUDES TOWARD THE USE
OF FOOD ADDITIVES 181

FRUIT AND GRAPES PRODUCTION FROM THE STANDPOINT OF HEALTHY SAFE FOOD

Elizabeta Miskoska-Milevska, Ana Selamovska, Katerina Nikolic
GENOTYPE EXPRESSION OF TRADITIONAL PEAR VARIETY SINEC
DEPENDING ON ECOLOGICAL FACTORS 191

<i>Prof. Ana Selamovska, Prof. Viktor Gjamovski, Prof. Milena Taseska-Gjorgjijevski, Prof. Dusko Nedelkovski, Prof. Katerina Bandjo Oreshkovikj</i>	
CONTENT OF SOME ANTIOXIDANTS IN THE FRUITS OF AUTOCHTHONOUS APPLE VARIETIES.....	199

<i>Prof. Dr. sc. Ljubica Karakashova, Assist. Prof. Dr. sc. Frosina Babanovska-Milenkovska, MSc. Jelena Mladenova, Assoc. Prof. Dr. sc. Lence Velkoska-Markovska, Assoc. Prof. Dr. sc. Viktorija Stamatovska, Assoc. Prof. Dr. sc. Namik Durmishi</i>	
COMPARISON OF QUALITY PROPERTIES OF FRESH AND SOLAR DRIED <i>GOJI BERRIES (LYCIUM BARBARUM)</i>	207

LIVESTOCK PRODUCTION FROM THE STAND POINT OF HEALTHY SAFE FOOD

<i>Franjo Poljak, Marcela Šperanda, Mirna Gavran, Mislav Đidara, Vera Popović, Vesna Gantner</i>	
THE VARIABILITY OF BIOCHEMICAL PARAMETERS IN BLOOD SERUM AND MILK OF HOLSTEIN COWS REGARDING THE AGE AND HEALTH STATUS.....	219

<i>Željko Mihaljev, Milica Živkov Baloš, Sandra Jakšić, Nenad Popov</i>	
LOAD OF SOIL, FOOD, ANIMAL FEED AND BIOLOGICAL MATERIAL WITH RADIOACTIVE RESIDUES	227

<i>Ivana Jožef, Dragan Solić, Marcela Šperanda, Muhamed Brka, Mislav Đidara, Vesna Gantner</i>	
THE CORRELATION BETWEEN SOME DAILY MILK TRAITS AND BIOCHEMICAL PARAMETERS IN PLASMA AND MILK OF DAIRY COWS	235

<i>Aleksandra Ivetić, Bojan Stojanović, Vesna Davidović, Milivoje Čosić</i>	
THE USE OF BIOWASTE AS A SILAGE BY-PRODUCT FOR SUSTAINABLE FOOD AND FEED PRODUCTION	245

APPLICATION OF FOOD TECHNOLOGY IN THE PRODUCTION OF HEALTHY SAFE FOOD

<i>Vladimir Filipović Ph.D., Jelena Filipović Ph.D., Milica Nićetin Ph.D., Biljana Lončar Ph.D., Violeta Knežević Ph.D.</i>	
NEW TYPE OF COOKIE PRODUCT WITH DEHYDRATED PEACH.....	255

<i>Doctor of Science, Dragana Stanisavljević, PhD student, Stanka Pešić, Master filolog Predrag Ilić, Doctor of Science, Violeta Mickovski Stefanović</i>	HOME BREWING OF CRAFT BEER	263
---	----------------------------------	-----

ECONOMIC STANDPOINT OF HEALTHY SAFE FOOD PRODUCTION AND MARKETING

<i>Goran Vitomir, Slobodan Popović, Ivanka Vranić, Vinka Vukeljić, Dragan Božović</i>	COOPERATION OF FORENSIC ACCOUNTING AND FORENSIC AUDIT IN FOOD PRODUCTION COMPANIES	275
---	---	-----

<i>Jelena Vitomir, Slobodan Popović, Sonja Tomas-Miskin</i>	MACRO-ECONOMIC RESPECT FOR HETEROGENEOUS ENVIRONMENTAL FACTORS IN COMPANIES ENGAGED IN THE PRODUCTION OF HEALTHY SAFE FOOD	281
---	--	-----

<i>Ivanka Vranić, Slobodan Popović, Goran Vitomir</i>	INDEPENDENT FORENSIC ACCOUNTING IN FOOD COMPANIES	287
---	---	-----

<i>Dragan Dokić, Mirna Gavran, Danko Šinka, Vesna Gantner</i>	THE IMPORTANCE OF OPTIMIZING THE VALUE-ADDED TAX RATE IN PIGLET PRODUCTION	293
---	---	-----

<i>Zsuzsanna Tóth, Csaba Limbek</i>	COMPARISON OF THE SYSTEM OF EU COMMUNITY TRADEMARKS AND HUNGARIKUMS	301
-------------------------------------	--	-----

<i>Sonja Tomas-Miskin, Jelena Vitomir, Slobodan Popović</i>	ECONOMIC RESPECT FOR HETEROGENEOUS FACTORS IN COMPANIES ENGAGED IN THE PRODUCTION OF HEALTHY SAFE FOOD	309
---	--	-----

<i>Slobodan Popović, Sonja Tomas-Miskin, Jelena Vitomir</i>	IMPLEMENTATION OF INTERNAL AUDIT IN COMPANIES MAINLY OPERATING IN THE FIELD OF HEALTHY SAFETY PRODUCTION.....	317
---	--	-----

<i>V.V. Zakrevskii</i>	THE ROLE OF THE BIOLOGICAL AND NUTRITIONAL VALUE OF THE DIET IN THE PREVENTION OF CORONAVIRUS INFECTION	325
------------------------	--	-----

<i>Ana Janžeković, Lect. Dr. Aleš Golja, Prof. Dr. Andrej Udovč</i>	CURRENT ISSUES ON LESSER-KNOWN FIELD CROPS FOR FOOD SAFETY PURPOSES	333
---	--	-----

NUTRITIONAL FOOD VALUE AND QUALITY NUTRITION

*Jela Ikanović, Vera Popović, Ljubiša Živanović, Ljubiša Kolarić,
Vera Rajičić, Vladimir Filipović, Ljubičić Nataša, Kristina Kajiš*
IMPORTANCE OF BROAD BEAN IN THE DIET – POSSIBILITY
OF MORE RATIONAL USE OF MAIN AND BY-PRODUCTS..... 343

*Vojislav Trkulja, Nikola Ilić, Vera Popović, Vladan Pešić, Ljubiša Kolarić,
Gordana Dražić, Nikola Rakašćan*
INFLUENCE OF GENOTYPES AND DIGESTATE
ON THE PRODUCTIVITY OF THE MORPHOLOGICAL INDICATOR
OF SILAGE-SORGHUM AS HIGH QUALITY FEED 353

Maja Milošević, Milica Perović, Zorica Knežević Jugović, Mirjana Antov
SOY HULL PECTIC FIBER – HIGHLY VALUABLE FIBER
EXTRACTED FROM BY-PRODUCT OF SOYBEAN INDUSTRY 363

Milica Perović, Maja Milošević, Zorica Knežević Jugović, Mirjana Antov
GREEN TECHNOLOGY FOR ENHANCED RECOVERY AND
PROPERTIES OF RuBisCO PROTEIN FROM PUMPKIN LEAVES 367

*Svetlana Tepavac, Vesna Vujasinović, Marija Vukadin, Jovana Bajkanović,
Đurkinjak Mirjana, Račićević Nikoleta*
STUDENT’S PERCEPTION OF SOURCES AND SIGNIFICANCE
VITAMINS A AND D IN THE DIET 371

*PhD student Stanka Pešić, PhD student, Bojan Laloš,
Doctor of Science, Violeta Mickovski Stefanović, Doctor of Science,
Jasmina Bačić, Doctor of Science, Msc Predrag Brković,
Dragana Stanisavljević, Master of Arts, Predrag Ilić*
NUTRITIONAL VALUE OF REFRESHING
NON-ALCOHOLIC BEVERAGES 379

PhD and DA Agota Vitkay-Kucsera
THE IMPACT OF DIET ON ELITE VOCAL PROFESSIONALS
WITH SPECIAL REFERENCE TO STOMACH ACID REFLUX
(GERD – GASTROESOPHAGEAL REFLUX DISEASE
AND LPR – LARYNGOPHAGEAL REFLUX) 385

LEGAL ASPECTS OF HEALTHY SAFE FOOD PRODUCTS PROTECTION

Prof. Dr. Joldžić Vladan
PRODUCTION OF HEALTHY SAFE FOOD
IN THE LIGHT OF LAND AND WATER PROTECTION..... 395

<i>Ana Jovičić Vuković, Bojana Kovačević Berleković, Snježana Gagić Jaraković</i>	
THE IMPORTANCE OF THE APPLICATION OF FOOD STANDARDS IN THE HOSPITALITY INDUSTRY	405

ECOLOGICAL MODELS AND SOFTWARE
IN THE PRODUCTION OF HEALTHY
SAFE FOOD

<i>Ranko Gantner, Gordana Bukvić, Goran Herman, Vesna Gantner</i>	
DESIGNING OF FORAGE SYSTEMS FOR IMPROVED SUSTAINABILITY OF CATTLE FARMING AND AGRICULTURE	415

<i>Ts. Gerasimova, G. Jovtchev, S. Gateva, M. Topashka-Ancheva, Ts. Angelova, A. Dobрева, M. Mileva</i>	
STUDY ON POTENTIAL CYTOTOXIC AND CLASTOGENIC EFFECTS OF <i>ROSA CENTIFOLIA</i> L. WASTEWATER	423

NAME REGISTRY	433
---------------------	-----



**Dejan Zejak^{1*}, Velibor Spalevic^{2,3}, Vera Popovic⁴, Mile Markoski⁵,
Branislav Dudic^{6,7}, Abdessalam Ouallali⁸, Artan Hysa⁹,
Ronaldo Luiz Mincato¹⁰, Shuraik Kader¹¹**

¹ Faculty of Agriculture, University of Belgrade, Nemanjina 6,
11000 Belgrade, Serbia;

² Biotechnical Faculty, University of Montenegro, 81000 Podgorica, Montenegro;

³ Geography, Philosophy, University of Montenegro, D. Bojovica bb,
81400 Niksic, Montenegro;

⁴ Institute of Field and Vegetable Crops, M. Gorkog 30, 21101 Novi Sad, Serbia;

⁵ Faculty of agricultural sciences and food Skopje, Skopje, North Macedonia

⁶ Faculty of Management, Comenius University in Bratislava,
81102 Bratislava, Slovakia;

⁷ Faculty of Economics and Engineering Management, University Business
Academy, Novi Sad, RS;

⁸ Faculty of Sciences and Techniques of Mohammedia, Hassan II University
of Casablanca, Morocco;

⁹ Faculty of Engineering and Architecture, Epoka University, 1032 Tirana, Albania;

¹⁰ Institute of Natural Sciences, Federal University of Alfenas, UNIFAL-MG,
Minas Gerais, Brazil;

¹¹ Department of Civil Engineering, Sri Lanka Institute of Information Technology,
Malabe, Sri Lanka.

* Corresponding author: zejakd@gmail.com

Original Scientific paper

**ANALYSIS OF THE PRESENCE OF HEAVY METALS
IN THE SOILS OF THE HILLY-MOUNTAINOUS AREAS
OF BALKAN PENINSULA WITH THE ASSESSMENT
OF ITS POTENTIAL FOR THE FRUIT GROWING:
CASE STUDY OF THE LJUBOVIDJA RIVER BASIN,
POLIMLJE, MONTENEGRO**

Abstract

The paper presents results of the analysis of the presence of heavy metals with an assessment of its potential for the further development of fruit growing in North Montenegro, and in this specific case at the territory of the River Basin Ljubovidja, which is the hilly-mountainous area

of the Balkan Peninsula. Sampling and further analysis showed that the dominant type of soil in this area is District Cambisol. In the second phase, we conducted further research on the presence of heavy metals in the area of Pavino Polje, in the municipality of Bijelo Polje, Montenegro. The results of the analysis finally showed that there is no presence of heavy metals in the samples we collected during the field visit. The studied area is recommended for the establishment of organic production. It is particularly suitable for the production of fruit growing, vegetables and fodder is also recommended.

Keywords: *Heavy metals, mountainous areas, fruit growing, organic production, Polimlje, Montenegro, Balkan Peninsula*

INTRODUCTION

At the end of the 20th century, significant changes occurred in the world: social, economic, and political (Zejak et al., 2022). Population explosion, industrialization, urbanization and exploitation of natural resources are creating problems globally (Nagpal et al., 2020). The problem of land degradation is more and more one of the most important issues that should be treated very seriously. Large quantities of different kinds of elements are discharged into the environment as contaminants each year by human activities. Heavy metal toxicity and its lack of biodegradability and long-lasting duration in the environment while entering the food chain are considered to be one the most serious potential pollutants of the ecosystem (Popović, 2002; Pakzad et al., 2016; Kastratovic et al., 2016). No or low degraded soil and well-protected lands are today a special asset of every society. Proper management of this resource is an obligation and is part of all wise decisions of those who deal with agrarian policies and sustainable management.

According to the official statistic (MONSTAT), Montenegro is with 0.84 ha per inhabitant of the total agricultural land. In relation to this criterion, the country is in the second position in the European continent after Ireland. On the other hand, this country is poor in fertile land, because many meadows have been turned into pastures, and pastures into forest land; and finally barren land counts for 23% of the territory. That is why Montenegrin agriculture is characterizing the so-called rural agriculture, i.e. production took place just for satisfying farmers' own needs (Fustic and Spalević, 2003; Fustic et al., 2005). Land degradation in Montenegro is caused by numerous natural and anthropogenic factors. The distinctly hilly and mountainous area of Montenegro (94% with a slope of over 5%), a very dynamic and scattered relief, a developed and dense network of watercourses, an abundance of precipitation (1000-5000 mm per year) and other climatic characteristics are factors of crucial importance for provoking of water and wind erosion. This is contributed by the high share of erodible and non-resistant soils substrates, the devastation of plant cover and bare land, which make up 23% of the total territory, the damages are enormous, multiple and incalculable. Another type of land degradation is conditioned by excess floods that threaten 26,000 ha in flat areas with varying intensity covering only 5% of Montenegro. Applying soil conservation measures, these potentially fertile lands should be brought to intensive agriculture production – culture for more intensive use in agriculture. Technological development

caused an increase in the use of natural resources by exploitation of mineral raw materials, industrialization and urbanisation – construction of infrastructure facilities and settlements, which "consumed" a considerable area, mostly the most fertile land. With this particular form of degradation through conversion – land use change, Montenegro has lost 800-1000 ha per year in the last 40-50 years (Fustic, 2005).

In Montenegro, the transition from conventional agriculture to organic production is increasing. This method of production is well accepted by those who are engaged in soil conservation. Unlike conventional agriculture, which, due to the application of pesticides and fertilizers, led to the loss of individual plant and animal species, organic farming is based on the principles of protection and preservation of plant and animal species and the environment. Organic production is based on the biological balance of the system land – plant – animal – human. According to the Codex Alimentarius definition, it is a "holistic" production system that promotes and strengthens the agro-ecosystem and health, including biodiversity, biological cycle, and soil. Organic production is one of the world's leading trends in agriculture that is continuing to grow. Organic production of agricultural and other products is based on the application of organic production methods at all stages of production, aims to reduce the use of chemicals, and excludes the use of GMOs and products consisting of or derived from GMOs, as well as the use of ionizing radiation. Organic farming aims to produce safe food, of high quality, in an environmentally sustainable way, maintaining the genetic diversity of agro and ecosystem, preserving the environment, maintaining and improving soil fertility, reducing all forms of pollution, and producing food of high nutritional value, improving health and making a profit (Popović et al., 2014; 2019; Zejak et al., 2022; Ikanović and Popović, 2020). The land we want for the future must be carefully treated and well protected today.

MATERIALS AND METHODS

Study area. Montenegro is a country with great variability, ranging from sand and rock coasts (some corresponding to a ria coast), karst plateaus, large (intra-montane) fields, high mountains holding a (peri) glacial imprint, canyons, and more; all of it comprised in only 13 812 km² and within an elevation range of 2535 m. It is characterized by a Mediterranean climate, with warm and dry summers and autumns, and relatively cold winters with heavy snowfall on the continent (Zejak et al., 2022). The average altitude is 1,086 m; the lowest point: is the Adriatic Sea 0 m; the highest point: is Prokletije (Zla Kolata, 2534 m a.s.l., Dobra Kolata, 2528 m); Durmitor Kuk (Bobotov Kuk, 2522 m). According to data provided in the papers of various authors (Zejak et al, 2022; Spalevic et al., 2019; 2020), the structure of land use in Montenegro is as follows: agricultural land: 38.2% (estimated in 2018); arable land: 12.9% (estimated in 2018); crops: 1.2% (estimated in 2018); pasture: 24.1% (estimated in 2018); forest: 40.4% (estimated in 2018); other: 21.4% (estimated 2018). Irrigated land 24 km² (2012). The Map of Montenegro with the studied region of River Basin Ljubovidja is presented in Figure 1. North Montenegro, where the studied region is located is mainly mountainous. In this neighbouring region, the highest peaks of Montenegro

are found, including Komovi (2487 m a.s.l.) and Zla Kolata (2535 m a.s.l.) in the Prokletije Mountains of the same Polimlje Region. Nevertheless, this area of Polimlje is quite densely populated and includes the towns of Plav, Andrijevica, Berane, and Bijelo Polje. The rivers in this region drain to the Black Sea, and some of them form deep canyons crossing limestone formations. Further downstream, they form broader valleys flowing through softer Palaeozoic material (Spalevic, 2011).




Figure 1. Map of MNE, Ljubovidja basin and Pavino Polje, (Zejak et al., 2022)

Table 1. Maximum allowed concentrations of dangerous and harmful substances in soils based on standards of Ex-Yugoslavia (mg/kg of air-dry soil)

Element	Official Gazette of the Republic of Serbia 23/1944	Rulebook Of Council Of Europe 2092/91	Rulebook on methods organic herbal of production, Official Gazette of FRY 51/2002
Cadmium	3	2.0	0.8
Lead	100	100.0	50.0
Mercury	2	1.0	0.8
Arsenic	25	-	10.0
Chrom	50	50.0	30.0
Nickel	300	-	-
Copper	100	50.0	50.0
Zinc	300	150.0	150.0
Boron	50	-	-
Molybdenum	-	-	10.0
Cobalt	-	-	30.0

Source: Kastori et.al., 2003.

Climate. Based data of more than 70 years (1948–2021) for temperature and precipitation, the study area is characterized by a mountain Mediterranean climate (Rajović, 2009) with rainy autumns and springs, cold winters, and a deficit of precipitation in summer months (Table 2.)

Table 2. Precipitation and temperature for the period 1948–2021 recorded at Bijelo Polje

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
<i>Max daily precipitation in mm</i>												
Max	68.6	92.8	73	93.3	42.6	58.5	97.8	55.8	95.6	157.6	101.6	79.4
Aver.	23.4	22.8	21.5	24	21.9	20.6	21.7	21.1	25.3	29	29	23.5
St.D.	15.6	18.6	13	15	9.9	12.4	15.2	11.9	17.2	24.6	16.2	14.7
<i>Mean monthly temperatures in °C</i>												
Max	2.9	5.8	7.8	12.6	15.8	18.2	20.8	20.9	17.7	12.6	8.6	4.4
Min	-5.6	-5.2	-0.7	6.1	9.8	14.1	16.2	14.3	11.3	6.2	-1.6	-4.7
Aver.	-1.6	0.8	4.6	8.9	13.3	16.3	18.1	17.7	14.3	9.4	4.5	0.1
St.D.	2.2	2.7	2.1	1.3	1.3	1	1.1	1.4	1.5	1.4	2.1	2.2
<i>Max. daily temperatures in °C</i>												
Max	15.4	20.9	25.6	28.1	32.4	35.5	36.8	39.2	36	29.5	23	19.2
Aver.	11.7	14.5	20.1	23.6	27.6	30.4	32.8	32.8	29.4	24.8	18.6	13.6
St.D.	2.8	3	3.1	2.3	2.2	2.5	2	2.5	2.6	2.6	2.8	3.3
<i>Min. daily temperatures in °C</i>												
Min	-27.6	-24.5	-16.5	-7.5	-4	0	1.2	2.6	-4	-7.2	-15.4	-21.7
Aver.	-15.1	-13	-8.4	-2.8	0.9	4.8	6.5	6.1	2.3	-2.5	-7.3	-12.6
St.D.	5.3	4.7	4.1	1.8	2	1.8	2.1	1.5	2.5	2.3	3.7	4.6

Source: Data from the Hydrometeorological Institute of Montenegro

The absolute maximum air temperature ever recorded was 39.2 °C. Winters are severe, with negative temperatures as low as -27.6 °C. The average annual air temperature, t_0 , was 8.9 °C. The average annual precipitation, H year, was 873 mm. The temperature coefficient for the region, T , was calculated at 0.99. The torrential rain, bb , was calculated at 84.7 mm.

The geological structure and soils of the area. The study area at large consists of various types of sediment, magmatic and metamorphic rocks generated in the long, Palaeozoic to Quaternary, interval. Most of the terrain is underlain by Mesozoic formations of carbonate composition, while magmatic and silico-clastic rocks are substantially less present. Paleozoic geological formations consist of sedimentary and metamorphic, silico-clastic rocks found mostly in the north-eastern parts of Montenegro, while Cainozoic rocks of carbonate and clastic composition occur almost in all

regions of Montenegro. The research part related to geology and soil is based on previous geological and pedological studies (Fustic and Djuretic, 2000), who analyzed all geological formations and soils of Montenegro. Furthermore, we collected some soil samples for chemical and physical analysis. The grain size composition of the soil was determined by the pipette method. The soil samples were air-dried at 105°C sifted through 2 mm sieve and dispersed using sodium pyrophosphate. Total carbonates were determined by the volumetric Scheibler method; the soil reaction (pH in H₂O and nKCl) was determined with a potentiometer; the content of the total organic matter was determined by the Kozman method; easily accessible phosphorous and potassium were determined by the Al-method and the adsorptive complex (y₁, S, T, V) was determined by the Kappen method (Spalevic et al., 2011). Certain soil analyzes in Montenegro for the presence of heavy metals were carried out as part of the "program for testing harmful substances in the soil for the year 2007". The analyzes were done by the ministry of tourism and environmental protection. The program was implemented by the public institution "Center for ecotoxicological testing of Montenegro" in accordance with the rulebook on permitted amounts of hazardous and harmful substances in soil and methods for their testing (official gazette of the Republic of Montenegro 18/97). The investigation includes analyzing land in 15 settlements that are most exposed to pollution. The total content of heavy metals in the soil was determined after preparing the samples in a closed microwave system under high pressure Milestone Ethos 1 according to US EPA method 3051A (2007).

RESULTS AND DISCUSSION

Soil is the basic substrate for plant production, and that is why it is necessary to determine whether it is suitable for this purpose, from the point of view of the content of harmful substances and heavy metals, in order to produce health-safe food (Vukeljić et al., 2002). This is possible if the determined content of heavy metals is compared with the limit value for the respective element. The limit values for the maximum allowed concentration of heavy metals in the soil (Table 1) represent significant support in the assessment of soil pollution with these predominantly toxic elements (Kastori et al., 2003; Đukić, 2003). With the sudden scientific and technological development in all branches of the economy, the emission of heavy metals increased, which thus became significant pollutants (Vukeljić et al., 2002). Plants grown on contaminated soils and useful in nutrition endanger the life and health of people and animals (Triphati et al., 2001). The concentrations of Pb, Cd, Cu and Mn in sixteen imported NPK fertilizers, which are most often used in Serbia, were determined by the method of flame atomic absorption spectrometry. The obtained results show that the content of heavy metals in samples of different fertilizers varies significantly, which depends on the ratio of N:P:K in the examined samples of fertilizers, as well as on the origin of the fertilizers themselves. The highest content of Cu was found in coloured, mixed NPK fertilizers originating from Hungary, Greece and the Netherlands, and exceeds the value of the maximum permissible concentration of Cu that can be found in the soil. The Mn content in the Hungarian NPK fertilizer (10:10:20) is ten times higher than the

average Mn content in the soil. These data indicate that it is necessary to permanently control the content of heavy metals in imported fertilizers, in order to reduce soil, underground and surface water pollution (Milinović et al., 2008). The maximum permitted amount of hazardous and harmful substances in soil (Tab. 3), in Montenegro, can lead to its pollution, and which is caused by the improper use of mineral fertilizers and plant protection products by legal and natural persons, as well as by the discharge of waste matter from various sources, are specified in the Rulebook on permitted amounts of hazardous and harmful substances in soil and methods for their testing ("Official Gazette of Montenegro", 18/97).

Table 3. Maximum allowed quantities (MDK) of dangerous and harmful substances, according to the current Ordinance

No.	Element	Chemical label	MDK in the soil in MNE, mg/kg	MDK in the soil in Serbia, mg/kg dry soil	Average value in the soil, mg/kg
1.	Cadmium	Cd	2.0	3.0	< 0.50
2.	Lead	Pb	50.0	100.0	17.37
3.	Mercury	Hg	1.5	2.0	0.0527
4.	Arsenic	As	20.0	25.0	< 5.0
5.	Nickel	Ni	50.0	50.0	17.75
6.	Fluorine	F	300.0	300.0	144.0
7.	Copper	Cu	100.0	100.0	22.5
8.	Zinc	Zn	300.0	300.0	95.0
9.	Boron	B	5.0	50	3.0
10.	Cobalt	Co	50.0	30.0	14.25
11.	Molybdenum	Mo	10.0	10.0	< 5.0
12.	Potassium	K	–	–	207.65
13.	Phosphorus	P	–	–	492.5
14.	Nitrogen	N	–	–	0.15

The research also covers the area of Bijelo Polje. The content of hazardous and harmful substances in the soil in the Municipality of Bijelo Polje was analyzed at four locations. The results of the analysis of the samples indicate the presence of lead (Pb) in concentrations above the MDK at the location of City Landfill 1. At the aforementioned location of City Landfill 1, an increased concentration of cadmium (Cd), chromium (Cr), copper (Cu), and zinc was also determined (Zn). The concentration of fluorine (F) at the Zaton 1 location is above the MDK. At the locations of City Landfill 1 and Zaton 1, there is an evident increase in lead concentration compared to the previous year, while at the other two locations, a decrease in lead concentration can be observed compared to the previous year. The content of polyaromatic hydrocarbons

exceeds MDK at the location of City Landfill 1. Except for the location Zaton 1, at all other locations, there is an increase in the concentration of polyaromatic hydrocarbons compared to the previous year.

In the final report of the Ministry, it was concluded that the condition of the soil in relation to the content of dangerous and harmful substances can be characterized as satisfactory. On the basis of the monitoring carried out, at the targeted locations, it was found that in nine municipalities, it was polluted as a result of inadequate disposal of municipal waste. In the municipality of Nikšić, inadequate disposal of industrial waste causes an increased content of dangerous and harmful substances. In most municipalities, there is a negative impact of traffic, i.e. exhaust emissions, which is a significant source of pollution of the surrounding land, at the same time endangering air quality. In relation to the study area of in the Polimlje, Montenegro region, a soil samples were analyzed, from a plot of pasture/meadow culture (Figure 1) in the Pavino Polje, municipality of Bijelo Polje, at a location of the altitude of 800 m.a.s.l., with a sampling depth of 0.3 and 0.5 m of the Ah portion of the profile. Then an analysis was carried out for the presence of heavy metals by the national and authorized laboratory at the Public institution Center for Ecotoxicological Testing – CETI in Podgorica (Montenegro). This analysis by using the method AOAC (Official, method 990.08 – Metals in solid wastes, AOAC Official methods of analysis (1995), Chapter 9, p. 31). The K content was determined on an AA-6701F Atomic Absorption Spectrophotometer, and the P content was determined on the Sequential Plasma Spectrometer ICPS-7500. Both instruments are products of Shimadzu. The results of the tests in CETI-Podgorica are given in Tab.3. The majority of microelements belong to the group of heavy metals, which are characteristically toxic to plants, animals and humans in larger quantities quote from Brankov et al. (2006).

Maximum permitted amounts (MDK) of dangerous and harmful substances, according to the current Ordinance (Official Gazette of the Montenegro, No. 18/97), it was concluded that the concentration of cadmium (Cd) is below the permitted concentration, more precisely, that it is only 0.05 mg/kg, while 2mg/kg is otherwise permitted. The concentration of 2mg/kg in the sample indicates that the presence of lead (Pb) in the controlled soil is about 30% lower than the permitted level, mercury (Hg) is only 0.005, which is a trace presence in the sample compared to the permitted 1.5mg/kg. Arsenic (As) is recorded with around 25%, and Chromium (Cr) the same with around 25%. Nickel (Ni) 17.75 mg/kg of the permitted 50 mg/kg. Fluorine (F) is recorded with about 50% of the permitted concentration, Copper (Cu) is present with 22.5mg/kg, while its presence is permitted in the amount of 100 mg/kg. Zinc (Zn) was found in the sample in a concentration of about 30% of the allowed: Boron (Bo) was recorded with 3 mg/kg, which is 2 mg/kg less than the allowed 5 mg/kg in the soil sample. Cobalt (Co) was found in the amount of 14.25 mg/kg, and the permitted concentration is 50 mg/kg. Molybdenum (Mo) is above 5mg/kg, while the permissible concentration of molybdenum is 10mg/kg of the sampled soil. The moisture content of the sampled soil during testing was 9.35%. The nitrogen content is 0.15%, phosphorus 492.5 mg/kg, and potassium 207.65 mg/kg. Based on the results (Tab. 3.) of laboratory tests in the report of Public institution CETI – Podgorica, it was determined that the tested soil samples

corresponds to the conditions of the Rulebook on permitted concentrations of hazardous and harmful substances in soil and methods for their testing (Official Gazette of Montenegro, No 18/97).

CONCLUSION

- At the studied area of Pavino Polje, analyzes of soil samples did not determine the presence of heavy metals. As a consequence it is possible to organize organic production in this region, but it is necessary to take care of the locality itself in such a way that possible existing or potential sources of its pollution would be avoided.
- The relief, climate and soils of the North Montenegro, their existing condition and soil use would be arranged for organic agriculture, so that this area can be used for the possible production of: (1) planting of continental fruits such as raspberries, blackberries, blueberries, apples, pears and plums; (2) field crops (buckwheat, millet, potato, barley, oats, rye), and the production of (3) certain types of bulk fodder (hay, silage) for domestic animals raised in the organic farming system.
- The soil, climate and relief, with their current state and existing basic features, favour the further development of organic agricultural production, especially in the studied area and region of northern Montenegro.

Acknowledgments. The paper is funded by the Ministry of Education, Science and Technological Development of the Republic of Serbia, Grant numbers: 451-3-68 / 2022-14/200032.

REFERENCE

1. Brankov, M., Ubavić, M., Sekulić, P., Vasin, J. Sadržaj mikroelemenata i teških metala u poljoprivrednim i nepoljoprivrednim zemljištima Banata, *Ins. za rat. i povr.*, Novi Sad. 42, 2006.
2. Đukić, D., Mandić, L., Zejak, D., Spalević, V. Dynamics of Microbial Activity of the Highly Present Soil Types of Monte Negro, *Acta Agriculturae Serbica*, VIII, Čačak. 15, 27-40, 2003.
3. Fuštić B. Problemi degradacije zemljišta Crne Gore, *Acta biologica Iugoslavica-serija A: Zemljište i biljka*, 54,1.1-8, 2005.
4. Fuštić B., Spalević V. Pedološka karata Sliva rijeke Lim, *Biotehnički Institut*, Podgorica 2003.
5. Fuštić, B., Đuretić, G. Zemljišta Crne Gore, Podgorica 2000. s. 134-490. 2000.
6. Fuštić, B., Topalović, Ana, Knežev, M. Zemljišta Crne Gore kao resurs održivog razvoja. XI Kongres Društva za proučavanje zemljišta Srbije i Crne Gore, Budva, 3-18. 2005.
7. Ikanović, J., Popović, V. Organic plant production. Book. Faculty of Agriculture, Bijeljina, 230.2020.
8. JDPZ: Priručnik za ispitivanje zemljišta, knj.1 Hemijske metode ispitivanja zemljišta, Beograd, 1966.

9. Kastori, R., Sekulić, P., Petrović, N., Arsenijević-Maksimović, I. Osvrt na granične vrijednosti sadržaja teških metala u zemljištu u nas i u svijetu, Naučni instit. za rat. i povrt., Novi Sad.38, 2003.
10. Kastratovic, V., Jacimovic, Z., Bigovic, M., Djurovic, D., Krivokapic, S. Environmental Status and geochemical assessment Sediments of Lake Skadar, Montenegro. Environmental Monitoring and Assessment. 188: 449-464. 10.1007/s10661-016-5459-0. 2016.
11. Milinović J., Lukić V., Nikolić-Mandić S., Stojanović D. Concentrations of heavy metals in NPK fertilizers imported in Serbia, Pestic. phytomed. (Belgrade), 23 (2008) 195-200. 2008.
12. Nagpal, A., Katnoria, J., Sharma, D. Heavy metals pollution. Biospectra 5(2): 369-374. 2020.
13. US EPA 3051A: Microwave Assisted Acid Digestion of Sediments, Sludges, Soils, and Oils, 2007.
14. Pakzad, H. R., Pasandi, M., Yeganeh, S., Lahijani, H. Assessment of heavy metal enrichment in the offshore fine-grained sediments of the Caspian Sea. Environ.l Monitoring and Assessment, 188, 303. 2016.
15. Popović V., Sikora V., Simić D., ZivanovićLj., Ugrenović V., Filipović V., Zejak D. Effect of foliar fertilization on buckwheat productivity (*Fagorypum esculentum* Moench) in organic farming system. JournalofInstitute of PKB Agroekonomik, Beograd.20(1-2):83-92. 2014.
16. Popović V., Jovović Z., Mirecki N., Lakić Ž. Organic production trend. Ed. Filipovic, Ugrenovic. Organic production and biodiversity. Open days of biodiversity, Pančevo, 3-32. 2019.
17. Pravilnik o dozvoljenim koncentracijama opasnih i štetnih materija u zemljištu i metodama za njihovo ispitivanje (Sl.RCG br.18/97).
18. Rajović G. Agroklimatski uslovi i izrada projekt plana za organsku poljoprivredu severoistočne Crne Gore, Ekonomika, Niš, 55, 1-2, 103-114., 2009.
19. Spalevic V. Assessment of Soil Erosion Processes by Using the 'IntErO' Model: Case Study of the DubokiPotok, Montenegro. Journal of Envir. Protection and Ecology, 20(2): 657–665. 2019.
20. Spalevic V. Barovic G., Vujacic D., Curovic M., Behzadfar M., Djurovic N., Dudic B., Billi P. The Impact of Land Use Changes on Soil Erosion in the River Basin of Miocki Potok, Montenegro. Water 12: 2973. <https://doi.org/10.3390/w12112973>. 2020.
21. Spalevic, V. Impact of Land Use on Runoff and Soil Erosion in Polimlje. Ph.D. Thesis, Faculty of Agriculture, University of Belgrade, Serbia, 1–260.2011.
22. Vukeljić (Popović) V. Određivanje rezidualnih količina teških metala odabranog lokaliteta u cilju zaštite životne sredine. Magistarski rad, Novi Sad,1-85. 2002.
23. Tripathi R. M., Raghunath R., Mahapatra S., Sadasivan S.Sci. Total Environ., 277, 161-168. 2001.
24. Zakon o organskoj poljoprivredi, (Sl. list RCG, br. 49/04).
25. Zakon o organskoj proizvodnji, (Službeni list SRJ br. 28, 2000.).
26. Zakon o poljoprivrednom zemljištu, (Sl. list RCG, br. 15/92 i 50/92).
27. Zejak D. Dinamika mikrobiološke aktivnosti najzastupljenijih tipova zemljišta Polimlja, Diplomski rad, Univerzitet u Kragujevcu, Agronomski fakultet, Čačak, 1-68. 2003.
28. Zejak, D., Popovic, V., Spalevic, V., Popovic, D., Radojevic, V., Ercisli, S., Glisic, I. State and economical benefit of organic production: fields crops and fruits in the world and Montenegro. Notulae Botanicae Horti Agrobotanici Cluj-Napoca, 50(3), 12815. 2022.

**Дејан Зејак^{1*}, Велибор Спалевић^{2,3}, Вера Поповић⁴, Миле Маркоски⁵,
Бранислав Дудић^{6,7}, Абдесалам Уалали⁸, Артан Хиса⁹,
Роналдо Луиз Минкато¹⁰, Шураик Кадер¹¹**

¹ Пољопривредни факултет Универзитета у Београду, Немањина 6,
Београд, Србија

² Биотехнички факултет Универзитета Црне Горе, Подгорица, Црна Гора

³ Географија, Филозофија, Универзитет Црне Горе, Бојовића бб, Никшић, ЦГ

⁴ Институт за ратарство и повртарство, Максима Горког 30, Нови Сад, Србија

⁵ Факултет пољопривредних наука и хране Скопље, Скопље,
Северна Македонија

⁶ Факултет за менаџмент, Универзитет Коменски у Братислави,
Братислава, Словачка

⁷ Факултет за економију и инжењерски менаџмент, Универзитет Привредна
Академија, Нови Сад, Србија

⁸ Факултет наука и технике Мохамедиа, Хасан II Универзитет
у Казабланки, Мароко

⁹ Факултет за инжењерство и архитектуру, Универзитет Епока,
Тирана, Албанија

¹⁰ Институт природних наука, Федерални универзитет Алфенас, УНИФАЛ-МГ,
Минас Жераис, Бразил

¹¹ Одељење за грађевинарство, Институт за информационе технологије
Шри Ланке, Малабе, Шри Ланка

* Одговорни аутор: zejakd@gmail.com

Оригинални научни рад

**АНАЛИЗА ПРИСУСТВА ТЕШКИХ МЕТАЛА У ЗЕМЉИШТИМА
БРДСКО-ПЛАНИНСКОГ ПОДРУЧЈА БАЛКАНСКОГ ПОЛУОСТРВА
СА ПРОЦЈЕНОМ ЊЕГОВОГ ПОТЕНЦИЈАЛА ЗА УЗГОЈ ВОЋАРСКИХ
КУЛТУРА: СТУДИЈА СЛУЧАЈА СЛИВА РИЈЕКЕ ЉУБОВИЋЕ,
ПОЛИМЉЕ, ЦРНА ГОРА**

Извод

У раду је приказана анализа присуства тешких метала са процјеном његовог потенцијала за узгој воћарских култура у земљиштима сјевера Црне Горе, слива ријеке Љубовиће, која се налази у брдско-планинском подручју Балканског полуострва. Узимање узорака је показало да је доминантни тип земљишта овог подручја дистрични камбисол, што се подударило са претходним истраживањима Фустића и Ђуретића (претходног вијека), Спалевића и Зејака (овога вијека). У другој фази, свели смо даље истраживања на присуство тешких метала на подручје Павиног Поља, које је по својим физичко-географским карактеристикама репрезент овог подручја из општине Бијело Поље. Полазна хипотеза је била да је ово подручје потенцијално погодно за даљи развој воћарства, а ова истраживања и предлози су ишли ка томе да се овдје даље иде ка успостављању органске производње. Резултати анализа су на крају показали да у узорцима отворених профила, а

који су прикупљени током рада на терену, не постоји присуство тешких метала. Предметни локалитет је препоручљив од стране струке и науке за даљи рад на заснивању органске производње. Посебно је погодан за производњу воћа, као примарне гране, а препоручљиво је и гајење поврћа и крмног биља.

Кључне речи: *Тешки метали, планинска подручја, воћарство, органска производња, Полимље, Црна Гора, Балканско полуострво.*

CIP – Каталогизација у публикацији
Библиотеке Матице српске, Нови Сад

502:711.4(082)

INTERNATIONAL Eco-Conference (26 ; 2022 ; Novi Sad)

Nikola Aleksić]. – Novi Sad : Ecological Movement of Novi Sad, 2022
(Novi Sad : Red copy). – 436 str. : ilustr. ; 23 cm

Tiraž 100. – Bibliografija uz svaki rad. – Rezime na srp. jeziku uz svaki rad.
– Registar.

ISBN 978-86-83177-59-2

а) Животна средина – Заштита – Градови – Зборници

COBISS.SR-ID 74631433



ISBN 978-86-83177-59-2



9 788683 177592