

- Zwatz, B. (1987): Host resistance and pathogen virulence in powdery mildew of spring barley in Austria. In: Integrated Control of Cereal Mildews: Monitoring the Pathogen, Proc. of a Seminar in the Community Programme of Coordinated Research on Energy in Agriculture held in Freising - Weihenstephan, p. 35-37, Germany.

(Primljeno: 10.03.1998.)

(Prihvaćeno: 28.05.1998.)

## STUDY OF THE *ERYSIPHE GRAMINIS* F. SP. *HORDEI* POPULATION, OF THE BARLEY PATHOGEN, USING MOBILE NURSERIES

S. Stojanović,<sup>1,3</sup> Jovanka Stojanović,<sup>1</sup> R. Jevtić,<sup>2</sup> Vesna Grujić<sup>1</sup> and S. Gudžić<sup>3</sup>

<sup>1</sup> Agricultural Research Institute „Serbia“, Belgrade, Center for Small Grains, Kragujevac; <sup>2</sup> Institute of Field and Vegetable Crops, Novi Sad; <sup>3</sup> Faculty of Agriculture, Pristina

### Summary

Spring population of *Erysiphe graminis hordei*, of the barley pathogen, was analyzed using of mobile nurseries in the Kragujevac locality (Serbia) during the period from 1995 to 1997. There were significant differences in the course of development of this parasite between years and depending on environmental conditions during plant standing in the field.

Maximal values of coefficient of infection in 1995 were in V (27-28 of April) and 1997 in VI (8-9 of May) mobile nurseries. In 1996 powdery mildew severity was very low, because of unfavourable environmental conditions for disease development.

All M1 genes of resistance had corresponding genes of virulence in the population of the parasite. The lowest frequency had v-05, V-p and V-(1402) virulency. Frequencies of V-a8, V-a, V-a3, V-a6+V-a14, V-a7+V-k+V-(No3), V-a7+V-(LG2), V-a9+V-k, V-a9, V-a10+V-(Du2), V-a12+V-(Em2); V-a13+V-(Ru3), V-c, V-(41/145), V-(Ru2), V-k, V-nn, V-at, V-g+V-(CP), V-(La) i V-h were very high.

The mobile nurseries have the great significance in the detection of rare virulences and in the investigation of development course of the causer of barley mildew. So, in *E. graminis hordei* population it is find the virulency from ml-05 genes for the first time.

**Key words:** barley; mildew; mobile nurseries; populations; *Erysiphe graminis hordei*.

Received: March 10, 1998.

Accepted: may 28, 1998.

UDK 632.9

YU ISSN 0372-7866

INSTITUT ZA ZAŠTITU BILJA I ŽIVOTNU SREDINU - BEOGRAD  
INSTITUTE FOR PLANT PROTECTION AND ENVIRONMENT - BELGRADE

# ZAŠTITA BILJA PLANT PROTECTION

VOL. 48 (3), No 221, 1997.

---

Zaštita bilja

vol. 48 (3)

Br. 221 (str. 141-204)

Beograd, 1997.

Schaud, N.W., White, W.C. (1974): A Selective Medium for Soil Isolation and Enumeration of *Xanthomonas campestris*. *Phytopathology* 64: 876-880.

Strider, D.L. (1980): Control of Bacterial Leaf Spot of Zinnia with Captan. *Plant Disease* 64: 920-922.

(Primljeno: 3.11.1997.)

(Prihvaćeno: 28.05.1998.)

## CHARACTERISTICS OF XANTHOMONAS CAMPESTRIS PV. CAMPESTRIS ISOLATES ORIGINATING FROM CABBAGE PLANTS

Olivera Jovanović, D. Antonijević and Dragana Jakovljević

Faculty of Agriculture, Belgrade - Zemun

### Summary

From the cabbage diseased leaves many isolates of the bacterium with yellow colonies were isolated. In this paper four of them were investigated.

Pathogenicity of the strains studied on cabbage plants and on other crucifers was proved by artificial inoculation. The bacteria formed yellow colonies on NA and YDC medium; oxidase test was negative, HR on tobacco leaves and catalase,  $\text{NH}_3$ ,  $\text{H}_2\text{S}$  and indol production were positive; they hydrolysed starch and aesculin and caused milk proteolysis; acid produced from arabinose, glucose, mannose and mannitol; gelatin liquification, nitrate reduction, reducing substances from sucrose, tetrasolium chloride at 0,1% tolerance, acid from dulcitol and lecithinase activity tests were negative; glucose metabolism was oxidative; they grow at 35°C and in liquid medium containing 2 and 5% NaCl.

On the basis of pathogenicity, morphological, cultural and biochemical characteristics it was shown that investigated isolated originating from diseased leaves of cabbage plants belong to the *Xanthomonas campestris* pv. *campestris* (Pammel) Dowson bacterium.

*Key words:* *Xanthomonas campestris* pv. *campestris*; bacterium; isolates; pathogenicity; morphological and cultural properties; biochemical characteristics; cabbage.

Received: November 3, 1997.

Accepted: may 28, 1998.

## TAXONOMY AND NOMENCLATURE OF PLANT PATHOGENIC BACTERIA (1864-1995)

M. Arsenijević

Faculty of Agriculture, Novi Sad and Institute for Plant Protection and Environment, Belgrade

### Summary

Considerable changings in contemporary nomenclature and taxonomy of phytopathogenic bacteria were made regarding names of some pathogens and new genera. In this paper the literature review of so far used names of bacteria and proposals for new names brought with the latest changings communicated by eminent phytobacteriologists (Young et al, 1996) were given.

- The former genus „*Corynebacterium*” had been already divided into several genera: *Arthrobacter*, *Clavibacter*, *Curtobacterium*, *Rathayibacter* and *Rhodococcus*.

- From the genus *Pseudomonas* two new genera were separated: *Acidovorax* and *Burkholderia*, the names not used so far in phytobacteriological literature. The genus *Xanthomonas* was also a subject of considerable changes, because many bacteria - pathogenic varieties of the species *X. campestris* - were renamed and they do not belong any more to this species, but to the species *X. arboricola*, *X. axonopodis* and *X. hortorum*.

- The bacterium *X. translucens* has now its pathovars: *X. translucens* pv. *arrenatheri*, pv. *cerealis*, pv. *graminis*, pv. *phlei*, pv. *phleipratensis*, pv. *poae*, pv. *secalis*, pv. *translucens* and pv. *undulosa*.

- Besides the mentioned genera, the naming of the new genus *Pantoea* for the bacteria *Pantoea agglomerans*, *P. ananas* and *P. stewartii* is suggested.

- We should be informed with all this, accept eventual new suggestions and changings and apply this in our scientific and teachers' work.

*Key words:* bacterium; genus; species; subspecies; pathovar; taxon; name of bacterium.

*Received: december 29, 1997.*

*Accepted: may 28, 1998.*

Plant Protection, Vol. 48(3), No 221: 145-164, 1997, Belgrade

## CONTENTS

### Reviews papers

- M. Arsenijević*  
Taxonomy and Nomenclature of plant Pathogenic Bacteria (1864-1995) ..... 163

### Original scientific papers

- S. Stojanović, J. Stojanović, R. Jevtić, V. Grujić and S. Gudžić*  
Study of the *Erysiphe graminis* f. sp. *hordei* population, of the barley  
pathogen, using mobile nurseries ..... 174
- O. Jovanović, D. Antonijević and D. Jakovljević*  
Characteristics of *Xanthomonas campestris* pv. *campestris* isolates  
originating from cabbage plants ..... 181
- M. Mijatović*  
Occurrence of cauliflower mosaic virus in Serbia ..... 187-188
- S. D. Stojanović*  
Incidence of mycoparasite *Colletotrichum gloeosporoides* and  
its effect on pathogen *Polystigma rubrum* subsp. *rubrum* development ..... 202