

CULTIVOS  
HERBACEOS

# 17<sup>th</sup> INTERNATIONAL SUNFLOWER CONFERENCE

Vol. 1



*Consejería de Agricultura y Pesca*



Proceedings of the  
**17th International  
Sunflower Conference**

**Vol. 1**



**Córdoba, Spain**  
June 8-12, 2008

Sponsored by The International Sunflower Association, Paris, France



Instituto de Investigación y Formación Agraria y Pesquera  
CONSEJERÍA DE AGRICULTURA Y PESCA



International Sunflower  
Association

**Título:** Proc. 17th International Sunflower Conference, Córdoba, Spain (2008), Vol. 1

© **Edita:** JUNTA DE ANDALUCÍA - Consejería de Agricultura y Pesca

**Publica:** Secretaría General Técnica. Servicio de Publicaciones y Divulgación

© **Textos:** Autores

**Colección:** Congresos y Jornadas

**Serie:** Cultivos Herbáceos

**Depósito Legal:** SE-3054-08

**Impresión:** German López - Servicios Gráficos

*Proceedings of the 17<sup>th</sup> International Sunflower Conference*  
Córdoba, Spain. June 8-12, 2008

**Editor:** Leonardo Velasco

**Editorial Committee:**

José M. Fernández-Martínez

Luis García-Torres

Pedro González

José M. Melero-Vara

Francisco Orgaz

Begoña Pérez-Vich

Francisco Villalobos

**Organizing Committee:**

Chairman: Juan Domínguez: Institute for Research and Training in Agriculture and Fisheries (IFAPA).

Deputy Chairman: Juan Fernández Pérez LIMAGRAIN IBERICA

ISA Secretariat: Laurencine Lot

ISA Secretary General: André Pouzet

Luis Carlos Alonso Arnedo. KOIPESOL S.A.

José M. Fernández Martínez. Institute for Sustainable Agriculture (CSIC)

José Antonio García de Tejada. ARLESA Semillas

José Rafael García Ruiz. Institute for Research and Training in Agriculture and Fisheries (IFAPA)

Luis López Bellido. University of Cordoba

José M. Melero Vara. Institute for Sustainable Agriculture (CSIC)

Leonardo Velasco. Institute for Sustainable Agriculture (CSIC)

*Photography: Spanish landrace of confectionary sunflower collected by L. Velasco and B. Pérez-Vich in Villarta de San Juan, Ciudad Real, Spain, on October 10, 2007.*



## Foreword

The proceedings of the 17<sup>th</sup> *International Sunflower Conference* contain 142 contributions from scientists of 24 countries. They include plenary lectures in several disciplines and regular communications presented in posters during the conference and discussed in the corresponding workshops. The manuscripts are classified by disciplines. They offer a good picture of the current state of the art of sunflower research and cultivation around the world.

The manuscripts in the *Proceedings* have been reviewed by an editorial committee with the main objective of helping the authors to improve their manuscripts through a critical reading. The authors received the edited manuscripts together with the comments of the reviewers and then went on to draft their final version. All the manuscripts received have been published in the *Proceedings*. The contents of the manuscripts are the responsibility of the authors. They should be considered as being privileged communications that require the express consent of the authors to be reprinted in part or as a whole. We wish to thank both the members of the Editorial Committee for their dedication to the task of editing such a large number of manuscripts, as well as all the authors for their collaboration throughout the whole edition process.

The Organizing Committee would also like to thank Diana Badder and José A. Palacios for their excellent editorial assistance in the preparation of these *Proceedings*. We are indebted to the Spanish Association of Sunflower Breeders (Asociación Española de Mejoradores de Girasol), which collaborated actively in the organization of the conference, and, very especially, to Juan Parejo, who was in charge of the financial side.

Finally, we would like to thank all the participants in the conference, who have contributed to its success by a careful preparation and revision of manuscripts and posters, presentation of their research in the workshops, and stimulating discussions throughout the conference on the scientific and technical aspects of sunflower research and cultivation in the world.

The Organizing Committee  
17<sup>th</sup> International Sunflower Conference  
Córdoba, Spain. June 8-12, 2008

# Volume 1

## Table of Contents

### PLENARY

|   |    |
|---|----|
| <b>Research progress in sunflower diseases and their management</b><br>Ferenc Virányi.....  | 1  |
| <b>Could a crop model be useful for improving sunflower crop management?</b><br>Francis Flénet, Philippe Debaeke, Pierre Casadebaig.....                | 13 |
| <b>Phenotypic plasticities of yield, phenological development and seed traits</b><br>Victor Sadras, Abelardo de la Vega.....                            | 19 |
| <b>Sunflower germplasm development utilizing wild <i>Helianthus</i> species</b><br>Chao-Chien Jan, Gerald J. Seiler, Thomas J. Gulya, Jiuhuan Feng..... | 29 |
| <b>Current advances in sunflower oil applications</b><br>Rafael Garcés, Joaquín J. Salas-Liñan, Mónica Venegas-Calderón, Enrique Martínez-Force.....    | 45 |
| <b>Sunflower in Spain: Past and present trends in an international context</b><br>Luis Carlos Alonso.....   | 53 |

### DISEASE RESISTANCE AND PATHOLOGY

|  |     |
|--|-----|
| <b>Phomopsis control in sunflower using products of biogenic origin</b><br>I.I. Begunov, V.T. Piven, A.T. Podvarko, V.Y. Ismailov, T. Gulya.....   | 69  |
| <b>Verticilosis en germoplasma de girasol</b><br>Julio González, Nora Mancuso, Pedro Ludueña, Antonio Ivancovich.....  | 73  |
| <b>Towards <i>Sclerotinia</i> resistance – <i>In vitro</i> screening of wild sunflower species</b><br>Ksenija T. Ajdukovic, Dragana Miladinovic, Nevena Nagl, Sreten Terzic, Sinisa Jocić, Vladimir Miklič .....   | 77  |
| <b>Correlation between macronutrient content and sunflower resistance to <i>Sclerotinia sclerotiorum</i> measured by sclerotia infection of stem</b><br>Dragana Miladinovic, Slobodanka Pajevic, Ana Marjanovic-Jeromela, Petar Sekulic, Sinisa Jocić, Vladimir Miklič ..... | 81  |
| <b>Races of <i>Plasmopara halstedii</i> on sunflower in separate agroecosystems of Adigeya Republic, Krasnodar and Rostov regions in Russia</b><br>Tatiana Antonova, Maria Iwebor, Nina Araslanova.....  | 85  |
| <b>Differences in some DNA RAPD-loci of <i>Plasmopara halstedii</i> races affecting sunflower in Krasnodar region of Russia</b><br>Tatiana Antonova, Saida Guchetl, Maria Iwebor, Tatiana Tchelustnikova.....  | 91  |
| <b>Relations between spring rainfall and infection of sunflower by <i>Plasmopara halstedii</i> (downy mildew)</b><br>Denis Tourvieille de Labrouhe, Pascal Walser, Frédéric Serre, Sylvie Roche, Félicity Vear.....  | 97  |
| <b>Structural aspects regarding formation and emission of <i>Diaporthe (Phomopsis) helianthi</i> ascospores</b><br>Valentina Androsova, Irina Balakhnina, Thomas Gulya.....  | 103 |



|   |     |
|---|-----|
| <b>Research on a growth chamber test to measure quantitative resistance to sunflower downy mildew</b><br>Frédéric Serre, Pascal Walser, Sylvie Roche, Félicity Vear, Denis Tourvieille de Labrouhe.....   | 109 |
| <b>Can management of <i>Pl</i> genes influence aggressiveness in <i>Plasmopara halstedii</i> (sunflower downy mildew)?</b><br>Nachaat Sakr, Jeanne Tourvieille, Pascal Walser, Félicity Vear, Mireille Ducher, Denis Tourvieille de Labrouhe.....                 | 115 |
| <b>Morphological and molecular identification of <i>Diaporthe helianthi</i> from <i>Xanthium italicum</i></b><br>Karolina Vrandecic, Drazenka Jurkovic, Jasenka Cosic, Luca Riccioni, Tomislav Duvnjak  | 121 |
| <b>Origins of major genes for downy mildew resistance in sunflower</b><br>Felicity Vear, Hervé Serieys, Aurélie Petit, Frédéric Serre, Jean-Pierre Boudon, Sylvie Roche, Pascal Walser, Denis Tourvieille de Labrouhe.....  | 125 |
| <b>Caracterización de la resistencia genética a podredumbre basal en girasol causada por <i>Sclerotinia sclerotiorum</i> en Argentina</b><br>Daniel Alvarez, Francisco Marraro.....   | 131 |
| <b>Molecular characterization of a novel <i>Sunflower chlorotic mottle virus</i> (SuCMoV) strain</b><br>Nicolás Bejerman, Fabián Giolitti, Sergio Lenardon.....   | 137 |
| <b>Estimation of sunflower breeding material tolerance to <i>Diaporthe/Phomopsis helianthi</i></b><br>Tomislav Duvnjak, Anto Mijić, Ivica Liović, Marija Vratarić, Aleksandra Sudarić, Miroslav Krizmanić, Karolina Vrandečić, Jasenka Čosić.....                 | 143 |
| <b>Study of the expression of shikimate dehydrogenase activity in sunflower genotypes treated with <i>Sclerotinia sclerotiorum</i></b><br>Sattar Tahmasebi Enferadi, Gian Paolo Vannozzi, Zohreh Rabiei, Ghasem Abbas Akbari..                                    | 149 |
| <b>Molecular changes in downy mildew-infected sunflower triggered by resistance inducers</b><br>Katalin Kőrösi, Ferenc Virányi.....   | 157 |
| <b>Effectiveness of the genetic resistance to <i>Plasmopara halstedii</i> under natural conditions and diversity of the pathogen within sunflower fields</b><br>Javier García-López, Rafael García-Ruiz, José Martínez-Aguilar, M. Leire Molinero-Ruiz            | 163 |
| <b>Determining the sunflower downy mildew risk by soil analysis</b><br>Denis Tourvieille de Labrouhe, Annette Penaud, Pascal Walser, Emmanuelle Mestries, Jacques Moinard, Frédéric Serre, Claire Thiery, Bernard Garric, François Delmotte, Xavier Pinochet..... | 169 |
| <b>Large scale field evaluations for <i>Sclerotinia</i> stalk rot resistance in cultivated sunflower</b><br>Thomas Gulya, Scott Radi, Nikolay Balbyshev.....  | 175 |
| <b>Study on an <i>in vitro</i> screening test for resistance to <i>Sclerotinia sclerotiorum</i> in sunflower</b><br>Miglena Drumeva, Nina Nenova, Ivan Kiryakov.....  | 181 |
| <b>EST-derived markers highlight genetic relationships among <i>Plasmopara halstedii</i> French races</b><br>François Delmotte, Xavier Giresse, Silvie Richard-Cervera, Felicity Vear, Jeanne Tourvieille, Pascal Walser, Denis Tourvieille de Labrouhe.....      | 187 |

|   |     |
|---|-----|
| <b>Effect of sowing date and initial inoculum of <i>Alternaria helianthi</i> on sunflower in the south region of Brazil</b>   |     |
| Regina M.V.B.C. Leite, Lilian Amorim, A. Bergamin Filho, Maria Cristina N. de Oliveira, César de Castro.....  | 193 |
| <b>Effects of nitrogen and water on premature ripening caused by <i>Phoma macdonaldii</i>, a fungal pathogen of sunflower</b>   |     |
| Célia Seassau, Emmanuelle Mestries, Philippe Debaeke, Grégory Dechamp-Guillaume....   | 199 |
| <b>Pathological and morphological evaluation of sunflower isohybrids carrying or not the <i>Rcm-1</i> gene for <i>Sunflower chlorotic mottle virus</i> resistance</b>                       |     |
| Maria Eugenia Bazzalo, Fabián Giolitti, María Teresa Galella, Alberto León, Sergio Lenardon.....  | 205 |
| <b>Study of resistance to <i>Sclerotinia</i> head disease in sunflower genotypes</b>  |     |
| Sándor Páricsi, Melinda Tar, Rozália Nagyné-Kutni.....  | 211 |
| <b><i>Puccinia helianthi</i> Schw., infecciones en híbridos comerciales en Argentina y su evolución durante dos décadas</b>   |     |
| N. Huguet, J. Pérez Fernandez, F. Quiroz.....   | 215 |
| <b>Results regarding the influence of <i>in vitro</i> stress induced by the <i>Phomopsis helianthi</i> filtrate on some physiological indices and on sunflower oil quantity and quality</b> |     |
| Florentina Raducanu, Elena Petcu, Constantin Raducanu, Maria Stanciu, Danil Stanciu....   | 219 |
| <b>The impact of the new races of broomrape (<i>Orobanche cumana</i> Wallr.) parasite in sunflower crop in Romania</b>  |     |
| Maria Pacureanu Joita, Steluta Raranciuc, Emilia Procopovici, Elisabeta Sava, Dumitru Nastase.....  | 225 |
| <b>Distribution and dissemination of sunflower broomrape (<i>Orobanche cumana</i> Wallr) race F in Southern Spain</b>   |     |
| Juan Fernández-Escobar, M. Isabel Rodríguez-Ojeda, Luis Carlos Alonso.....  | 231 |
| <br><b>ENTOMOLOGY</b>   |     |
| <b>Development of resistance to insect pests attacking the stem and head of cultivated sunflower in the central and northern production areas of North America</b>                          |     |
| Laurence D. Charlet, Robert M. Aiken, Gerald J. Seiler, Jerry F. Miller, Kathleen A. Grady, Janet J. Knodel.....  | 237 |
| <b>Integrated pest management of the banded sunflower moth in cultivated sunflower in North Dakota</b>  |     |
| Janet J. Knodel, Laurence D. Charlet.....   | 243 |
| <br><b>CROP PRODUCTION – MANAGEMENT</b>   |     |
| <b>Epicuticular wax content in the pericarp of sunflower fruits (<i>Helianthus annuus</i> L.) grown under moderate water deficit</b>  |     |
| María Clara Franchini, Lilia I. Lindström, Luis F. Hernández .....  | 249 |



|   |     |
|---|-----|
| <b>The pattern of foraging paths of the Honey bee (<i>Apis mellifera</i> L.) can also explain the appearance of located regions with incompletely developed fruits in the sunflower capitulum</b> |     |
| Luis F. Hernández.....  | 255 |
| <b>The influence of weather conditions on economic characteristics of sunflower hybrids in macro experiments from 1997 to 2007</b>  |     |
| Branimir Simic, Jasenka Cosic, Ivica Liovic, Miroslav Krizmanic, Jelena Postic.....   | 261 |
| <b>The appropriate technique for collecting and measuring the amount of floral nectar in sunflower (<i>Helianthus annuus</i> L.)</b>  |     |
| Zvonimir Sakac, Sreten Terzic, Vladimir Miklič .....  | 265 |
| <b>Sunflower and peanut emergence: initial development under sugarcane mulch</b>  |     |
| Nilza Patrícia Ramos, Maria do Carmo de Salvo Soares Novo, Maria Regina Gonçalves Ungaro, Antônio Augusto Lago.....   | 269 |
| <b>La agricultura de conservación como sistema viable para combatir el jopo en el girasol</b>   |     |
| J. R. García-Ruiz, F. Perea-Torres, R. Ordóñez-Fernández, J. García-López.....  | 275 |
| <b>Changes in seed oil content of sunflower (<i>Helianthus annuus</i> L.) as affected by harvesting date</b>  |     |
| Vladimir Miklič, Siniša Jocić, Dragana Vasić, Nenad Dušanić, Nada Hladni.....   | 281 |
| <b>Sunflower yield and root system development under water stress in tropical conditions</b>  |     |
| Evandro M. Gomes, Maria Regina G. Ungaro, Dirceu B. Vieira.....   | 287 |
| <b>Insecticide residues cross-contamination of oilseeds during storage</b>  |     |
| Sylvie Dauguet, Jean-Philippe Loison.....   | 293 |
| <b>Evaluating irrigation performance of sunflower in an irrigation scheme of Southern Spain</b>   |     |
| Franco Castillo-Llanque, Cristina Santos, Ignacio J. Lorite, Margarita García-Vila, Masahiro Tasumi, Javier Estévez, Pedro Gavilán.....   | 299 |
| <b>Size reduction of ornamental sunflowers by the application of daminozide</b>   |     |
| Maria Carolina Sabbagh, Francine Lorena Cuquel, Ana Cláudia Barneche de Oliveira, Edson Perez Guerra.....   | 305 |
| <b>Participatory on-farm sunflower variety evaluation in northern and eastern Uganda</b>  |     |
| Walter O. Anyanga, Moses Biruma.....  | 309 |
| <b>Initial growth of sunflower in soils with high concentrations of boron and heavy metals</b>  |     |
| Adriana M. Moreno Pires, Cleide A. Abreu, Aline Renee Coscione, Vinicius Alberti da Silva, Nilza Patrícia Ramos.....  | 315 |
| <b>Impact des facteurs limitants du rendement du Tournesol (<i>Helianthus annuus</i> L.) en conditions réelles d'utilisation par les agriculteurs, en Midi-Pyrénées – Etude de cas</b>            |     |
| V. Thebaud, J.D. Scheiner, J. Dayde.....  | 319 |

|  |     |
|--|-----|
| <b>Planting date effect on yield and yield components of sunflower in Miyaneh region</b>   |     |
| Ali Faramarzi, Mohammad-Bagher Khorshidi.....  | 325 |
| <b>Some aspects of sunflower crop management in Romania</b>  |     |
| Gheorghe Sin, Marius Botea, Lenuța Drăgan.....   | 329 |
| <b>Screening and drying conditions for early harvested sunflower</b>   |     |
| Genta Kanai, Hitoshi Kato, Naonobu Umeda, Kensuke Okada, Morio Matsuzaki.....  | 333 |
| <br><b>CROP PRODUCTION – PHYSIOLOGY</b>  |     |
| <br><b>Physiological maturity in sunflower. Correspondence between the quantitative and the visual definition</b>                                    |     |
| Luis F. Hernández, Adelina O. Larsen, Lilia I. Lindström, Liliana B. Iriarte.....  | 337 |
| <b>Influence of desiccation on germination and field emergence of sunflower</b>  |     |
| Ivica Liović, Marijan Bilandžić, Miroslav Krizmanić, Anto Mijić, Ruža Popović, Ilonka Ivanišić, Tomislav Duvnjak, Branimir Šimić, Jasenka Čosić..... | 341 |
| <b>Physiological traits for quantification of drought tolerance in sunflower</b>   |     |
| Elena Petcu, Maria Stanciu, Danil Stanciu, Florentina Raducanu.....  | 345 |
| <b>El vuelco en el cultivo de girasol: características anatómicas y mecánicas del sistema radical</b>  |     |
| Milena E. Manzur, Antonio J. Hall, Claudio A. Chimenti.....  | 351 |
| <b>Influence of drought stress on growth, protein expression and osmolyte accumulation in sunflower <i>Helianthus annuus</i> L. c.v. Peredovik</b>   |     |
| Sabine Fulda, Hendrikje Stegmann, Renate Horn.....   | 357 |
| <b>Development and validation of a model of lodging for sunflower</b>  |     |
| Mariano M. Sposaro, Pete M. Berry, Mark Sterling, Antonio J. Hall, Claudio A. Chimenti.....  | 363 |
| <b>Exploring genotypic strategies for sunflower drought resistance by means of a dynamic crop simulation model</b>                                   |     |
| Pierre Casadebaig, Philippe Debaeke.....   | 369 |
| <b>Dynamics of dry matter accumulation in sunflower</b>  |     |
| Nenad Dusanic, Vladimir Miklič, Igor Balalic, Velimir Radic, Jovan Crnobarac.....  | 375 |
| <b>IAA/GA3 quantitative ratio of some sunflower genotypes representing CMS-Rf system</b>   |     |
| Maria Duca, Angela Port.....   | 381 |
| <b>Genetic-phytohormonal interactions in male fertility and male sterility phenotype expression in sunflower (<i>Helianthus annuus</i> L.)</b>       |     |
| Maria Duca, Angela Port.....   | 387 |
| <b>Direct and indirect effects of morphophysiological traits on seed yield of sunflower (<i>Helianthus annuus</i> L.)</b>                            |     |
| Nada Hladni, Siniša Jocić, Vladimir Miklič, Anto Mijić, Dejana Saftić Panković.....  | 393 |
| <b>Determination of maximum achene size in sunflower</b>   |     |
| A. Mantese, D. Rondanini, D. Medan, A.J. Hall.....   | 399 |
| <b>Abscisic acid content of a nondormant sunflower (<i>Helianthus annuus</i> L.) mutant</b>  |     |
| Brady A. Vick, C.C. Jan.....   | 405 |
| <b>Producción de girasol (<i>Helianthus annuus</i> L.) en valles altos de México</b>   |     |
| J. Alberto Escalante-Estrada, M <sup>a</sup> Teresa Rodríguez-González.....  | 411 |



## The appropriate technique for collecting and measuring the amount of floral nectar in sunflower (*Helianthus annuus* L.)

Zvonimir Sakac, Sreten Terzic, Vladimir Miklic

Institute of field and vegetable crops, Maksima Gorkog 30, 21000 Novi Sad, Serbia  
E-mail: sakacz@ifvcns.ns.ac.yu; terzic@ifvcns.ns.ac.yu; miklicv@ifvcns.ns.ac.yu

### ABSTRACT

The available techniques for collecting and measuring the amount of floral nectar are applicable but often found to be unrepresentative. Centrifugation yields larger samples but they also include nectar that is not actually accessible to insects, the capillary method has been described as unsuitable because of possible damage to the nectary tissue, the method including filter paper is considered to be unreliable because of evaporation and nectar extraction methods including washing are considered limited because the solution may include sugars from plant tissue cells. We have found that capillary tubes with inner diameter of 0.25-0.5mm and outer diameter of 0.5-0.75mm are suitable for nectar collection in sunflower. To determine the amount of nectar, we isolate five inflorescences per sunflower line at the start of flowering and collect the nectar two days after the isolation. The capillary tube is inserted between the style and filaments down to the nectary. After the level of nectar stops rising the next flower is processed. The tubes can be measured on an analytical scale and the amount of nectar is obtained as the weight increase in comparison to the empty tube. Faster determination of floral nectar amount can be provided by using calibrated capillary tubes of a known and uniform inner diameter. The appropriate outer diameter of the capillary tubes reduces the risk of tissue damage and allows more precise collecting so that the capillary method is preferable to others for nectar collecting in sunflower.

**Key words:** capillary technique – nectar quantity – sunflower

### INTRODUCTION

Sunflower is one of the plant species that produces pollen which is too heavy for wind dispersal (Putt, 1940). Even though the cultivated sunflower has a reasonable percentage of self-compatibility it still benefits from insect pollination. One of the major components influencing pollinator choice is certainly the production of nectar, whose amount and quality are often studied.

The nectaries in *Asteraceae* family form on top of the ovary and surround the style base (Mani and Saravanan, 1999), (Fig. 1). The nectar can be accessed for quantification purposes by capillary tubes (Hocking, 1953), volumetric centrifugation (Bosi, 1973), filter paper strips (McKenna and Thomson, 1988), flushing of water into the corolla (Cresswell and Galen, 1991) or floating the flowers inverted in water (Manetas, 2000) depending on flower structure. These techniques were used with a variable success, but the overall conclusion is that no single method can be considered satisfactory for all plant species (Mesquida et al, 1988). The capillary method can be used in sunflower but it is necessary to use capillary tubes of appropriate dimensions. If the outer diameter is too large it is not possible to access the nectar without destroying the surrounding corolla tissue (Fig. 1.) and if the tube is too thin then the intake of nectar is slower and the strong capillary force may make the extraction of nectar from the tubes difficult.




Fig. 1. Longitudinal section of a sunflower disk flower showing the location of the nectary

#### MATERIALS AND METHODS

We found that capillary tubes with inner diameter of 0.25-0.5mm and outer diameter of 0.5-0.75mm are suitable for nectar collection. The length of the tube then determines the capacity and can be picked for its suitability to the collecting design but should not be smaller than 15  $\mu$ l.

To determine the amount of nectar, we isolate five inflorescences per sunflower line at the start of flowering with linen bags to prevent insects from collecting nectar. The best moment is when the first two rows of disk flowers have opened. Two days after the isolation, at approximately 8 AM, the inflorescences are cut and taken to the laboratory in a portable refrigerator to minimize evaporation and the change in nectar volume. It is advisable for the transport duration to be as short as possible. The following should be prepared to analyze one inflorescence:

1. Four capillary tubes (previously weighed on an analytical scale) each placed in a separate tube labeled with sample and replication to ease the work of collecting and measuring
2. A clean vial (previously weighed on an analytical scale)
3. A clean HPLC vial with sample label on it filled with 1 ml mixture of AcCN:H<sub>2</sub>O in a ratio of 75:25
4. A plastic dish with sample label on it for deep freezing

Four groups of five analyzed flowers are equally far from each other on an inflorescence. We collect nectar from 5 fully opened disk flowers with one non-calibrated capillary tube. The tube is inserted between the style and filaments down to the nectary (Fig. 1.). After the level of nectar stops rising the next flower is processed. When a total of 20 flowers are finished, the tubes can be measured on an analytical scale and the amount of nectar is obtained as the weight increase in comparison to the empty tube. Faster determination of nectar amount in flowers can be provided by using calibrated tubes (capillary tubes with a uniform known inner diameter) for nectar collecting, in which case the height of nectar in tubes can be correlated with the nectar volume. This method is suitable when it is necessary to determine the amount of nectar in field conditions, without cutting the sunflower head and taking it to the lab.

The next step in method developing is a qualitative and quantitative HPLC analysis of nectar extracted from a single inflorescence as a collection from 20 disk flowers. For this purpose, ten disk flowers are also pulled off with tweezers, put in a glass and weighed on an analytical scale to obtain the information about the flower mass and possible correlation with nectar production.

The nectar collected in capillary tubes, after weight measurement, can also be kept for subsequent analysis. The contents of all capillary tubes from a single inflorescence are transferred into a HPLC vial (2 ml) filled with 1 ml mixture of AcCN:H<sub>2</sub>O in a ratio of 75:25 and placed in a refrigerator for



subsequent HPLC nectar quality analysis. Twenty flowers are pulled out of the disk with tweezers, frozen in liquid nitrogen and then kept at  $-72^{\circ}\text{C}$ .

### DISCUSSION

The rest of the techniques cited are applicable but often unrepresentative. Centrifugation yields larger quantities but they also include nectar that is not actually accessible to insects and modified chemical composition due to tissue lesion (Mesquida et al., 1988). The method including filter paper is considered to be unreliable because of evaporation (Livtzieva, 1954). Nectar extraction methods including washing are considered limited because the solution may include sugars from plant tissue cells (Kenoyer, 1917).

A combination of capillary method and filter paper can be used so that the nectar is extracted with capillary tubes and then ejected on to a filter paper, which is measured for total nectar. After the evaporation has finished the amount of sugars is obtained as a difference between wet and dry filter paper.

The capillary method has been used on sunflower (Pham-Delegue et al., 1985) but it has also been described as unsuitable for collecting nectar amounts less than  $1\ \mu\text{l}$  and to cause damage to the nectary tissue (McKenna, 1988). The appropriate outer diameter of the capillary tubes reduces the risk of tissue damage and allows more precise collecting so that the capillary method is preferable to others for sunflower nectar collecting.

### REFERENCES

- Bosi, G. 1973. Méthode rapide pour la détermination par chromatographie en phase gazeuse des glucides du nectar: technique de prélèvement du nectar et de préparation des éthers triméthylsilylés en présence d'eau. *Apidologie* 4:57-64.
- Cresswell, J.E., and C. Galen. 1991. Frequency-dependent selection and adaptive surfaces for floral character combinations: the pollination of *Polemonium viscosum*. *Amer. Natur.* 138:1342-1353.
- Hocking, B. 1953. The intrinsic range and speed of flight of insects. *Trans. Royal Entomol. Soc. London* 104:223-345.
- Kenoyer, L.A. 1917. Environmental influences on nectar secretion. *Bot. Gaz.* 63:249-365.
- Livtzieva, E.K. 1954. Methods used in nectar secretion investigations. (In Russian). *Pchelovodstvo*, 31(11):33-39.
- Mani, M.S., and J.M. Saravanan. 1999. *Pollination ecology and evolution in Compositae (Asteraceae)*. Enfield, NH, USA: Science Publishers Inc.
- McKenna, M.A., and J.D. Thomson. 1988. A technique for sampling and measuring small amounts of floral nectar. *Ecology* 69:1306-1307.
- Mesquida, J., R. Marilleau, and M. Pham-Delegue. 1988. A study of rapeseed (*Brassica napus* L. var. *oleifera* Metzger) flower nectar secretions. *Apidologie* 19:307-318.
- Pham-Delegue, M.H., C. Masson, A. Bonjean, P. Etievant, and R. Marilleau. 1985. Etude des paramètres chimiques impliqués dans la pollinisation du tournesol par les abeilles en production de semences hybrides. p. 559-564. In: Proc. 11<sup>th</sup> Int. Sunflower Conf., 10-13 March, Mar Del Plata, Argentina.
- Putt, E.D. 1940. Observations on morphological characters and flowering processes in sunflower (*Helianthus annuus* L.). *Sci. Agric.* 25:185-190.